

Review

The Yeast *Torulaspora delbrueckii*: An Interesting But Difficult-To-Use Tool for Winemaking

Manuel Ramírez * and Rocío Velázquez

Departamento de Ciencias Biomédicas (Área de Microbiología), Facultad de Ciencias, Universidad de Extremadura, 06006 Badajoz, Spain; rociovelazquez1981@gmail.com

* Correspondence: mramirez@unex.es; Tel.: +34-924-289426

Received: 27 September 2018; Accepted: 9 November 2018; Published: 12 November 2018



Abstract: *Torulaspora delbrueckii* is probably the non-*Saccharomyces* yeast that is currently most used for winemaking. Multiple advantages have been claimed for it relative to conventional *S. cerevisiae* strains. However, many of these claimed advantages are based on results in different research studies that are contradictory or non-reproducible. The easiest way to explain these discrepancies is to attribute them to the possible differences in the behaviour of the different strains of this yeast that have been used in different investigations. There is much less knowledge of the physiology, genetics, and biotechnological properties of this yeast than of the conventional yeast *S. cerevisiae*. Therefore, it is possible that the different results that have been found in the literature are due to the variable or unpredictable behaviour of *T. delbrueckii*, which may depend on the environmental conditions during wine fermentation. The present review focusses on the analysis of this variable behaviour of *T. delbrueckii* in the elaboration of different wine types, with special emphasis on the latest proposals for industrial uses of this yeast.

Keywords: *Torulaspora delbrueckii*; winemaking; yeast inoculation; yeast dominance; wine quality; genetic improvement

1. Introduction

Of the non-*Saccharomyces* yeasts, *Torulaspora delbrueckii* is probably the most suitable for use in winemaking. This is because it has a good fermentation performance compared to other non-*Saccharomyces* yeasts that might be considered for winemaking, such as *Hanseniaspora uvarum*, *H. vineae*, *Candida zemplinina*, *C. pulcherrima*, *C. stellata*, *Schizosaccharomyces pombe*, *Hansenula anomala*, *Metschnikowia pulcherrima*, *Lachancea thermotolerans*, *Pichia fermentans*, *P. kluyveri*, and *Kazachstania aerobia*. Additionally, it has been claimed that *T. delbrueckii* can be used to optimise some wine parameters with respect to usual *S. cerevisiae* wines such as a low amount of acetic acid, lower ethanol concentration, increased amount of glycerol, greater mannoprotein and polysaccharide release, promoted malolactic fermentation, increased amounts of interesting aromatic compounds (fruity esters, lactones, thiols, and terpenes), and decreased amounts of unwanted aromatic compounds (such as higher alcohols); these may improve wine quality or complexity (reviewed by Benito in [1]). These additional features constitute the main reason why modern oenologists are conducting trials with this yeast, as well as with other non-*Saccharomyces* yeasts, for industrial winemaking. However, this option entails additional complications and economic costs in controlling must fermentation with respect to the use of conventional selected strains of *S. cerevisiae*, which is undoubtedly the most reliable yeast species for this purpose. Some of the complications of using *T. delbrueckii* in commercial wineries come from its physiological properties under the stressing conditions that are usual in the winemaking process. In this work, we review these specific properties, with a focus on their relevance for some applied aspects of winemaking.

2. The *Torulaspora* Yeasts

Torulaspora are fermentative yeasts that can be found in wild and anthropic habitats, where they may coincide with other fermentative yeasts such as *Saccharomyces* and *Zygosaccharomyces*. In the past, there has been some misclassification of some species in these three genera, because they share various morphological and physiological features. The genus *Torulaspora* includes at least six species: *T. delbrueckii* (anamorph *Candida colliculosa*), *T. franciscae*, *T. pretoriensis*, *T. microellipsoides*, *T. globosa*, and *T. maleeae* [2]. Other species have also been proposed for inclusion in this genus following their characterisation by molecular tools used to discriminate new isolates, examples being *T. indica* [3] and *T. quercuum* [4]. The taxonomy of *Torulaspora* is changing rapidly, and one can expect species reassignments and new species in the near future. In this sense, based on the nucleotide sequence of the internal transcribed spacer (ITS) region located between the 18S and 28S rRNA genes, it has already been proposed that *Zygosaccharomyces mrakii* and *Z. microellipsoides* might be reclassified in the genus *Torulaspora* [5]. Similarly, four strains presumed to be *T. delbrueckii* were found to be considerably different from the type strain, and were reclassified into the genera *Debaryomyces* and *Saccharomyces* [6]. In particular, one must bear in mind that some error in the presumptive assignment of a new isolate to *Torulaspora* may explain some of the controversial results found for the biotechnological properties that have been recently claimed for these yeasts (see Section 5 below).

Torulaspora cells mostly have a spherical shape (*torulu*), although ovoid and ellipsoidal shapes are also frequent. Its cellular size, $2\text{--}6 \times 3\text{--}7 \mu\text{m}$, uses to be smaller than that of *S. cerevisiae* (Figure 1A,E). *Torulaspora* may rarely produce pseudohyphae, but never true hyphae. Its life cycle has yet to be elucidated. In contrast with wild *S. cerevisiae* strains, which are diploids or polyploids during vegetative propagation, *T. delbrueckii* was long believed to be a haploid yeast, mainly because of its small cell size [7], and because tetrads are absent or rarely observed in sporulation media. However, it has recently been suggested that *T. delbrueckii* may actually be mostly diploid and homothallic [8]. If this is the case, the small cell size of *Torulaspora* may still be explained by its having only 16 chromosomes in the diploid stage, instead of the 32 chromosomes that are present in *S. cerevisiae* diploid yeasts. It is believed that *T. delbrueckii* diverged from the *S. cerevisiae* lineage before the latter underwent its genome polyploidisation [9].

All species of *Torulaspora* reproduce asexually by multilateral budding (mitosis). Sexual reproduction may occur in sporulation media through asci that contain one to four spherical ascospores (Figure 1B). The ascus can be originated by the heterogamic conjugation between a cell and its bud, or between two independent cells. Three types of conjugation have been proposed in the *Torulaspora* genus [10]: (i) between mother and daughter cells (the daughter cell may be one small bud still attached to the mother cell, but separated from it by a cross-wall); (ii) between two cells with their respective conjugation tubes; and (iii) between a cell with conjugation tube and another cell without it. The ascospores (one to four per ascus, frequently two, $2\text{--}4 \mu\text{m}$ diameter) are spherical with a smooth or warty cell wall. It should be noted that starvation is not required for *T. delbrueckii* to sporulate [8,11]; i.e., spores can be formed on rich media such as yeast extract peptone dextrose (YEPD) agar (Figure 1D), as also may occur for some wine strains of *S. cerevisiae*.

Cells with conjugation tubes are frequent in *T. delbrueckii*, mainly in non-growing cells in a stationary growth stage or in sporulation media (Figure 1B), but most of them seem uninvolved in the yeast conjugation process [12]. Probably, most of the several buds that have been frequently observed in single non-growing yeasts are actually incipient cell wall protuberances destined to potentially become a conjugation tube for yeast mating.

Given this lack of precise knowledge about the life cycle of *Torulaspora*, it is complicated to design strategies for the biotechnological improvement of *T. delbrueckii* by using classical genetic techniques such as those proposed for *S. cerevisiae* wine yeasts [13–16].




Figure 1. Vegetative cells (A,C,E) and spores (B,D,F) of *T. delbrueckii* and *S. cerevisiae* for comparison. Photomicrographs were taken using Nomarski (vegetative cells) or bright-field illumination (spores) with a Nikon Eclipse 600 microscope equipped with a 60× objective. (B,F) Spores from Minimal Sporulation Medium. (D) Spores from YEPD agar stained with malachite green.

3. Growth and Fermentation Capability

With respect to the features required to perform industrial alcoholic fermentation, of the non-*Saccharomyces* yeasts, *T. delbrueckii* is the most similar to the referent, best-considered, *S. cerevisiae*. This is the main reason why *T. delbrueckii* was the first non-*Saccharomyces* yeast proposed for industrial use in wine fermentation [17]. However, slight but relevant differences in the physiological characteristics of the two yeasts affect their choice as options for different industrial applications. In particular, the specific rates of CO₂ production and O₂ consumption are higher in *T. delbrueckii* than in *S. cerevisiae*, which results in lower biomass yields from batch cultures, and may represent a handicap for the large-scale production of *T. delbrueckii* with respect to *S. cerevisiae* [18]. The two yeasts' sugar utilisation and regulation patterns are very similar, with similar biomass yields in glucose-limited oxygen-sufficient chemostat cultures, as is consistent with fully respiratory growth. Interestingly, *S. cerevisiae* is the first to switch to a respirofermentative metabolism as the oxygen feed rate decreases, showing a lower biomass yield at low oxygen tensions with respect to *T. delbrueckii*, which is still able to sustain full respiration under these conditions [19]. Unfortunately, however, *T. delbrueckii* shows less growth than *S. cerevisiae* under strict anaerobic conditions [20,21], and has less fermentation vigour and slower growth rate than *S. cerevisiae* under usual wine fermentation conditions, being quickly overcome by wild or inoculated *S. cerevisiae* strains [18,22]. Obviously, this is an important issue for winemaking, which is usually done under strict anaerobic conditions for white and sparkling wines, or in the presence of very low amounts of oxygen for red wines, where the cap of grape skins is punched down into the must several times a day. A poorer performance of *T. delbrueckii* with respect

to *S. cerevisiae* should always be expected under these working conditions, especially for white and sparkling wines.

When inoculated in fresh grape must, the dominance ratio of *T. delbrueckii* depends on many factors: the size of the inoculum, amount and types of viable wild microbes initially present in the must, fermentation stage, sugar and ethanol concentrations, killer phenotype and killer sensitivity of the inoculated yeast and the wild yeasts, SO₂ (sulfur dioxide) concentration, Cu (copper) concentration, pesticides coming in with the grapes, etc. Some of these have been analysed in several related works. The *T. delbrueckii* dominance ratio during the must fermentation reported in the literature of specialised research works ranges from 0 to 100%, being in most cases around 50% in tumultuous fermentation [22–27]. This dominance ratio can be expected to be lower in commercial wineries, where the working conditions are not as controlled as in research laboratories or experimental wineries. This ratio is much lower than that which is expected for the reference yeast *S. cerevisiae*, which is assumed to be close to 100%, even considering that this yeast is usually inoculated at a lower concentration (around 10⁶ CFU/mL) than *T. delbrueckii* (around 10⁷ CFU/mL).

A strong dominance of *T. delbrueckii* is usually achieved by using sterile or much clarified must, and a killer strain for inoculation. In this way, the relative initial growth of the *T. delbrueckii* population is high, and favours its becoming the clear protagonist during fermentation. Nonetheless, since this yeast is less resistant than *S. cerevisiae* to high ethanol concentrations (Figure 2), the fermentation rate slows down, and cell death increases after the tumultuous fermentation of sugar-rich musts (Figure 4). As a consequence, fermentation may stop, slow down and become sluggish, or continue mainly because of the presence of some contaminating wild *Saccharomyces* yeasts, which may continue must fermentation if the required nutrients are still available. Obviously, any such sequence of events will reduce the domination ratio of *T. delbrueckii* by the end of fermentation, even down to its disappearance [22,24]. This situation is more evident and occurs more rapidly when mixed-inoculating with the two yeasts (co-inoculation or sequential inoculation) in grape must be supplemented with yeast nutrients. In this case, the dominance of *T. delbrueckii*, if it occurs at all, is usually ephemeral, lasting no more than two days [18,22,24].




Figure 2. Resistance of *T. delbrueckii* (Td) and *S. cerevisiae* (Sc) to ethanol, SO₂, and Cu. Two strains of each yeast species are shown (Td-1 and Td-2, and Sc-1 and Sc-2, respectively). YEPD: rich YEPD medium. SD: minimal SD medium.

This situation makes it difficult to reliably assign a specific feature to *T. delbrueckii*, and may explain why some disagreement is found in the literature about the effect of this yeast on winemaking and wine quality. For example, the effects related to the production of some aroma compounds remain confusing. Mixed *T. delbrueckii*/*S. cerevisiae* inoculation has been described as increasing the total ester concentration (mainly isoamyl acetate and ethyl hexanoate, octanoate, and 3-hydroxybutanoate) relative to inoculation with *S. cerevisiae* [28], while the contrary has also been reported [29,30]. Such contradictory results may occur just because the production and degradation of these compounds depend on the proportion of each species of yeast during must fermentation. Likewise, even when a putative technological improvement of *T. delbrueckii* affecting the quality of wine has been accepted, it is still very complicated to achieve the desired result in commercial wineries if the dominance of this yeast cannot be ensured, and this may dissuade oenologists from spending money purchasing new non-*Saccharomyces* yeasts for must inoculation. Consequently, it is necessary to thoroughly control the level of dominance of *T. delbrueckii* throughout the wine fermentation process, analysing adequate numbers of samples and colonies of viable microorganisms sampled at different times during that process. Frequently, one finds in the literature that few samples and yeast colonies were analysed (10 colonies, only twice per fermentation [31]), or simply that the inoculated *T. delbrueckii* strain was not monitored, but was simply assumed to dominate the wine fermentation [32]. Moreover, in some cases, the authors themselves recognise the presence of contaminating yeasts in the pasteurised must; these are yeasts that, in parallel experiments in which there was no dominance of the inoculated yeast, were able to complete the fermentation. Even more so, in some cases, the fermentations were carried out by continuous mixing of the inoculated must in an orbital shaker to increase the oxygen supply [31], contrary to what is usually done in commercial cellars where the must is not continuously agitated. Therefore, in sum, the different environmental conditions used in the experimental fermentations of *T. delbrueckii*, and the uncertainty about the desired dominance of this yeast during fermentation, suggest that the observed effect of a given strain on the quality of wine may be different, or even contradictory, depending on the working conditions. Consequently, it would be necessary to establish criteria that are more precise regarding the experimental conditions and the minimum level of dominance of *T. delbrueckii* in order to obtain more homogeneous and reliable conclusions about any of the advantages of this yeast in winemaking.

In the absence of general agreement on these criteria, one possibility that might explain some of the conflicting results found is that different *T. delbrueckii* strains might have different fermentation capabilities. This would not be too surprising, given the possible lack of a reliable and definitive taxonomic classification. However, to confirm this argument, it would be necessary to test in parallel the different yeast strains under identical experimental conditions, and achieve a clear dominance of each strain tested. Moreover, in addition to the idea that the production of esters by *T. delbrueckii* might be strain-dependent, it has been proposed that this production may even vary when this yeast is associated with *S. cerevisiae* in mixed cultures [25,33,34]. This situation may become even more complicated, although clearly very interesting, when considering the use of *T. delbrueckii* for red winemaking, since this differs from white winemaking in certain aspects that affect this yeast's growth during must fermentation, and hence its effect on wine quality. Namely, oxygen availability, alcohol content, the amount of initial wild microorganisms, and nutrient availability are usually greater in red than in white wine fermentation. Given these differences, and since fermentations dominated by *T. delbrueckii* are slower than those dominated by *S. cerevisiae*, the eventual development of lactic acid bacteria may promote malolactic fermentation occurring simultaneously with alcoholic fermentation. This promotion is enhanced in situations where these bacteria may be more frequent, as when using poorly clarified musts or crushed grapes as starting material for winemaking. Indeed, we found that in such situations all of the *T. delbrueckii* wines, but none of the *S. cerevisiae* wines, underwent malolactic fermentation, and putative lactic acid bacteria were always found in the *T. delbrueckii* wines, but none or very few were found in the *S. cerevisiae* wines. The highest malic acid degradation was found in those wines where *T. delbrueckii* reached the greatest dominance ratios, and at the same time, had the

slowest fermentation kinetics [23,24]. Likewise, some companies, such as Chr. Hansen (Hoersholm, Denmark), recommend inoculation with *T. delbrueckii* to promote malolactic fermentation, although in this case, the effect is claimed to be due to a decreased production of toxic medium-chain fatty acids that favour the growth of lactic acid bacteria.

I.E., the confluence of so many environmental factors in winemaking that may change the behaviour of *T. delbrueckii* with respect to the production of aroma compounds makes it difficult to design a reliable strategy for wine quality improvement in commercial wineries. In any case, until the handicap of low dominance of *T. delbrueckii* is resolved, one cannot determine with precision the real usefulness of these yeasts in the cellar. To address this problem, further knowledge is required of how *T. delbrueckii* interacts with other wine microorganisms during wine fermentation so as to be able to ensure the dominance of this yeast species.

4. Interaction with Other Fermentative Yeasts and Ability to Dominate Must Fermentation

As noted above, the ability of *S. cerevisiae* to overcome other non-*Saccharomyces* yeasts and dominate wine fermentation can be explained mainly by its excellent biological fitness and its tolerance to the stress factors that converge in this process. This circumstance makes it possible that an almost undetectable contamination of *S. cerevisiae* in a must that has been previously inoculated with *T. delbrueckii* can be responsible for carrying out most of the fermentation process. This may occur because *T. delbrueckii* dies prematurely, or because its metabolic activity declines markedly as a result of the environmental stress. However, the opposite case is clearly unlikely, so that non-*Saccharomyces* yeasts usually play an irrelevant part in the fermentations of musts inoculated with *S. cerevisiae*. Biotechnologically, this situation will not change until *T. delbrueckii* strains whose biological efficacy is similar to that of *S. cerevisiae* are available, which is an aspect that we shall return to later in this review. Apart from this, other types of direct interactions between the two yeast species have been described that can influence, or even determine, the preponderance of one yeast over the other.

The yeast interactions in pure and mixed fermentations have been the subject of a thorough review [35] which clearly stated that the competitiveness of any given strain can be influenced by many factors, which are both abiotic (pH, temperature, ethanol, osmotic pressure, nitrogen availability, molecular sulfur dioxide concentration, etc.) and biotic (type of microorganisms, presence of antimicrobial molecules, cell-to-cell contact, grape variety, etc.). This competitiveness will change depending on the specific environmental conditions of a given food fermentation, which will in turn influence the ability of a given strain to out-compete other yeasts. Moreover, in mixed fermentation, the choice of the specific strains of *T. delbrueckii* and *S. cerevisiae* can be another biotic factor to consider as an origin of variability in the competitiveness of each yeast strain. However, the use of a fixed couple of yeast strains could produce predictable results related to the dynamic of each yeast population and, thereby to the wine quality.

With respect to the biotic factors, a new wine *T. delbrueckii* killer strain has recently been characterised. It secretes a killer toxin (Kbarr-1) that is encoded in a double-stranded RNA virus (ScV-Mbarr-1) with broad antifungal activity against *S. cerevisiae* (killer and non-killer strains) and other non-*Saccharomyces* yeasts [36,37]. This negative interaction has already been applied to promote the dominance of *T. delbrueckii* in white [24] and red [23] table wines. This killer strain had the advantage of dominating must fermentation in the presence of *S. cerevisiae* relative to the non-killer isogenic strains. However, full dominance is not ensured if a fairly large population of *S. cerevisiae* yeasts (about 10^6 CFU/mL) is co-inoculated or was already present as wild yeasts in the grape must. Therefore, a large inoculum (about 10^7 CFU/mL) of *T. delbrueckii* should be considered to achieve satisfactory dominance in these situations.

Apart from this, yeast growth inhibition by cell-to-cell contact is another biotic factor that may influence the competitiveness of each strain in multi-starter fermentations. Physical contact between *S. cerevisiae* and *T. delbrueckii* seems to induce the rapid death of the latter yeast. However, when the two yeasts were separated from each other in a double-compartment fermenter, *T. delbrueckii*

maintained its viability and metabolic activity [38]. The details of this mechanism of cell-to-cell contact inhibition are still unknown. It has recently been proposed that in the case of the early death of *L. thermotolerans* during anaerobic fermentation mixed-inoculated with *S. cerevisiae*, this phenomenon is caused by a combined effect of cell-to-cell contact plus antimicrobial peptides [39]. Similarly, it has been found that cell-free *S. cerevisiae* supernatants induce fast culturability loss in other non-*Saccharomyces* yeasts, meaning that this inhibitory effect is mainly caused by some metabolites secreted by *S. cerevisiae*. To further complicate the issue, the culturability loss of non-*Saccharomyces* yeasts induced by *S. cerevisiae* is species-dependent and strain-dependent [40]. The effect of these biotic factors on the competitiveness of each strain can be influenced by an abiotic factor, an example being the addition of assimilable nitrogen, which was found to partially mitigate the inhibitory interaction between different yeasts [41].

Given this situation in which many different factors may influence a given yeast's competitiveness with respect to others, and the lack of precise knowledge about the mechanisms of this influence, the degree of compatibility of co-inoculated strains of *T. delbrueckii* and *S. cerevisiae* must be known with precision in order to program and control the industrial use of that combination. This is especially important if a specific proportion of the fermentation must be performed by *T. delbrueckii* in order to obtain a determined effect on the final wine's organoleptic quality. Mixed inoculation results may not be fully satisfactory if the production of aromas by *T. delbrueckii* varies when it is associated in a variable and difficult-to-control proportion with *S. cerevisiae*, or when the type and amount of the aromas produced in mixed cultures depend on a synergistic interaction between these two yeasts [33]. If this is the case, the use of *T. delbrueckii* in mixed-inoculated fermentation could be a situation that is far too complicated to control satisfactorily at an industrial level. However, if total or major dominance of *T. delbrueckii* is desired, this can be achieved using killer strains, a large yeast inoculum (approximately 10^7 CFU/mL), and grape must containing only a discrete amount of wild *Saccharomyces* yeasts (less than 10^5 CFU/mL) [23,24,42]. In this situation, if the effect of *T. delbrueckii* on the quality of the wine is more intense than desired, this could be resolved by mixing with conventional *S. cerevisiae* wine to obtain the required wine quality.

5. Influence on the Chemical Composition and Aroma of Table Wine

As mentioned in the Introduction section, several wine parameters can be improved by inoculating the must with *T. delbrueckii*. However, some of these claimed improvements can be achieved only in certain specific circumstances. For example, the production of reduced amounts of acetic acid is usually found only in high-sugar fermentations, which are conditions where *S. cerevisiae* produces high amounts of this compound [43]. It is possible that no such effect will be achieved with usual sugar concentrations, because *T. delbrueckii*-dominated fermentation takes a long time to complete, and this may lead to the additional oxidation of the wine [23,24]. Sometimes, the claimed improvement may not be very relevant; an example is the reduction in ethanol concentration, which is usually low under anaerobic conditions, and correlates with increased acetic acid and residual sugar contents [44,45], both of which may be undesirable. Moreover, the supposed improvement may not be generally reproducible at commercial wineries, given that contradictory results have been described in experimental vinifications, as is the case for the claimed increased amount of important aromatic compounds: fruity esters, lactones, thiols, and terpenes [23–26,28–30,33,34,46,47].

These circumstances may result in eventual disappointment for oenologists attempting to improve some wine types, such as young white or rosé wines, by using *T. delbrueckii*. The *T. delbrueckii* wines may contain slightly greater amounts of volatile acidity and residual sugars, have roughly the same amount of ethanol, have decreased fresh fruity aromas, or be slightly oxidised with respect to *S. cerevisiae* wines. Considering that using *T. delbrueckii* for winemaking is more expensive and time consuming than using conventional *S. cerevisiae* yeasts, and that most of the commercially available wine strains of the latter yeast are of proven virtuosity for winemaking, most oenologists may refuse to further try *T. delbrueckii* to improve wine quality. Therefore, wine yeast researchers and producer firms should be very careful in advising very precisely those winemaking conditions that may really be appropriate

for wine quality improvement by using *T. delbrueckii*, perhaps even more careful than when the original recommendations were given for inoculating with *S. cerevisiae* yeasts (about one century ago) to improve winemaking with respect to traditional spontaneous must fermentation. Otherwise, the use of *T. delbrueckii* for commercial winemaking may become just anecdotal and circumscribed to experimental research, as will probably end up being the case for many other non-conventional wine yeast species.

Given that these important prior considerations must be kept in mind, we shall now look at some specific approaches using *T. delbrueckii* in terms of its most common effects on wine quality and aroma composition, and which have so far been described as being independent of the winemaking conditions. These are decreased acetic acid production in grape must with a high sugar concentration, decreased concentrations of the main esters, and increased concentrations of lactones and some minor ethyl esters, which are all relative to *S. cerevisiae*-fermented wines. Overall, these effects have led to decreased fresh-fruit aroma intensity, but increased aroma complexity with a dried-fruit/pastry flavour, which is achieved mainly when the *T. delbrueckii* strains reached a high dominance ratio (above 70% of viable cells) at the tumultuous fermentation stage [23,24,26,29,30]. It should be noted that part of this effect on the aroma complexity of wine may be due to the growth of lactic acid bacteria [23]. According to these common effects, the use of *T. delbrueckii* for young (white or rosé) table wines should not be recommended mostly because of the loss of fresh-fruit aromas, which are usually wanted in these types of wine. However, an exception should be considered for those specific situations in which a low dominance ratio of *T. delbrueckii* does not reduce the fresh-fruit aroma intensity of the wine, or it interacts with *S. cerevisiae* to increase the amounts of specific aromas such as volatile thiols (3-sulfanylhexas-1-ol and 3-sulfanylhexasyl acetate) that clearly improve the quality of certain wines, with Sauvignon Blanc being an example [33,48]. Besides this, we think that inoculation with *T. delbrueckii* can mostly be recommended for full-bodied red wines made from grapes with undesirably high malic acid content, natural sweet wines made from grapes with high sugar content, or any aged wine in which the dried-fruit aromas and oxidised character that are usually produced by this yeast may be greatly appreciated by the consumers.

6. Traditional Sparkling Wine Making

There have also recently been trials of the use of *T. delbrueckii* for sparkling wine making. The sequential inoculation of this yeast with *S. cerevisiae* has been shown to be an interesting tool with which to obtain base wines (first fermentation of the grape must) with different characteristics [22]. However, those authors did not analyse the usefulness of *T. delbrueckii* to finally complete the production of sparkling wine (second fermentation of the base wine), probably because they suspected that this yeast could not complete this task. We confirmed these results in a recent work, finding that *T. delbrueckii* could indeed be used to improve base wine quality, but was unable to complete second fermentation, because it cannot survive above 3.5 atm of CO₂ pressure [49]. However, we should point out that base wine that was single-inoculated with *T. delbrueckii* was able to complete second fermentation when there was a very small contamination (4.3×10^{-5} CFU/mL) of *Saccharomyces* yeasts, which became dominant by the time CO₂ pressure increased above 3 atm, and *T. delbrueckii* quickly became inviable (Figure 3). This clearly indicates that CO₂ pressure is another abiotic factor that may decrease the competitiveness of *T. delbrueckii* relative to other potential competitor microorganisms such as *S. cerevisiae*, and that its resistance to high pressure should be genetically improved to get a dominance of this yeast during the second fermentation of sparkling wine.

It should also be mentioned that two *T. delbrueckii* strains (DiSVA 130 and DiSVA 313) single-inoculated in a base wine with high ethanol concentration (11.65% vol) were able to complete the secondary fermentation of sparkling wine even more efficiently than single-inoculated *S. cerevisiae* [32]. This was an unexpected result, mainly because there had been no previous data reported related to the resistance of these DiSVA strains at high ethanol concentrations. Whichever the case, these authors unfortunately did not confirm the dominance of the inoculated strains during second fermentation.

Therefore, some contamination with *S. cerevisiae* yeasts may have occurred, with these being the yeast that was actually responsible for second fermentation completion, as was the case mentioned above (Figure 3).




Figure 3. (Top) Evolution of CO₂ pressure during the second fermentation of cava wine inoculated with *S. cerevisiae* (*Sc*) sensitive to cycloheximide (cyh) and *T. delbrueckii* (*Td*) resistant to cyh. (Bottom) Methylene blue stain to show cell death, and cyh-resistance of viable cells (YEPD+cyh) to monitor the inoculated yeast, for the *T. delbrueckii* inoculated fermentation (7 days, 30 days, and 60 days of fermentation).

7. Proposals for Genetic Improvement of *T. delbrueckii*

The main disadvantages of *T. delbrueckii* that hinder it from becoming as good as *S. cerevisiae* for industrial use are probably its small cell size [50], slow growth rate, and low fermentation vigour under the usual alcoholic fermentation conditions [18,22,24]. These peculiarities result in small harvest quantities in industrial biomass production, and extra difficulties in recovering, washing, and dehydrating the yeast biomass using conventional microfiltration, which may take a long time because the filters become clogged by the small yeast cells, and have to be replaced frequently.

The small cell size of *T. delbrueckii* cells seems to be because it propagates vegetatively as a haploid yeast [7], while the industrial strains of *S. cerevisiae* propagate as diploid or polyploid yeasts. This cell size can be enlarged by increasing the number of chromosomes (ploidy) inducing diploids through perturbed protoplast regeneration or the UV irradiation of intact cells [51–53]. Stable diploid strains have already been constructed from baker's haploid *T. delbrueckii* strains. These diploid strains were about three times larger than the original parental haploid, while both types had similar biomass yield,

stress resistance, gassing power, and sweet dough-leavening ability. The propagation and manipulation of this artificially constructed diploid seemed to be enhanced under industrial conditions, and its stress tolerance and CO₂ production after direct freezing make it possible to reduce the yeast dose in the formulation of frozen sweet bakery products [51,53]. These interesting results for baker's *T. delbrueckii* strains should serve to encourage attempts to perform similar genetic improvements with wine strains, which is a task that remains to be completed.

With regard to the properties that are desired of yeasts for them to perform industrial alcoholic fermentations efficaciously, the tolerance to high glucose concentrations (up to 700 g/L) of *S. cerevisiae* (a poor osmotolerant) was increased by fusing it with the heat-treated protoplasts of an osmotolerant *T. delbrueckii* strain. The interspecific hybrids were able to grow in high glucose concentration media, producing enhanced amounts of ethanol. Further investigation of the practical application of these hybrids is pending [54]. Other than this one-off achievement, there has been little published research related to this issue. In particular, there remains the problem of improving *T. delbrueckii* with respect to the properties of *S. cerevisiae* that make this latter yeast the best choice for most industrial alcoholic fermentations. It is accepted that *T. delbrueckii* may be more resistant than *S. cerevisiae* to various stressors such as freeze–thaw or growth in high osmolality media, which can be very important properties for a baker's yeast used in frozen and frozen sweet dough technology [52,53]. While these properties may be interesting for certain specific wine fermentations, there are other stress resistance properties that are generally considered to be of major importance in winemaking technology, such as the resistance to ethanol, SO₂, or Cu, each of which affects the grape must fermentation vigour of *T. delbrueckii*.

Based on our experience, some strains of *T. delbrueckii* can fully dominate and complete crushed grape fermentation to reach above 14° GL, which is a very high alcohol content for non-*Saccharomyces* yeast fermentation (Figure 4). However, this achievement occurs only rarely, under certain favourable environmental conditions: the presence of only very low amounts of competitor *Saccharomyces* yeasts; a very large inoculum of healthy *T. delbrueckii* cells to start must fermentation quickly and overcome the growth of any other yeasts; frequent agitation to provide extra oxygen; the addition of some extra amounts of nutrients that are usually contained in crushed grapes but which may be scarce in clarified grape must; low amounts of toxic compounds such as SO₂, Cu, pesticides, etc. Even under these favourable conditions, *T. delbrueckii*-dominated fermentation took much longer to complete than *S. cerevisiae*-dominated fermentation, because fermentation vigour and cell viability declined more quickly in *T. delbrueckii* than in *S. cerevisiae* after the wine reached about 5% alcohol (Figure 4B). The differences in ethanol resistance between the two yeast species can be seen on YEPD agar plates supplemented with different ethanol concentration (Figure 2). Similarly, while *T. delbrueckii* is able to complete fermentation in the presence of 50 mg/L SO₂ (Figure 5), although a partial lethal effect cannot be ruled out, it is clearly less resistant to this compound than *S. cerevisiae*. A concentration of 125 mg/L SO₂, which is frequently used in winemaking, is clearly lethal for *T. delbrueckii* (Figure 2). Moreover, there may be an undesirable amount of Cu in grape juice, because Cu compounds are frequently used as fungicides in the vineyard. Again, *S. cerevisiae* is more resistant to Cu than *T. delbrueckii* (Figure 2), meaning that the possible presence of high concentrations (20–30 mg/L) of Cu would reduce fermentation vigour more drastically in the latter yeast than in the former. Experiments to obtain new *T. delbrueckii* strains resistant to ethanol, SO₂, and Cu (as well as to high CO₂ pressures) are currently underway to improve the overall fermentation performance of this species of yeast to bring it as close as possible to that usually shown by selected *S. cerevisiae* wine yeasts. *A priori*, new strains with improved fermentation capacities could be isolated and selected, since there could be intraspecific variations in *T. delbrueckii* as they exist in other yeast species such as *S. cerevisiae*.




Figure 4. Must fermentation kinetics (A) and proportion of dead cells (B) of Garnacha grape fermentation inoculated with *S. cerevisiae* (Sc) and *T. delbrueckii* (Td). The concentrations of ethanol and residual sugars in the wines are given.




Figure 5. Must fermentation kinetics (A) and yeast population dynamics (B) of Pinot Noir grape fermentation inoculated with *S. cerevisiae* (Sc) and *T. delbrueckii* (Td) in the absence and presence of 50 mg/L SO₂.

8. Concluding Remarks

Given that *T. delbrueckii* is not as good a grape must fermenter as the selected *S. cerevisiae* wine yeasts that are currently available in the market at reasonable prices, further research effort will be required to select new *T. delbrueckii* strains or genetically improve the strains that have already been described before this yeast can become generally useful for commercial winemaking. The currently available strains of this yeast present certain handicaps that both limit their performance as good wine fermenters and increase their production costs (and hence their market selling price).

Perhaps most of the considerations that we have made here regarding the possible erroneous assignment of interesting properties to non-*Saccharomyces* yeasts such as *T. delbrueckii* should also have been made for strains of *S. cerevisiae* when they were originally selected for winemaking. However, in this latter case, any possible misassignment would not have been so critical as to dissuade the oenologist from using the selected yeast industrially, because the fermentation vigour of most *S. cerevisiae* strains is fairly good, and one could have expected certain success with the inoculation of these yeasts in the winery compared with the traditional spontaneous fermentation. In particular, it would be rare for any relevant growth of a contaminating non-*Saccharomyces* yeast in must fermentations inoculated with *S. cerevisiae* to occur, and such contaminations should not have any negative impact on wine quality. However, on the contrary, one may expect the massive growth of *Saccharomyces* yeasts in fermentations inoculated with non-*Saccharomyces* such as *T. delbrueckii* to occur frequently, with the consequence of failing to obtain the desired effect on wine quality. Given this current situation, in order to obtain the required improvement of wine quality, we would propose for the time being to use single inoculation with *T. delbrueckii* under appropriate conditions to achieve its full dominance, and then to mix the resulting wines with other *S. cerevisiae* wines.

Author Contributions: Conceptualization, M.R. and R.V.; Methodology, M.R. and R.V.; Formal Analysis, M.R. and R.V.; Investigation, M.R. and R.V.; Resources, M.R.; Writing-Original Draft Preparation, M.R.; Writing-Review & Editing, M.R.; Supervision, M.R. and R.V.; Project Administration, M.R.; Funding Acquisition, M.R.

Funding: This research was funded by grants GR18117 of the Extremadura Regional Government and AGL2017-87635-R of the Spanish Ministry of Economy, Industry and Competitiveness (co-financed with FEDER funds).

Conflicts of Interest: The authors declare no conflict of interest.

References

- Benito, S. The impact of *Torulaspora delbrueckii* yeast in winemaking. *Appl. Microbiol. Biotechnol.* **2018**, *102*, 3081–3094. [[CrossRef](#)] [[PubMed](#)]
- Kurtzman, C.P. *Torulaspora lindner* (1904). In *The Yeasts: A Taxonomic Study*, 5th ed.; Kurtzman, C.P., Fell, J.W., Boekhout, T., Eds.; Elsevier: London, UK, 2011; Volume 2, pp. 867–874.
- Saluja, P.; Yelchuri, R.K.; Sohal, S.K.; Bhagat, G.; Paramjit; Prasad, G.S. *Torulaspora indica* a novel yeast species isolated from coal mine soils. *Antonie van Leeuwenhoek* **2012**, *101*, 733–742. [[CrossRef](#)] [[PubMed](#)]
- Wang, Q.M.; Xu, J.; Wang, H.; Li, J.; Bai, F.Y. *Torulaspora quercuum* sp. Nov. and *Candida pseudohumilis* sp. Nov., novel yeasts from human and forest habitats. *FEMS Yeast Res.* **2009**, *9*, 1322–1326. [[CrossRef](#)] [[PubMed](#)]
- James, S.A.; Collins, M.D.; Roberts, I.N. Use of an rRNA internal transcribed spacer region to distinguish phylogenetically closely related species of the genera *Zygosaccharomyces* and *Torulaspora*. *Int. J. Syst. Bacteriol.* **1996**, *46*, 189–194. [[CrossRef](#)] [[PubMed](#)]
- Oda, Y.; Yabuki, M.; Tonomura, K.; Fukunaga, M. Reexamination of yeast strains classified as *Torulaspora delbrueckii* (lindner). *Int. J. Syst. Bacteriol.* **1997**, *47*, 1102–1106. [[CrossRef](#)] [[PubMed](#)]
- Yarrow, D. Genus 29. *Torulaspora lindner*. In *The Yeasts, A Taxonomic Study*, 3rd ed.; Kreger-van Rij, N.J.W., Ed.; Elsevier Science: Amsterdam, The Netherlands, 1984; pp. 434–439.
- Albertin, W.; Chasseriaud, L.; Comte, G.; Panfili, A.; Delcamp, A.; Salin, F.; Marullo, P.; Bely, M. Winemaking and bioprocesses strongly shaped the genetic diversity of the ubiquitous yeast *Torulaspora delbrueckii*. *PLoS ONE* **2014**, *9*, e94246. [[CrossRef](#)] [[PubMed](#)]
- Wolfe, K.H.; Shields, D.C. Molecular evidence for an ancient duplication of the entire yeast genome. *Nature* **1997**, *387*, 708–713. [[CrossRef](#)] [[PubMed](#)]
- Kreger-van Rij, N.J.; Veenhuis, M. Ultrastructure of the ascospores of some species of the *Torulaspora* group. *Antonie Leeuwenhoek* **1976**, *42*, 445–455. [[CrossRef](#)] [[PubMed](#)]
- Codón, A.C.; Gasent-Ramírez, J.M.; Benítez, T. Factors which affect the frequency of sporulation and tetrad formation in *Saccharomyces cerevisiae* baker's yeasts. *Appl. Environ. Microbiol.* **1995**, *61*, 630–638. [[PubMed](#)]
- Limtong, S.; Imanishi, Y.; Jindamorakot, S.; Ninomiya, S.; Yongmanitchai, W.; Nakase, T. *Torulaspora maleeae* sp. Nov., a novel ascomycetous yeast species from Japan and Thailand. *FEMS Yeast. Res.* **2008**, *8*, 337–343. [[CrossRef](#)] [[PubMed](#)]

13. Ramírez, M.; Ambrona, J. Construction of sterile *ime1Δ* transgenic *Saccharomyces cerevisiae* wine yeasts unable to disseminate in nature. *Appl. Environ. Microbiol.* **2008**, *74*, 2129–2134. [[CrossRef](#)] [[PubMed](#)]
14. Ramírez, M.; Pérez, F.; Regodón, J.A. A simple and reliable method for hybridization of homothallic wine strains of *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.* **1998**, *64*, 5039–5041. [[PubMed](#)]
15. Ramírez, M.; Rebollo, J.E. Technological wine yeast improvement by classical genetic techniques. In *Recent Research Developments in Microbiology*; Pandalai, S.G., Ed.; Research Signpost: Kerala, India, 2003; pp. 87–101.
16. Ramírez, M.; Regodón, J.A.; Pérez, F.; Rebollo, J.E. Wine yeast fermentation vigor may be improved by elimination of recessive growth-retarding alleles. *Biotechnol. Bioeng.* **1999**, *65*, 212–218. [[CrossRef](#)]
17. Castelli, T. Yeasts of wine fermentations from various regions of Italy. *Am. J. Enol. Vitic.* **1955**, *6*, 18–20.
18. Mauricio, J.C.; Millán, C.; Ortega, J.M. Influence of oxygen on the biosynthesis of cellular fatty acids, sterols and phospholipids during alcoholic fermentation by *Saccharomyces cerevisiae* and *Torulaspora delbrueckii*. *World J. Microbiol. Biotechnol.* **1998**, *14*, 405–410. [[CrossRef](#)]
19. Alves-Araujo, C.; Pacheco, A.; Almeida, M.J.; Spencer-Martins, I.; Leao, C.; Sousa, M.J. Sugar utilization patterns and respiro-fermentative metabolism in the baker's yeast *Torulaspora delbrueckii*. *Microbiology* **2007**, *153*, 898–904. [[CrossRef](#)] [[PubMed](#)]
20. Visser, W.; Scheffers, W.A.; Batenburg-van der Vegte, W.H.; van Dijken, J.P. Oxygen requirements of yeasts. *Appl. Environ. Microbiol.* **1990**, *56*, 3785–3792. [[PubMed](#)]
21. Hanl, L.; Sommer, P.; Arneborg, N. The effect of decreasing oxygen feed rates on growth and metabolism of *Torulaspora delbrueckii*. *Appl. Microbiol. Biotechnol.* **2005**, *67*, 113–118. [[CrossRef](#)] [[PubMed](#)]
22. González-Royo, E.; Pascual, O.; Kontoudakis, N.; Esteruelas, M.; Esteve-Zarzoso, B.; Mas, A.; Canals, J.M.; Zamora, F. Oenological consequences of sequential inoculation with non-*Saccharomyces* yeasts (*Torulaspora delbrueckii* or *Metschnikowia pulcherrima*) and *Saccharomyces cerevisiae* in base wine for sparkling wine production. *Eur. Food Res. Technol.* **2014**, *240*, 999–1012. [[CrossRef](#)]
23. Ramírez, M.; Velázquez, R.; Maqueda, M.; Zamora, E.; López-Piñeiro, A.; Hernández, L.M. Influence of the dominance of must fermentation by *Torulaspora delbrueckii* on the malolactic fermentation and organoleptic quality of red table wine. *Int. J. Food Microbiol.* **2016**, *238*, 311–319. [[CrossRef](#)] [[PubMed](#)]
24. Velázquez, R.; Zamora, E.; Alvarez, M.L.; Hernández, L.M.; Ramírez, M. Effects of new *Torulaspora delbrueckii* killer yeasts on the must fermentation kinetics and aroma compounds of white table wine. *Front. Microbiol.* **2015**, *6*, 1222. [[CrossRef](#)] [[PubMed](#)]
25. Renault, P.; Coulon, J.; de Revel, G.; Barbe, J.-C.; Bely, M. Increase of fruity aroma during mixed *T. delbrueckii*/*S. cerevisiae* wine fermentation is linked to specific esters enhancement. *Int. J. Food Microbiol.* **2015**, *207*, 40–48. [[CrossRef](#)] [[PubMed](#)]
26. Azzolini, M.; Fedrizzi, B.; Tosi, E.; Finato, F.; Vagnoli, P.; Scrinzi, C.; Zapparoli, G. Effects of *Torulaspora delbrueckii* and *Saccharomyces cerevisiae* mixed cultures on fermentation and aroma of Amarone wine. *Eur. Food Res. Technol.* **2012**, *235*, 303–313. [[CrossRef](#)]
27. Azzolini, M.; Tosi, E.; Lorenzini, M.; Finato, F.; Zapparoli, G. Contribution to the aroma of white wines by controlled *Torulaspora delbrueckii* cultures in association with *Saccharomyces cerevisiae*. *World J. Microbiol. Biotechnol.* **2015**, *31*, 277–293. [[CrossRef](#)] [[PubMed](#)]
28. Herraiz, G.; Reglero, M.; Herraiz, P.; Alvarez, M.; Cabezudo, M. The influence of the yeast and type of culture on the volatile composition of wine fermented without sulphur dioxide. *Am. J. Enol. Vitic.* **1990**, *41*, 313–318.
29. Comitini, F.; Gobbi, M.; Domizio, P.; Romani, C.; Lencioni, L.; Mannazzu, I.; Ciani, M. Selected non-*Saccharomyces* wine yeasts in controlled multistarter fermentations with *Saccharomyces cerevisiae*. *Food Microbiol.* **2011**, *28*, 873–882. [[CrossRef](#)] [[PubMed](#)]
30. Sadoudi, M.; Tourdot-Maréchal, R.; Rousseaux, S.; Steyer, D.; Gallardo-Chacón, J.-J.; Ballester, J.; Vichi, S.; Guérin-Schneider, R.; Caixach, J.; Alexandre, H. Yeast–yeast interactions revealed by aromatic profile analysis of Sauvignon Blanc wine fermented by single or co-culture of non-*Saccharomyces* and *Saccharomyces* yeasts. *Food Microbiol.* **2012**, *32*, 243–253. [[CrossRef](#)] [[PubMed](#)]
31. Escribano, R.; Gonzalez-Arenzana, L.; Portu, J.; Garijo, P.; Lopez-Alfaro, I.; Lopez, R.; Santamaria, P.; Gutierrez, A.R. Wine aromatic compound production and fermentative behaviour within different non-*Saccharomyces* species and clones. *J. Appl. Microbiol.* **2018**, *124*, 1521–1531. [[CrossRef](#)] [[PubMed](#)]
32. Canonico, L.; Comitini, F.; Ciani, M. *Torulaspora delbrueckii* for secondary fermentation in sparkling wine production. *Food Microbiol.* **2018**, *74*, 100–106. [[CrossRef](#)] [[PubMed](#)]

33. Renault, P.; Coulon, J.; Moine, V.; Thibon, C.; Bely, M. Enhanced 3-sulfanylhexasan-1-ol production in sequential mixed fermentation with *Torulaspora delbrueckii*/*Saccharomyces cerevisiae* reveals a situation of synergistic interaction between two industrial strains. *Front. Microbiol.* **2016**, *7*, 293. [[CrossRef](#)] [[PubMed](#)]
34. Renault, P.; Miot-Sertier, C.; Marullo, P.; Hernández-Orte, P.; Lagarrigue, L.; Lonvaud-Funel, A.; Bely, M. Genetic characterization and phenotypic variability in *Torulaspora delbrueckii* species: Potential applications in the wine industry. *Int. J. Food Microbiol.* **2009**, *134*, 201–210. [[CrossRef](#)] [[PubMed](#)]
35. Ciani, M.; Capece, A.; Comitini, F.; Canonico, L.; Siesto, G.; Romano, P. Yeast interactions in inoculated wine fermentation. *Front. Microbiol.* **2016**, *7*, 555. [[CrossRef](#)] [[PubMed](#)]
36. Ramírez, M.; Velázquez, R.; Maqueda, M.; López-Piñero, A.; Ribas, J.C. A new wine *Torulaspora delbrueckii* killer strain with broad antifungal activity and its toxin-encoding double-stranded RNA virus. *Front. Microbiol.* **2015**, *6*, 983. [[CrossRef](#)] [[PubMed](#)]
37. Ramirez, M.; Velazquez, R.; Lopez-Pineiro, A.; Naranjo, B.; Roig, F.; Llorens, C. New insights into the genome organization of yeast killer viruses based on “atypical” killer strains characterized by high-throughput sequencing. *Toxins* **2017**, *9*, 292. [[CrossRef](#)] [[PubMed](#)]
38. Renault, P.E.; Albertin, W.; Bely, M. An innovative tool reveals interaction mechanisms among yeast populations under oenological conditions. *Appl. Microbiol. Biotechnol.* **2013**, *97*, 4105–4119. [[CrossRef](#)] [[PubMed](#)]
39. Kemsawasd, V.; Branco, P.; Almeida, M.G.; Caldeira, J.; Albergaria, H.; Arneborg, N. Cell-to-cell contact and antimicrobial peptides play a combined role in the death of *Lachancea thermotolerans* during mixed-culture alcoholic fermentation with *Saccharomyces cerevisiae*. *FEMS Microbiol. Lett.* **2015**, *362*, fmv103. [[CrossRef](#)] [[PubMed](#)]
40. Wang, C.; Mas, A.; Esteve-Zarzoso, B. The interaction between *Saccharomyces cerevisiae* and non-*Saccharomyces* yeast during alcoholic fermentation is species and strain specific. *Front. Microbiol.* **2016**, *7*, 502. [[CrossRef](#)] [[PubMed](#)]
41. Taillandier, P.; Lai, Q.P.; Julien-Ortiz, A.; Brandam, C. Interactions between *Torulaspora delbrueckii* and *Saccharomyces cerevisiae* in wine fermentation: Influence of inoculation and nitrogen content. *World J. Microbiol. Biotechnol.* **2014**, *30*, 1959–1967. [[CrossRef](#)] [[PubMed](#)]
42. Velázquez, R.; Zamora, E.; Álvarez, M.L.; Álvarez, M.L.; Ramírez, M. Using mixed inocula of new killer strains of *Saccharomyces cerevisiae* to improve the quality of traditional sparkling-wine. *Food Microbiol.* **2016**, *59*, 150–160. [[CrossRef](#)] [[PubMed](#)]
43. Bely, M.; Stoeckle, P.; Masneuf-Pomarède, I.; Dubourdiou, D. Impact of mixed *Torulaspora delbrueckii*–*Saccharomyces cerevisiae* culture on high-sugar fermentation. *Int. J. Food Microbiol.* **2008**, *122*, 312–320. [[CrossRef](#)] [[PubMed](#)]
44. Contreras, A.; Hidalgo, C.; Schmidt, S.; Henschke, P.A.; Curtin, C.; Varela, C. The application of non-*Saccharomyces* yeast in fermentations with limited aeration as a strategy for the production of wine with reduced alcohol content. *Int. J. Food Microbiol.* **2015**, *205*, 7–15. [[CrossRef](#)] [[PubMed](#)]
45. Arslan, E.; Celik, Z.D.; Cabaroglu, T. Effects of pure and mixed autochthonous *Torulaspora delbrueckii* and *Saccharomyces cerevisiae* on fermentation and volatile compounds of Narince wines. *Foods* **2018**, *7*, 147. [[CrossRef](#)] [[PubMed](#)]
46. Zhang, B.-Q.; Luan, Y.; Duan, C.-Q.; Yan, G.-L. Use of *Torulaspora delbrueckii* co-fermentation with two *Saccharomyces cerevisiae* strains with different aromatic characteristic to improve the diversity of red wine aroma profile. *Front. Microbiol.* **2018**, *9*, 606. [[CrossRef](#)] [[PubMed](#)]
47. Belda, I.; Navascues, E.; Marquina, D.; Santos, A.; Calderon, F.; Benito, S. Dynamic analysis of physiological properties of *Torulaspora delbrueckii* in wine fermentations and its incidence on wine quality. *Appl. Microbiol. Biotechnol.* **2015**, *99*, 1911–1922. [[CrossRef](#)] [[PubMed](#)]
48. Belda, I.; Ruiz, J.; Beisert, B.; Navascues, E.; Marquina, D.; Calderon, F.; Rauhut, D.; Benito, S.; Santos, A. Influence of *Torulaspora delbrueckii* in varietal thiol (3-sh and 4-msp) release in wine sequential fermentations. *Int. J. Food Microbiol.* **2017**, *257*, 183–191. [[CrossRef](#)] [[PubMed](#)]
49. Velázquez, R.; Zamora, E.; Álvarez, M.L.; Ramírez, M. Using *Torulaspora delbrueckii* killer yeasts in the elaboration of base wine and traditional sparkling. *Int. J. Food Microbiol.* **2018**, in press.
50. Spencer, J.F.T.; Spencer, D.M. Taxonomy: The names of the yeasts. In *Yeasts in Natural and Artificial Habitats*; Spencer, J.F.T., Spencer, D.M., Eds.; Springer: Berlin/Heidelberg, Germany, 1997; pp. 11–32.
51. Ohshima, Y.; Sugaura, T.; Horita, M.; Sasaki, T. Industrial application of artificially induced diploid strains of *Torulaspora delbrueckii*. *Appl. Environ. Microbiol.* **1987**, *53*, 1512–1514. [[PubMed](#)]

52. Sasaki, T.; Ohshima, Y. Induction and characterization of artificial diploids from the haploid yeast *Torulaspota delbrueckii*. *Appl. Environ. Microbiol.* **1987**, *53*, 1504–1511. [[PubMed](#)]
53. Hernandez-Lopez, M.J.; Pallotti, C.; Andreu, P.; Aguilera, J.; Prieto, J.A.; Randez-Gil, F. Characterization of a *Torulaspota delbrueckii* diploid strain with optimized performance in sweet and frozen sweet dough. *Int. J. Food Microbiol.* **2007**, *116*, 103–110. [[CrossRef](#)] [[PubMed](#)]
54. Lucca, M.E.; Loray, M.A.; de Figueroa, L.I.C.; Callieri, D.A. Characterisation of osmotolerant hybrids obtained by fusion between protoplasts of *Saccharomyces cerevisiae* and heat treated protoplasts of *Torulaspota delbrueckii*. *Biotechnol. Lett.* **1999**, *21*, 343–348. [[CrossRef](#)]



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).