

Odd Jacobi Manifolds: General Theory and Applications to Generalised Lie Algebroids

ANDREW JAMES BRUCE

Pembrokeshire College, Haverfordwest, Pembrokeshire, SA61 1SZ, UK
andrewjamesbruce@googlemail.com

Presented by Janusz Grabowski

Received February 1, 2012

Abstract: In this paper we define a Grassmann odd analogue of Jacobi structure on a supermanifold. The basic properties are explored. The construction of odd Jacobi manifolds is then used to reexamine the notion of a Jacobi algebroid. It is shown that Jacobi algebroids can be understood in terms of a kind of *curved* Q-manifold, which we will refer to as a quasi Q-manifold.

Key words: supermanifolds, Jacobi structures, Lie algebroids, Q-manifolds.

AMS Subject Class. (2010): 17B70, 53D10, 53D17, 58A50, 83C47.

CONTENTS

1	Introduction	92
2	General Theory	96
2.1	Odd Jacobi structures	96
2.2	Basic examples	103
2.3	The Schoutenization of an odd Jacobi structure	107
2.4	Exact QS-manifolds and odd Jacobi structures	108
3	Jacobi Algebroids	109
3.1	Quasi Q-manifolds and Jacobi algebroids	109
3.2	Lie algebroids in the presence of a 1-cocycle	112
3.3	Schoutenization and Lie algebroids	116
3.4	Odd contact manifolds and Jacobi algebroids	117
4	Concluding remarks	119
5	Appendix: Canonical double vector bundle morphisms	120
	References	122

1. INTRODUCTION

Lichnerowicz [17] introduced the notion of a Poisson manifold as well as a Jacobi manifold. Such manifolds have found applications in classical mechanics and play an important role in quantisation. In this paper we introduce the notion of (Grassmann) odd Jacobi (super)manifolds. The main motivation for doing so is the fact that odd Poisson brackets, also known as Schouten brackets or in the physics literature as antibrackets have found important application in the Batalin-Vilkoviski antifield formalism of gauge theories [4, 5].

Let us briefly recall the definition of a (classical or even) Jacobi manifold¹ as a manifold equipped with a bi-vector Γ and a vector field E such that

$$[[\Gamma, \Gamma]] = 2E \wedge \Gamma, \quad L_E \Gamma = [[E, \Gamma]] = 0, \quad (1.1)$$

where $[[\bullet, \bullet]]$ is the Schouten-Nijenhuis bracket. (There is some leeway here in signs due to conventions). Given these structures one builds a Lie algebra on $C^\infty(M)$ viz

$$\{f, g\}_J = \Gamma(df, dg) + fE(g) - E(f)g.$$

Importantly this Jacobi bracket is not a derivation unless $E = 0$, which reduces the theory to that of Poisson geometry. That is if $E \neq 0$ the Leibniz rule is not exactly satisfied but contains a correction proportional to $\{f, \mathbb{1}\}_J$. For a modern review of Jacobi structures, including \mathbb{Z} -graded versions see [8, 9].

In this paper we construct a Grassmann odd analogue of Jacobi structures on supermanifolds².

Studying geometric structures on supermanifolds provides a method of confirming the philosophy that *supermanifolds can informally be thought of as “manifolds” with commuting and anticommuting coordinates*. More than this, the inclusion of Grassmann odd degrees of freedom allow one to construct geometric structures that have no classical analogue on manifolds. These “odd structures” such are of mathematical interest and often find applications in physics. For example, odd symplectic and Schouten structures are at the heart of the Batalin-Vilkoviski antifield formalism.

The theory of Schouten manifolds and Q-manifolds (supermanifolds with a homological vector field) are both well established within the mathematical

¹This can all be generalised to supermanifolds, however doing so is inessential for this paper.

²We refrain from calling odd Jacobi structures *Schouten-Jacobi structures* as this name has been already allocated in [8] to mean something specific and not identical to constructions found in this work.

physics literature. However, construction of an odd analogue of classical Jacobi structures appears to have been overlooked. The expected properties of such a construction cannot be taken for granted. In this work we perform an examination of the basic properties of odd Jacobi structures.

The theory of odd Jacobi structures is described by “almost Schouten structure” $S \in C^\infty(T^*M)$, that is a Grassmann odd fibre-wise polynomial of degree two and a homological vector field $Q \in \text{Vect}(M)$, together with natural conditions analogous to Eqn.(1.1). The associated odd Jacobi brackets satisfy the standard properties of Schouten brackets, with the exception of the derivation property.

A large proportion of this paper is devoted to making the previous statement concrete and exploring the elementary properties of supermanifolds equipped with odd Jacobi structures. We then proceed to use this technology to reexamine Jacobi algebroids.

Jacobi algebroids were first introduced by Iglesias and Marrero [13] under the name of generalised Lie algebroids. Such structures were the *revisited* by Grabowski and Marmo [8]. Jacobi algebroids represent a nice generalisation of the concept of a Lie algebroid. Indeed, Jacobi algebroids can be understood as Lie algebroids in the presence of a 1-cocycle [13]. Recall the notion of a Lie algebroid as a vector bundle $E \rightarrow M$ equipped with a Lie bracket on the sections $[\bullet, \bullet] : \Gamma(E) \otimes \Gamma(E) \rightarrow \Gamma(E)$ together with an anchor $a : \Gamma(E) \rightarrow \Gamma(TM)$ that satisfy the Leibniz rule

$$[u, fv] = a(u)[f]v + (-1)^{\tilde{u}\tilde{f}}f[u, v],$$

where $u, v \in \Gamma(E)$ and $f \in C^\infty(M)$. The Leibniz rule implies that the anchor is actually a Lie algebra morphism: $a([u, v]) = [a(u), a(v)]$. A Lie algebroid can also be understood in terms of:

1. a homological vector field of weight one on the total space of ΠE ;
2. a weight minus one Poisson structure on the total space of E^* ;
3. a weight minus one Schouten structure on the total space of ΠE^* .

Here Π is the parity reversion functor which shifts the Grassmann parity of the fibre coordinates, but does not effect the base coordinates. Also the parity reversion functor does not effect the assignment of the weights.

At this point we must remark that we will be working the the category of graded manifolds when discussing algebroids. That is we will be working with supermanifolds equipped with a privileged class of atlases where the

coordinates are assigned weights taking values in \mathbb{Z} and the coordinate transformations are polynomial in coordinates with nonzero weights respecting the weight. Generally the weight will be independent of the Grassmann parity. Moreover, any sign factors that arise will be due to the Grassmann parity and we do not include any possible extra signs due to the weight. In simpler terms, we have a manifold equipped with a distinguished class of charts and diffeomorphisms between them respecting the \mathbb{Z}_2 -grading as well as the additional \mathbb{Z} -grading. These gradings then pass over to geometric objects (tensor and tensor-like objects) on graded manifolds. For further details about graded manifolds one can consult [12, 23, 25].

Jacobi algebroids c.f. [8, 13] are understood in terms of a weight minus one even Jacobi bracket on $C^\infty(E^*)$ for a given vector bundle $E \rightarrow M$. By an even Jacobi bracket we mean a Poisson-like bracket in which the Leibnitz rule is weakened in very specific way. We say more about this shortly.

In this paper we address the “odd” approach to Jacobi algebroids generalising 1. and 3. of the above list mimicking the odd-super constructions related to Lie algebroids. We shall show that Jacobi algebroids can be described in terms of an odd vector field $\mathbb{D} \in \text{Vect}(\Pi E)$ of weight one that satisfies the condition that $[\mathbb{D}, \mathbb{D}] = q \mathbb{D}$ for some odd zero form $q \in C^\infty(\Pi E)$.

The description of Lie algebroids in terms of homological vector fields is due to Vaintrob [24]. The deep links between Poisson geometry and Lie algebroids can be traced back to Coste, Dazord & Weinstein [7].

Remark. Antunes and Laurent-Gengoux [2] studied Jacobi structures using the supergeometric formulation of *Buttin’s big bracket*. They use this formalism to efficiently describe Jacobi algebroids and Jacobi bialgebroids. The supergeometric approach of the big bracket is different, but certainly related to that presented here. Grabowski [11], just after this work was completed, provided a systematic and quite general approach to contact and Jacobi structures on graded supermanifolds. Also note the work of Mehta [22], which we will comment on later.

PRELIMINARIES

All vector spaces and algebras will be \mathbb{Z}_2 -graded. We will generally omit the prefix *super*. By *manifold* we will mean a *smooth supermanifold*. We denote the Grassmann parity of an object by *tilde*: $\tilde{A} \in \mathbb{Z}_2$. By *even* or *odd* we will be referring explicitly to the Grassmann parity.

A *Poisson* ($\varepsilon = 0$) or *Schouten* ($\varepsilon = 1$) *algebra* is understood as a vector

space A with a bilinear associative multiplication and a bilinear operation $\{\bullet, \bullet\} : A \otimes A \rightarrow A$ such that:

- *grading* $\widetilde{\{a, b\}}_\varepsilon = \widetilde{a} + \widetilde{b} + \varepsilon,$
- *skewsymmetry* $\{a, b\}_\varepsilon = -(-1)^{(\widetilde{a}+\varepsilon)(\widetilde{b}+\varepsilon)}\{b, a\}_\varepsilon,$
- *Jacobi Identity* $\sum_{\text{cyclic}(a,b,c)} (-1)^{(\widetilde{a}+\varepsilon)(\widetilde{c}+\varepsilon)}\{a, \{b, c\}_\varepsilon\}_\varepsilon = 0,$
- *Leibniz Rule* $\{a, bc\}_\varepsilon = \{a, b\}_\varepsilon c + (-1)^{(\widetilde{a}+\varepsilon)\widetilde{b}}b\{a, c\}_\varepsilon,$

for all homogenous elements $a, b, c \in A$.

If the Leibniz rule does not hold identically, but is modified as

$$\{a, bc\}_\varepsilon = \{a, b\}_\varepsilon c + (-1)^{(\widetilde{a}+\varepsilon)\widetilde{b}}b\{a, c\}_\varepsilon - \{a, \mathbb{1}\}bc, \quad (1.2)$$

then we have even ($\varepsilon = 0$) or odd ($\varepsilon = 1$) *Jacobi algebras*.

A manifold M such that $C^\infty(M)$ is a Poisson/Schouten algebra is known as a *Poisson/Schouten manifold*. In particular the cotangent of a manifold comes equipped with a canonical Poisson structure.

Let us employ natural local coordinates (x^A, p_A) on T^*M , with $\widetilde{x}^A = \widetilde{A}$ and $\widetilde{p}_A = \widetilde{A}$. Local diffeomorphisms on M induce vector bundle automorphism on T^*M of the form

$$\bar{x}^A = \bar{x}^A(x), \quad \bar{p}_A = \left(\frac{\partial x^B}{\partial \bar{x}^A} \right) p_B.$$

We will in effect use the local description as a *natural vector bundle* to define the cotangent bundle of a supermanifold. The canonical Poisson bracket on the cotangent is given by

$$\{F, G\} = (-1)^{\widetilde{A}\widetilde{F}+\widetilde{A}} \frac{\partial F}{\partial p_A} \frac{\partial G}{\partial x^A} - (-1)^{\widetilde{A}\widetilde{F}} \frac{\partial F}{\partial x^A} \frac{\partial G}{\partial p_A}. \quad (1.3)$$

A manifold equipped with an odd vector field Q , such that the non-trivial condition $Q^2 = \frac{1}{2}[Q, Q] = 0$ holds, is known as a *Q-manifold*. The vector field Q is known as a *homological vector field* for obvious reasons.

The structure of a Schouten manifold is encoded in an “odd Hamiltonian” $\bar{S} \in C^\infty(T^*M)$ which Poisson self-commutes, $\{\bar{S}, \bar{S}\} = 0$. The associated Schouten bracket is given by

$$\llbracket f, g \rrbracket = (-1)^{\widetilde{f}+1} \{ \{ \bar{S}, f \}, g \},$$

with $f, g \in C^\infty(M)$.

A Schouten manifold equipped with a homological vector field such that $L_{\bar{Q}}\bar{S} = 0$ is known as a *QS-manifold*, see [25] and for the odd symplectic case see [1]. Importantly the homological vector field is a derivation over the Schouten bracket.

DEFINITION 1. The triple (M, \mathbb{D}, q) with M a manifold, $\mathbb{D} \in \text{Vect}(M)$ an odd vector field and $q \in C^\infty(M)$ an odd function such that:

$$\mathbb{D}^2 = \frac{1}{2} [\mathbb{D}, \mathbb{D}] = q\mathbb{D} \quad \text{and} \quad \mathbb{D}[q] = 0, \quad (1.4)$$

shall be called a *quasi Q-manifold*. The vector field \mathbb{D} shall be known as an *almost homological vector field*. The odd function q shall be known as the *curving function*.

DEFINITION 2. Let (M, \mathbb{D}_M, q_M) and (N, \mathbb{D}_N, q_N) be quasi Q-manifolds and let $\phi : M \rightarrow N$ be a smooth map. Then ϕ is said to be a *morphism of quasi Q-manifolds* if and only if

1. $\mathbb{D}_M(\phi^*f) = \phi^*(\mathbb{D}_Nf)$ for all $f \in C^\infty(N)$, that is the almost homological vector fields are ϕ -related;
2. $\phi^*q_N = q_M$, that is the curving functions match.

Quasi Q-manifolds and their morphisms form a category. Also note that if $q = 0$ then we have the category of *Q-manifolds* and \mathbb{D} is a *homological vector field*, [1]. The curving functions “measure” the failure of the homological condition of \mathbb{D} and thus represent a kind of “curvature”. The other extreme is to set $\mathbb{D} = 0$ and then keep q as some distinguished odd function. For example, one could consider (*higher*) *Schouten manifolds* as examples of quasi Q-manifolds. The far extreme is the trivial structure of $\mathbb{D} = 0$ and $q = 0$ and we recover the full category of supermanifolds.

2. GENERAL THEORY

2.1. ODD JACOBI STRUCTURES

In this section we define odd Jacobi manifolds and show that much of the theory of classical Jacobi manifolds carries over to the odd case. One should of course keep in mind the similarities and difference with Schouten manifolds. To some extent many of the results here are known to experts, though they appear not to have been written down in one place.

DEFINITION 3. An *odd Jacobi structure* (S, Q) on a manifold M consists of

- an odd function $S \in C^\infty(T^*M)$, of degree two in fibre coordinates,
- an odd vector field $Q \in \text{Vect}(M)$,

such that the following conditions hold:

1. the homological condition $Q^2 = \frac{1}{2}[Q, Q] = 0$,
2. the invariance condition $L_Q S = 0$,
3. the compatibility condition $\{S, S\} = -2QS$.

Here $Q \in C^\infty(T^*M)$ is the principle symbol or “Hamiltonian” of the vector field Q . The brackets $\{\bullet, \bullet\}$ are the canonical Poisson brackets on the cotangent bundle of the manifold.

Remark. Note $[E, E] = 0$ automatically for the even vector field E in the definition of an even or classical Jacobi structure. For odd structures this is a non-trivial condition. Specifically, the underlying manifold M is in fact a Q-manifold in the odd case.

Note that the above conditions 1. and 2. can be written entirely in terms of S and Q and the canonical Poisson bracket as

- 1'. $\{Q, Q\} = 0$,
- 2'. $\{Q, S\} = 0$,

which will be very convenient for calculational purposes.

Remark. Note that a QS-manifold can also be understood in a very similar way by taking the principle symbol of the homological vector field. In particular we have the triple (M, \bar{S}, \bar{Q}) , such that $\{\bar{S}, \bar{S}\} = 0$, $\{\bar{Q}, \bar{S}\} = 0$ and $\{\bar{Q}, \bar{Q}\} = 0$.

In natural local coordinates (x^A, p_A) on T^*M the odd Jacobi structure is given by

$$S = \frac{1}{2!} S^{AB}(x) p_B p_A \quad \text{and} \quad Q = Q^A(x) p_A, \quad (2.1)$$

the homological vector field is in local coordinates given by $Q = Q^A \frac{\partial}{\partial x^A}$. In local coordinates the conditions on the structures can be written as

$$\begin{aligned} \{\Omega, \Omega\} &= 2Q^B \frac{\partial Q^A}{\partial x^B} p_A = 0, \\ \{\Omega, S\} &= \left(\frac{1}{2} Q^C \frac{\partial S^{BA}}{\partial x^D} + (-1)^{\tilde{B}} S^{BC} \frac{\partial Q^A}{\partial x^C} \right) p_A p_B = 0, \\ \{S, S\} + 2\Omega S &= (-1)^{\tilde{C}} \left(S^{CD} \frac{\partial S^{BA}}{\partial x^D} + Q^C S^{BA} \right) p_A p_B p_C = 0. \end{aligned}$$

DEFINITION 4. A manifold equipped with an odd Jacobi structure (S, Q) shall be known as an *odd Jacobi manifold*.

As we shall see, the algebra of smooth functions $C^\infty(M)$ of an odd Jacobi manifold is in fact an odd Jacobi algebra. Following the natural analogue of Lichnerowicz's constructions we have the following definition:

DEFINITION 5. The *odd Jacobi bracket* on $C^\infty(M)$ is defined as

$$\begin{aligned} \llbracket f, g \rrbracket_J &= (-1)^{\tilde{f}+1} \{\{S, f\}, g\} - (-1)^{\tilde{f}+1} \{\Omega, fg\} \\ &= (-1)^{(\tilde{B}+1)\tilde{f}+1} S^{BA} \frac{\partial f}{\partial x^A} \frac{\partial g}{\partial x^B} + (-1)^{\tilde{f}} \left(Q^A \frac{\partial f}{\partial x^A} \right) g + f \left(Q^A \frac{\partial g}{\partial x^A} \right), \end{aligned} \quad (2.2)$$

with $f, g \in C^\infty(M)$.

THEOREM 1. *The odd Jacobi bracket defines an odd Jacobi algebra on $C^\infty(M)$. That is the odd Jacobi bracket has the following properties:*

- (i) *Symmetry:* $\llbracket f, g \rrbracket_J = -(-1)^{(\tilde{f}+1)(\tilde{g}+1)} \llbracket g, f \rrbracket_J$;
- (ii) *Jacobi identity:* $\sum_{\text{cyclic}(f,g,h)} (-1)^{(\tilde{f}+1)(\tilde{h}+1)} \llbracket f, \llbracket g, h \rrbracket_J \rrbracket_J = 0$;
- (iii) *Generalised Leibniz rule:* $\llbracket f, gh \rrbracket_J = \llbracket f, g \rrbracket_J h + (-1)^{(\tilde{f}+1)\tilde{g}} g \llbracket f, h \rrbracket_J - \llbracket f, \mathbb{1} \rrbracket_J gh$.

Proof. We proceed to prove the above theorem by making use of the local descriptions.

- (i) The symmetry is clear from the definition given that

$$S^{AB} = (-1)^{\tilde{A}\tilde{B}} S^{BA}.$$

(ii) As the odd Jacobi bracket is odd and skew-antisymmetric it is sufficient to examine the *even diagonal* in proving the Jacobi identity. That is we only need to consider $[[f, [[f, f]]_J]]_J = 0$ for an arbitrary even function. Thus, via direct computation we have

$$\begin{aligned} [[f, [[f, f]]_J]]_J &= (-1)^{\tilde{C}} \left(S^{CD} \frac{\partial S^{BA}}{\partial x^D} + Q^C S^{BA} \right) \frac{\partial f}{\partial x^A} \frac{\partial f}{\partial x^B} \frac{\partial f}{\partial x^C} \\ &\quad - f \left(2(-1)^{\tilde{B}} S^{BC} \frac{\partial Q^A}{\partial x^C} + Q^C \frac{\partial S^{BA}}{\partial x^C} \right) \frac{\partial f}{\partial x^A} \frac{\partial f}{\partial x^B} \\ &\quad + 2f^2 Q^B \frac{\partial Q^A}{\partial x^B} \frac{\partial f}{\partial x^A}, \end{aligned}$$

collecting terms order by order in “ $\frac{\partial f}{\partial x}$ ”. Note that all terms involving higher order derivatives exactly cancel. Then we see that $[[f, [[f, f]]_J]]_J = 0$ given the conditions on S and Q to form an odd Jacobi structure on M . Thus, the Jacobi identity is satisfied.

(iii) Via direct computation in local coordinates it is easy to see that

$$[[f, gh]]_J = [[f, g]]_J h + (-1)^{(\tilde{f}+1)\tilde{g}} g [[f, h]]_J - (-1)^{\tilde{f}} Q^A \frac{\partial f}{\partial x^A} gh.$$

Thus the “anomaly” is given by $[[f, 1]]_J$. ■

DEFINITION 6. Given a function $f \in C^\infty(M)$ the associated *Hamiltonian vector field* is given by

$$\begin{aligned} f &\rightsquigarrow X_f \in \text{Vect}(M) \\ X_f(g) &= (-1)^{\tilde{f}} [[f, g]]_J - Q(f)g. \end{aligned} \tag{2.3}$$

In natural local coordinates the Hamiltonian vector field of a function f is

$$X_f = (-1)^{\tilde{A}\tilde{f}+1} S^{AB} \frac{\partial f}{\partial x^B} \frac{\partial}{\partial x^A} + (-1)^{\tilde{f}} f Q^A \frac{\partial}{\partial x^A}. \tag{2.4}$$

We will explore the properties of Hamiltonian vector fields later. Before we do this, let us examine morphisms of odd Jacobi manifolds.

DEFINITION 7. Let (M_1, S_1, Q_1) and (M_2, S_2, Q_2) be odd Jacobi manifolds. Then a smooth map

$$\phi : M_1 \rightarrow M_2, \tag{2.5}$$

is said to be an *odd Jacobi morphism* if and only if

$$\phi^* \llbracket f, g \rrbracket_{J_2} = \llbracket \phi^* f, \phi^* g \rrbracket_{J_1}, \quad (2.6)$$

for all $f, g \in C^\infty(M_2)$.

In other words, a morphism $\phi : M_1 \rightarrow M_2$ is an odd Jacobi morphism if the associated pull-back morphism is a homomorphism of odd Lie algebras. Odd Jacobi manifolds form a category under composition of odd Jacobi morphisms.

PROPOSITION 1. *Let (M, S, Q) be an odd Jacobi manifold and $\phi : M \rightarrow M$ a diffeomorphism. Then the following are equivalent:*

- (i) ϕ is an odd Jacobi (auto)morphism; $\phi^* \llbracket f, g \rrbracket_J = \llbracket \phi^* f, \phi^* g \rrbracket_J$;
- (ii) $\phi^* S = S$ and $\phi^* Q = Q$;
- (iii) X_f and $X_{\phi^* f}$ are ϕ -related.

Proof. (i) \iff (ii) follows from the fact that pull-back associated with the diffeomorphism ϕ is a symplectomorphism on the cotangent bundle and the definition of the odd Jacobi bracket. Explicitly:

$$\begin{aligned} \phi^* \llbracket f, g \rrbracket_J &= \phi^* \left((-1)^{\tilde{f}+1} \{ \{ S, f \}, g \} - (-1)^{\tilde{f}+1} \{ Q, fg \} \right) \\ &= (-1)^{\tilde{f}+1} \{ \{ \phi^* S, \phi^* f \}, \phi^* g \} - (-1)^{\tilde{f}+1} \{ \phi^* Q, \phi^*(fg) \}. \end{aligned}$$

Then via (i) we obtain (ii).

(ii) \iff (iii) follows similarly. Explicitly:

$$\begin{aligned} \phi^*(X_f(g)) &= \phi^* \left((-1)^f \llbracket f, g \rrbracket_J - Q(fg) \right) \\ &= (-1)^f \llbracket \phi^* f, \phi^* g \rrbracket_J - \{ Q, \phi^*(fg) \} \end{aligned}$$

where we have used (ii). Thus,

$$\phi^*(X_f(g)) = X_{\phi^* f}(\phi^* g).$$

Thus X_f and $X_{\phi^* f}$ are ϕ -related. ■

Taking the nomenclature from classical mechanics, we will say that an odd Jacobi automorphism is a *canonical transformation* with respect to the odd Jacobi bracket.

DEFINITION 8. A vector field $X \in \text{Vect}(M)$ is said to be a *Jacobi vector field* if and only if

$$L_X S = \{\chi, S\} = 0 \quad \text{and} \quad L_X Q = \{\chi, Q\} = 0, \quad (2.7)$$

where $\chi \in C^\infty(T^*M)$ is the symbol or ‘‘Hamiltonian’’ of the vector field X .

The above definition is the infinitesimal version of a Jacobi automorphism. Note that the homological vector field Q is a Jacobi vector field.

LEMMA 1. *Let $X \in \text{Vect}(M)$ be a vector field on an odd Jacobi manifold. Then the following are equivalent:*

- (a) X is a Jacobi vector field;
- (b) X is a derivation over the odd Jacobi bracket

$$X(\llbracket f, g \rrbracket_J) = \llbracket X(f), g \rrbracket_J + (-1)^{\tilde{X}(\tilde{f}+1)} \llbracket f, X(g) \rrbracket_J;$$

- (c) $[X, Y_f] = (-1)^{\tilde{X}} Y_{X(f)}$, for all Hamiltonian vector fields Y_f .

Proof. Let $X \in \text{Vect}(M)$ be a vector field. Then consider the symbol or ‘‘Hamiltonian’’ of such a vector field: $X^A \frac{\partial}{\partial x^A} \rightarrow \chi = X^A p_A \in C^\infty(T^*M)$. The Lie derivative with respect to the vector field acting on $C^\infty(T^*M)$ is just $L_X = \{\chi, \bullet\}$.

(a) \iff (b) is proved via successive use of the Jacobi identity for the canonical Poisson bracket together with (a). Explicitly:

$$\begin{aligned} X(\llbracket f, g \rrbracket_J) &= (-1)^{(\tilde{f}+\tilde{X})+1} \{\{S, \{\chi, f\}\}, g\} - (-1)^{(\tilde{f}+\tilde{X})+1} \{Q, \{\chi, f\}\}g \\ &\quad + (-1)^{\tilde{X}(\tilde{f}+1)+\tilde{f}+1} \{\{S, f\}, \{\chi, g\}\} \\ &\quad - (-1)^{\tilde{X}(\tilde{f}+1)+\tilde{f}+1} \{Q, f\{\chi, g\}\}, \end{aligned}$$

which establishes the result.

(a) \iff (c) via direct computation. Explicitly:

$$\begin{aligned} X(Y_f(g)) &= (-1)^{\tilde{f}} \llbracket X(f), g \rrbracket_J + (-1)^{\tilde{f}+\tilde{X}(\tilde{f}+1)} \llbracket f, X(g) \rrbracket_J \\ &\quad - X(Q(f))g - (-1)^{\tilde{X}(\tilde{f}+1)} Q(f)Q(g), \end{aligned}$$

using the derivation property of X over the odd Jacobi bracket. Then

$$Y_f(X(g)) = (-1)^{\tilde{f}} \llbracket f, X(g) \rrbracket_J - Q(f)X(g),$$

gives

$$[X, Y_f](g) = (-1)^{\tilde{f}} \llbracket X(f), g \rrbracket - (-1)^{\tilde{X}} Q(X(f))g,$$

using 1. Thus the result is established. ■

COROLLARY 1. *The homological vector field Q satisfies the following:*

- (a) $Q(\llbracket f, g \rrbracket_J) = \llbracket Q(f), g \rrbracket + (-1)^{\tilde{f}+1} \llbracket f, Q(g) \rrbracket,$
- (b) $[Q, X_f] = -X_{Q(f)},$

for all $f, g \in C^\infty(M)$.

PROPOSITION 2. *The assignment $f \rightsquigarrow X_f$ is a morphism between the odd Lie algebra on $C^\infty(M)$ provided by the odd Jacobi brackets and the Lie algebra of vector fields. Specifically, the following holds:*

$$[X_f, X_g] = -X_{\llbracket f, g \rrbracket_J} \quad (2.8)$$

for all $f, g \in C^\infty(M)$.

Proof. Writing out the commutator explicitly we obtain

$$\begin{aligned} [X_f, X_g](h) &= (-1)^{\tilde{f}+\tilde{g}} \llbracket f, \llbracket g, h \rrbracket \rrbracket_J - (-1)^{\tilde{f}+\tilde{g}+(\tilde{f}+1)(\tilde{g}+1)} \llbracket g, \llbracket f, h \rrbracket \rrbracket_J \\ &\quad - (-1)^{\tilde{f}} \llbracket f, Q(g) \rrbracket_J h + (-1)^{\tilde{g}+(\tilde{f}+1)(\tilde{g}+1)} \llbracket g, Q(f) \rrbracket_J h \\ &\quad + \text{terms that cancel.} \end{aligned}$$

In the above we have explicitly used the generalised Leibniz rule for the odd Jacobi bracket. Importantly all other possible terms cancel. Then using the Jacobi identity for the odd Jacobi brackets in the form

$$\llbracket \llbracket f, g \rrbracket_J, h \rrbracket_J = \llbracket f, \llbracket g, h \rrbracket_J \rrbracket_J - (-1)^{(\tilde{f}+1)(\tilde{g}+1)} \llbracket g, \llbracket f, h \rrbracket_J \rrbracket_J,$$

and the differential property of Q over the odd Jacobi bracket we obtain

$$[X_f, X_g] = (-1)^{\tilde{f}+\tilde{g}} \llbracket \llbracket f, g \rrbracket_J, h \rrbracket_J + (Q(\llbracket f, g \rrbracket_J)) h,$$

and thus the result is established. ■

Unlike the Schouten or indeed the Poisson case, Hamiltonian vector fields on a Jacobi manifold (both even or odd) in general do not generate infinitesimal automorphisms of the bracket structure.

PROPOSITION 3. *A Hamiltonian vector field $X_f \in \text{Vect}(M)$ is a Jacobi vector field if and only if $f \in C^\infty(M)$ is Q -closed.*

Proof. If and only if f is Q -closed, i.e. $Q(f) = 0$ then $X_f = (-1)^{\tilde{f}} \llbracket f, \bullet \rrbracket_J$. Using Proposition 2. we have

$$[X_f, X_g] = -X_{\llbracket f, g \rrbracket_J} = (-1)^{\tilde{f}+1} X_{X_f(g)},$$

for any $g \in C^\infty(M)$. Then via Lemma 1. the result is established. ■

2.2. BASIC EXAMPLES

Let us turn our attention to straight forward examples to show that the category of odd Jacobi manifolds is not completely empty.

EXAMPLE 1. *The superline.* Consider the supermanifold $\mathbb{R}^{1|1}$, which we equip with local coordinates (t, ξ) . Here t is commuting and ξ anticommuting. There is a canonical odd Jacobi structure on $\mathbb{R}^{1|1}$ given by

$$\begin{aligned} S &= -\pi p, \\ \Omega &= -\pi, \end{aligned}$$

where we have used fiber coordinates (p, π) on $T^*(\mathbb{R}^{1|1})$. Direct computation produces

$$\{S, S\}_{T^*(\mathbb{R}^{1|1})} = -2(-\pi)(-\pi p),$$

which established the fact we have an odd Jacobi structure. Note that as $\pi^2 = 0$, S can also be considered as a Schouten structure. The odd Jacobi brackets are given by

$$\llbracket f, g \rrbracket_J = (-1)^{\tilde{f}} \frac{\partial f}{\partial \xi} \frac{\partial g}{\partial t} - \frac{\partial f}{\partial t} \frac{\partial g}{\partial \xi} - (-1)^{\tilde{f}} \left(\frac{\partial f}{\partial \xi} \right) g - f \left(\frac{\partial g}{\partial \xi} \right),$$

which we recognise as the canonical Schouten bracket on $\mathbb{R}^{1|1}$ plus a term that spoils the strict Leibniz rule.

EXAMPLE 2. *Schouten manifolds.* These are understood as odd Jacobi manifolds for which $Q = 0$. Schouten manifolds are of particular interest in mathematical physics due to their connection with the BV-antifield formalism.

1. Lie-Schouten structures: A vector space \mathfrak{g} is a Lie algebra if and only if $\Pi\mathfrak{g}$ comes equipped with a weight minus one (“linear”) homological vector field. Here Π is the parity reversion functor. If we employ local coordinates (ξ^α) on $\Pi\mathfrak{g}$ then the homological vector field is given by:

$$Q_{\mathfrak{g}} = \frac{1}{2} \xi^\alpha \xi^\beta Q_{\beta\alpha}^\gamma \frac{\partial}{\partial \xi^\gamma}.$$

A weight minus one Schouten structure on $T^*(\Pi\mathfrak{g}^*)$ can be associated with the homological vector field. Employing natural local coordinates $(\eta_\alpha, \pi^\alpha)$ the Schouten structure is given by

$$S = \frac{1}{2} (-1)^{\tilde{\alpha}+\tilde{\beta}} \pi^\alpha \pi^\beta Q_{\alpha\beta}^\gamma \eta_\gamma.$$

The Schouten condition $\{S, S\} = 0$, is equivalent to the homological condition on $Q_{\mathfrak{g}}$ and in turn is equivalent to the Jacobi identity on the initial Lie algebra structure \mathfrak{g} .

2. Odd symplectic manifolds: An odd symplectic manifold is defined as a manifold M of dimensions $(n|n)$ equipped with a closed non-degenerate odd two form denoted ω , understood as a function on the total space of ΠTM . In natural local coordinates the odd symplectic form is given by

$$\omega = \frac{1}{2} dx^A dx^B \omega_{BA}(x).$$

The non degeneracy condition means that, as a matrix ω_{BA} is invertible. Let us denote this inverse by ω^{AB} . Then associated with an odd symplectic form is a Schouten structure given by

$$S = \frac{1}{2} (-1)^{\tilde{B}} \omega^{AB} p_B p_A \in C^\infty(T^*M).$$

The Schouten condition $\{S, S\} = 0$ is directly equivalent to the closedness of ω .

EXAMPLE 3. *Q-manifolds.* These are understood as odd Jacobi manifolds with $S = 0$. The odd Jacobi bracket on a Q-manifold is given by $\llbracket f, g \rrbracket_Q = (-1)^{\tilde{f}} Q(fg)$. Such manifold, and in particular their algebra of functions is of wide interest in mathematics due to the fact that many algebraic structures can be encoded on formal Q-manifolds.

1. The de Rham complex: The algebra of differential (pseudo)forms over a manifold M is understood as the algebra of functions on the manifold ΠTM . The de Rham differential is the canonical homological vector field $d = dx^A \frac{\partial}{\partial x^A}$, where we have employed natural local coordinates (x^A, dx^A) . Then over any manifold, the algebra of differential forms comes equipped with an odd Jacobi bracket given by $\llbracket \alpha, \beta \rrbracket_d = (-1)^{\tilde{\alpha}}(d\alpha)\beta + \alpha(d\beta)$ with $\alpha, \beta \in C^\infty(\Pi TM)$.
2. Lie algebras: Let the vector space \mathfrak{g} be a Lie algebra. Then as seen in the previous example, $(\Pi \mathfrak{g}, Q_{\mathfrak{g}})$ is a Q-manifold. Thus, $C^\infty(\Pi \mathfrak{g})$ has an odd Jacobi bracket (of weight one in linear coordinates) given by

$$\llbracket f, g \rrbracket_Q = \left((-1)^{\tilde{f}} \frac{1}{2} \xi^\alpha \xi^\beta Q_{\beta\alpha}^\gamma \frac{\partial f}{\partial \xi^\gamma} \right) g + f \left(\frac{1}{2} \xi^\alpha \xi^\beta Q_{\beta\alpha}^\gamma \frac{\partial g}{\partial \xi^\gamma} \right).$$

Remark. It is worth noting that this construction of an odd Jacobi bracket generalises directly to L_∞ -algebras (c.f. [15]), which can be understood in terms of homological vector fields inhomogeneous in the linear coordinate ξ . Thus, the associated odd Jacobi bracket is also inhomogeneous in weight, yet remarkably binary. Similarly, one can build an odd Jacobi bracket associated with a Lie algebroid or even an L_∞ -algebroid.

EXAMPLE 4. *Odd contact manifolds.* Rather than diverge into the general theory of contact manifolds let us explore a specific example. Consider the manifold $M := \Pi T^*N \times \mathbb{R}^{0|1}$, where N is a pure even (classical) manifold³.

Let us employ natural local coordinates (x^a, x_a^*, τ) . The coordinates x^a are even, while the other coordinates x_a^* and τ are odd. The dimension of M is $(n|n + 1)$ assuming the dimension of N is n .

The manifold ΠT^*N comes equipped with a canonical odd symplectic structure $\omega = -dx_a^* dx^a$. Also note that all odd symplectic manifolds are equivalent to an anticotangent bundle and that the base manifold can be chosen to be a classical manifold [14, 21]. The manifold M comes equipped with an *odd contact one form*, which is the even one form

$$\alpha = d\tau - x_a^* dx^a.$$

We will for the purposes of this example stipulate that the above is the correct form via “superisation” of the even contact structure on \mathbb{R}^{2n+1} . Clearly

³ Due to a Darboux theorem for contact manifolds this example is also generic. One can generalise the arguments of Arnold (see [3, Appendix 4]) without much difficulty to include odd contact structures on supermanifolds.

the two form $d\alpha = -dx_a^* dx^a$ is, unsurprisingly, the canonical odd symplectic structure on $\Pi T^*N \subset M$. We employ natural fibre coordinates $(dx^a, dx_a^*, d\tau)$ on ΠTM and (p_a, p_a^*, π) on T^*M . The coordinates $dx_a^*, d\tau, p_a$ are even and dx^a, p_a^*, π are odd. Then let us define the almost Schouten structure via

$$\phi_S^*(\alpha) = 0 \quad \text{and} \quad \phi_S^*(d\alpha) = S,$$

where we have the standard fibre-wise morphism $\phi_S : T^*M \rightarrow \Pi TM$ associated with any almost Schouten structure. Let us take the *Ansatz*

$$S = p_a^* (p_a + x_a^* \pi) \in C^\infty(T^*M),$$

based on standard constructions related to even contact structures. Then

$$\begin{aligned} \phi_S^*(dx^a) &= \frac{\partial S}{\partial p_a} = p_a^*, \\ \phi_S^*(dx_a^*) &= -\frac{\partial S}{\partial p_a^*} = -(p_a + x_a^* \pi), \\ \phi_S^*(d\tau) &= -\frac{\partial S}{\partial \pi} = -p_a^* x_a^*. \end{aligned}$$

Direct computation gives

$$\phi_S^*(\alpha) = -p_a^* x_a^* - x_a^* p_a^* = 0 \quad \text{and} \quad \phi_S^*(d\alpha) = (p_a + x_a^* \pi) p_a^* = S.$$

Thus our *Ansatz* is confirmed to be correct. To extract the homological vector field one needs to calculate the self Poisson bracket of the almost Schouten structure. Explicitly

$$\begin{aligned} \{S, S\}_{T^*M} &= 2 \frac{\partial S}{\partial p_a^*} \frac{\partial S}{\partial x_a^*} \\ &= 2((p_a + x_a^* \pi)(-1)p_a^* \pi) \\ &= -2(-\pi)(p_a^*(p_a + x_a^* \pi)), \end{aligned}$$

thus $\mathcal{Q} = -\pi$. It is easy to see that $\{\mathcal{Q}, S\}_{T^*M} = 0$ and $\{\mathcal{Q}, \mathcal{Q}\}_{T^*M} = 0$. Furthermore, notice that the homological vector field $Q = -\frac{\partial}{\partial \tau}$ satisfies

$$i_Q \alpha = 1 \quad \text{and} \quad i_Q(d\alpha) = 0.$$

STATEMENT: Adding an extra odd variable (“odd time”) to the canonical odd symplectic manifold ΠT^*N produces a manifold with an odd Jacobi bracket rather than a Schouten bracket.

In natural local coordinates the odd Jacobi bracket is given by

$$\begin{aligned} \llbracket f, g \rrbracket_J &= (-1)^{\tilde{f}+1} \frac{\partial f}{\partial x_a^*} \frac{\partial g}{\partial x^a} - \frac{\partial f}{\partial x^a} \frac{\partial g}{\partial x_a^*} \\ &+ x_a^* \frac{\partial f}{\partial x_a^*} \frac{\partial g}{\partial \tau} - (-1)^{\tilde{f}+1} \frac{\partial f}{\partial \tau} x_a^* \frac{\partial g}{\partial x_a^*} \\ &+ f \frac{\partial g}{\partial \tau} - (-1)^{\tilde{f}+1} \frac{\partial f}{\partial \tau} g. \end{aligned}$$

We will return to this example in Section 3.4.

2.3. THE SCHOUTENIZATION OF AN ODD JACOBI STRUCTURE

In this section we show that via an extension of an odd Jacobi manifolds one can canonically construct a QS-manifold. This mimics very closely the classical situation of Poissonization.

Consider the manifold $M \times \mathbb{R}$, which we equip with natural coordinates (x^A, t) . Here t is the commuting coordinate on the factor \mathbb{R} . Assuming that M is in fact an odd Jacobi manifold one can build

$$\bar{S} = e^{-t} (S - Qp) \in C^\infty(T^*(M \times \mathbb{R})), \quad (2.9)$$

where p is the momenta associated with t .

THEOREM 2. *Let (M, S, Q) be an odd Jacobi manifold. Then the triple $(M \times \mathbb{R}, \bar{S}, Q)$ is a QS-manifold in the sense of Voronov.*

Proof. First observe that \bar{S} is of order two in the momenta and is Grassmann odd. Second note the Poisson bracket on $T^*(M \times \mathbb{R})$ has the natural decomposition as $\{, \} = \{, \}_{T^*M} + \{, \}_{T^*\mathbb{R}}$. It is then a straight forward exercise to take into account terms that contain conjugate variables and those that do not to show that

1. $\{\bar{S}, \bar{S}\} = e^{-2t} (\{S, S\} + 2Q - 2p\{S, Q\} + p^2\{Q, Q\}),$
2. $\{\bar{S}, Q\} = e^{-t}\{S, Q\} - p\{Q, Q\},$

and thus as (M, S, Q) be an odd Jacobi manifold we establish that $\{\bar{S}, \bar{S}\} = 0$ and $\{\bar{S}, Q\} = 0$. This establishes the proposition. ■

STATEMENT: Adding “time” to an odd Jacobi manifold produces a QS-manifold.

2.4. EXACT QS-MANIFOLDS AND ODD JACOBI STRUCTURES

In this section we define the notion of an exact QS-manifold, understood as a fairly direct generalisation of an exact or homogeneous Poisson manifold. We then, taking Petalidou [20] as our inspiration establish a link between exact QS-structures and odd Jacobi structures. In particular these structures are on the same manifold and no extension is required. This is in contrast to the process of Schoutenization of an odd Jacobi manifold, which requires the manifold to be extended by one even direction.

DEFINITION 9. An *exact QS-manifold* is the quadruple $(M, \bar{S}, \bar{Q}, \bar{E})$, where (M, \bar{S}, \bar{Q}) is a QS-manifold and $\bar{E} \in \text{Vect}(M)$ is an even vector field, referred as the *homothety vector field* that satisfies

$$L_{\bar{E}}\bar{S} = -\bar{S} \quad \text{and} \quad L_{\bar{E}}\bar{Q} = -\bar{Q}. \quad (2.10)$$

The existence of the homothety vector field on a QS-manifold means that both \bar{Q} and \bar{S} are *exact*

$$\begin{aligned} \{\mathcal{E}, \bar{S}\} &= -\bar{S} \quad \longrightarrow \quad \bar{S} = \{\bar{S}, \mathcal{E}\}, \\ \{\mathcal{E}, \bar{Q}\} &= -\bar{Q} \quad \longrightarrow \quad \bar{Q} = \{\bar{Q}, \mathcal{E}\}, \end{aligned}$$

with respect to the operators on $C^\infty(T^*M)$ they generate. Here $\mathcal{E} \in C^\infty(T^*M)$ is the symbol of the homothety vector field. In other words, the Schouten structure is itself a trivial element in the *Schouten cohomology* as generated by $\delta_{\bar{S}} := \{\bar{S}, \bullet\}$. The homological structure is similarly a trivial element in the cohomology of the operator $L_{\bar{Q}}$. Poisson cohomology goes back to Lichnerowicz [17], who also introduced the notion of (even) Jacobi manifolds. For a discussion of the cohomology of a Q-manifold see [18, 19].

Let us now proceed to the theorem relating exact QS structures to odd Jacobi structures on the *same underlying manifold*.

THEOREM 3. *Let $(M, \bar{S}, \bar{Q}, \bar{E})$ be an exact QS-manifold. Then the pair $(S = \bar{S} + \mathcal{E}\bar{Q}, Q = \bar{Q})$, provides an odd Jacobi structure on the manifold M .*

Proof. The proof requires one to examine the the invariance and compatibility conditions for odd Jacobi structures. The homological condition is given.

- Writing out the self-Poisson bracket of S one obtains

$$\begin{aligned} \{S, S\} &= \{\bar{S}, \bar{S}\} + 2\{\bar{S}, \varepsilon\}\bar{Q} + 2\varepsilon\{\bar{S}, \bar{Q}\} \\ &\quad - \{\varepsilon, \varepsilon\}\bar{Q}^2 - 2\varepsilon\{\varepsilon, \bar{Q}\}\bar{Q} + \varepsilon^2\{\bar{Q}, \bar{Q}\} \\ &= 2\bar{Q}(\{\varepsilon, \bar{S}\} + \varepsilon\{\varepsilon, \bar{Q}\}) \\ &= -2\bar{Q}(\bar{S} + \varepsilon\bar{Q}). \end{aligned}$$

- Writing out the Poisson bracket between Q and S one obtains

$$\{Q, S\} = \{\bar{Q}, \bar{S}\} + \{\bar{Q}, \varepsilon\}\bar{Q} + \varepsilon\{\bar{Q}, \bar{Q}\} = \bar{Q}^2 = 0.$$

Thus S and Q define an odd Jacobi structure on the manifold M . ■

Via a mild generalisation of the above proof we arrive at the following corollary:

COROLLARY 2. *Associated with any exact QS-structure on M is a pencil of odd Jacobi structures also on M given by $(S = a\bar{S} + b\varepsilon\bar{Q}, Q = b\bar{Q})$ where a, b are even parameters (or just real numbers).*

STATEMENT: Every exact QS-manifold is also an odd Jacobi manifold.

Setting $a = b = 1$ produces a “canonical” odd Jacobi structure on M . Setting $a = 1$ and $b = 0$ confirms the notion that a Schouten manifold can be thought of as an odd Jacobi manifold with the trivial homological vector field. Setting $a = 0$ and $b = 1$ confirms the notion that a Q-manifolds can be thought of as an odd Jacobi manifold with the trivial Schouten structure. In a loose sense, intermediate values of a and b interpolate between the extremes of Schouten manifolds and Q-manifolds understood as examples of odd Jacobi manifolds.

3. JACOBI ALGEBROIDS

3.1. QUASI Q-MANIFOLDS AND JACOBI ALGEBROIDS

In this section we propose a definition of a Jacobi algebroid in terms of an odd Jacobi structure on the total space of ΠE^* , given a vector bundle $E \rightarrow M$. It will turn out that this definition is equivalent to that given by Grabowski & Marmo [8] (also see Iglesias & Marrero [13]). We postpone the details of this equivalence to the next section and take the following definition as the starting point of this work.

DEFINITION 10. A vector bundle $E \rightarrow M$ is said to have the structure of a *Jacobi algebroid* if and only if the total space of ΠE^* comes equipped with a weight minus one odd Jacobi structure.

Recall that an odd Jacobi structure on a manifold is a pair of odd functions on the total space of the cotangent bundle quadratic and linear in the fibre coordinates together with a series of conditions expressed in terms of the canonical Poisson bracket. Let us employ natural local coordinates $(x^A, \eta_\alpha, p_A, \pi^\alpha)$ on the total space of $T^*(\Pi E^*)$. The weight is assigned as $w(x^A) = 0$, $w(p_A) = 0$, $w(\eta_\alpha) = +1$ and $w(\pi^\alpha) = -1$. This is the *natural weight* associated with the vector bundle structure $E^* \rightarrow M$. The parity of the coordinates is given by $\tilde{x}^A = \tilde{A}$, $\tilde{\eta}_\alpha = (\tilde{\alpha} + 1)$, $\tilde{p}_A = \tilde{A}$ and $\tilde{\pi}^\alpha = (\tilde{\alpha} + 1)$. In these natural local coordinates the odd Jacobi structure is given by

$$\begin{aligned} S &= (-1)^{\tilde{\alpha}} \pi^\alpha Q_\alpha^A(x) p_A + (-1)^{\tilde{\alpha} + \tilde{\beta}} \frac{1}{2} \pi^\alpha \pi^\beta Q_{\beta\alpha}^\gamma \eta_\gamma, \\ \mathcal{Q} &= \pi^\alpha Q_\alpha(x), \end{aligned} \quad (3.1)$$

which are both functions on the total space of $T^*(\Pi E^*)$. The notation and the sign factors employed make clear the relation with Lie algebroids.

This structure satisfies the conditions:

1. $\{\mathcal{Q}, \mathcal{Q}\}_{T^*(\Pi E^*)} = 0$;
2. $\{\mathcal{Q}, S\}_{T^*(\Pi E^*)} = 0$;
3. $\{S, S\}_{T^*(\Pi E^*)} = -2\mathcal{Q}S$.

Setting $\mathcal{Q} = 0$ means that S is a Schouten structure and thus we have a genuine Lie algebroid. Note that due to the fact that the function \mathcal{Q} does not contain conjugate variables the condition 1. is automatically satisfied. This is not generally the case and typically 1. will be a non-trivial condition.

The Jacobi algebroid structure on the vector bundle $E \rightarrow M$ is directly equivalent to the existence of a weight minus one odd Jacobi bracket on $C^\infty(\Pi E^*)$. That is the algebra of “multivector fields” comes equipped with the structure of an odd Jacobi algebra *viz*

$$\begin{aligned} \llbracket X, Y \rrbracket_E &= (-1)^{\tilde{X}+1} \{ \{ S, X \}_{T^*(\Pi E^*)}, Y \}_{T^*(\Pi E^*)} \\ &\quad - (-1)^{\tilde{X}+1} \{ \mathcal{Q}, XY \}_{T^*(\Pi E^*)}, \end{aligned}$$

with $X, Y \in C^\infty(\Pi E^*)$.

In natural local coordinates this bracket is given by

$$\begin{aligned} \llbracket X, Y \rrbracket_E &= Q_\alpha^A \left((-1)^{(\tilde{X}+\tilde{\alpha}+1)(\tilde{A}+1)} \frac{\partial X}{\partial \eta_\alpha} \frac{\partial Y}{\partial x^A} - (-1)^{(\tilde{X}+1)\tilde{\alpha}} \frac{\partial X}{\partial x^A} \frac{\partial Y}{\partial \eta_\alpha} \right) \\ &\quad - (-1)^{(\tilde{X}+1)\tilde{\alpha}+\tilde{\beta}} Q_{\alpha\beta}^\gamma \eta_\gamma \frac{\partial X}{\partial \eta_\beta} \frac{\partial Y}{\partial \eta_\alpha} \\ &\quad + (-1)^{\tilde{X}} Q_\alpha \frac{\partial X}{\partial \eta_\alpha} Y + X Q_\alpha \frac{\partial Y}{\partial \eta_\alpha}, \end{aligned}$$

where

$$X = X(x, \eta) = X(x) + X^\alpha(x) \eta_\alpha + \frac{1}{2!} X^{\alpha\beta}(x) \eta_\beta \eta_\alpha + \dots$$

The above odd Jacobi bracket is the natural generalisation of the weight minus one Schouten bracket associated with a Lie algebroid, which itself is a generalisation of the Schouten-Nijenhuis bracket between multivector fields over a manifold.

THEOREM 4. *The existence of a Jacobi algebroid structure on the vector bundle $E \rightarrow M$ is equivalent to ΠE being a weight one quasi Q -manifold.*

Proof. Recall that the canonical double vector bundle morphism

$$T^*(\Pi E^*) \xrightarrow{R} T^*(\Pi E),$$

is a symplectomorphism between the respective canonical symplectic structures. We place details of this morphism in an appendix. Thus we can move the odd Jacobi structure from ΠE^* to ΠE . However the resulting structure over ΠE will not be a genuine odd Jacobi structure as the degree in momenta (fibre coordinates of the cotangents) is not conserved under the canonical double vector bundle morphism.

Let us employ natural local coordinates $(x^A, \xi^\alpha, p_A, \pi_\alpha)$ on $T^*(\Pi E)$. The weight of the coordinates is assigned as $w(\xi^\alpha) = -1$ and $w(\pi_\alpha) = +1$. The parities are $\tilde{\xi}^\alpha = \tilde{\pi}_\alpha = (\tilde{\alpha} + 1)$. Then the canonical double vector bundle morphism is given by

$$R^*(\pi_\alpha) = \eta_\alpha, \quad R^*(\xi^\alpha) = (-1)^{\tilde{\alpha}} \pi^\alpha.$$

Then let us consider

$$\begin{aligned} \hat{S} &:= (R^{-1})^* S = \xi^\alpha Q_\alpha^A(x) p_A + \frac{1}{2} \xi^\alpha \xi^\beta Q_{\beta\alpha}^\gamma(x) \eta_\gamma, \\ \hat{Q} &:= (R^{-1})^* Q = (-1)^{\tilde{\alpha}} \xi^\alpha Q_\alpha(x), \end{aligned}$$

both of which are functions on the total space of $T^*(\Pi E)$. Note that the function \hat{S} is now linear in moneta and that \hat{Q} is independent of momenta. As R is a symplectomorphism we naturally have

$$\{\hat{S}, \hat{S}\}_{T^*(\Pi E)} = -2\hat{Q}\hat{S} \quad \text{and} \quad \{\hat{Q}, \hat{S}\}_{T^*(\Pi E)} = 0.$$

Then we can “undo” the symbol map which gives an odd vector field on ΠE and an odd function linear in the fibre coordinate:

$$\begin{aligned} \hat{S} &\longrightarrow \mathbb{D} = \xi^\alpha Q_\alpha^A(x) \frac{\partial}{\partial x^A} + \frac{1}{2} \xi^\alpha \xi^\beta Q_{\beta\alpha}^\gamma(x) \frac{\partial}{\partial \xi^\gamma} \in \text{Vect}(\Pi E), \\ \hat{Q} &\longrightarrow q = -(-1)^{\tilde{\alpha}} \xi^\alpha Q_\alpha(x) \in C^\infty(\Pi E). \end{aligned}$$

Note the extra minus sign in the definition of q . As the symbol map takes commutators of vector fields to Poisson brackets, etc., it is not hard to see that the conditions that (S, Q) be an odd Jacobi structure translates to ΠE being a quasi Q-manifold:

$$[\mathbb{D}, \mathbb{D}] = 2q\mathbb{D} \quad \text{and} \quad \mathbb{D}[q] = 0.$$

The grading is with respect to the natural grading associated with the vector bundle structure $E \rightarrow M$. That is we assign the weight as $\bar{w}(x^A) = 0$ and $\bar{w}(\xi^\alpha) = 1$. Note that $\bar{w} = -w$. ■

3.2. LIE ALGEBROIDS IN THE PRESENCE OF A 1-COCYCLE

In this section we in essence restate Grabowski & Marmo’s theorem (see [8, Theorem 5]) giving a one-to-one correspondence between Jacobi algebroids and Lie algebroids in the presence of a 1-cocycle.

As we are considering the total space ΠE to be a graded manifold we naturally have an Euler vector field, which counts the weight of objects via it’s Lie derivative. In natural local coordinates the Euler vector field is given by $\Xi = \xi^\alpha \frac{\partial}{\partial \xi^\alpha}$ as we have assigned weight $\bar{w}(x) = 0$ and $\bar{w}(\xi) = 1$. A “differential form” $\omega \in C^\infty(\Pi E)$ is homogeneous and of weight p if $\Xi(\omega) = p\omega$. In a similar way, a vector field, $V \in \text{Vect}(\Pi E)$ is homogeneous and of weight r if $[\Xi, V] = rV$. The action of the Euler vector field can be extended to higher tensor objects, but we will have no call to use it in this work. In relation to Jacobi algebroids, we will be exclusively interested in object of weight one. Such objects are invariant under the action of the Euler vector field, or in more classical language they are *linear* objects.

PROPOSITION 4. *Let $(\Pi E, \mathbb{D}, q)$ be the weight one quasi Q-manifold associated with a Jacobi algebroid. Then*

$$Q := \mathbb{D} - q\Xi,$$

defines a homological vector field on ΠE of weight one and thus a Lie algebroid structure on $E \rightarrow M$. Furthermore, we have $Q(q) = 0$ and thus we have a Lie algebroid in the presence of a 1-cocycle.

Proof. The weight conditions are clear from the definitions. We need to prove that Q is homological. Explicitly

$$\begin{aligned} Q^2\omega &= \mathbb{D}^2\omega + q\Xi(q\Xi(\omega)) - \mathbb{D}(q\Xi(\omega)) - q\Xi(\mathbb{D}\omega) \\ &= \mathbb{D}^2\omega - q[\Xi, \mathbb{D}]\omega \\ &= \mathbb{D}^2\omega - q\mathbb{D}\omega, \end{aligned}$$

for any $\omega \in C^\infty(\Pi E)$. Then using the fact that we have a quasi Q-manifold gives

$$Q^2 = 0.$$

It is clear that $Q(q) = 0$ and thus we have a 1-cocycle. ■

PROPOSITION 5. *Let $(\Pi E, Q)$ be a Lie algebroid and let $\phi \in C^\infty(\Pi E)$ be an odd 1-cocycle, that is $\Xi(\phi) = 1$, $Q(\phi) = 0$ and $\tilde{\phi} = 1$. Then*

$$(\Pi E, \mathbb{D} = Q + \phi\Xi, q = \phi),$$

defines a quasi Q-manifold of weight one, and thus a Jacobi algebroid.

Proof. The conditions on the weights is clear. Then via calculation we obtain

$$\begin{aligned} \mathbb{D}^2\omega &= Q^2\omega + \phi\Xi(\phi\Xi(\omega)) + Q(\phi\Xi(\omega)) + \phi\Xi(Q\omega) \\ &= \phi[\Xi, Q]\omega = \phi(Q + \phi\Xi)\omega = \phi\mathbb{D}\omega. \end{aligned}$$

The 1-cocycle condition implies $\mathbb{D}(\phi) = 0$. ■

Remark. The above proposition partially generalises to higher order odd cocycles, one loses the homogeneity in weight of the quasi Q-manifold structure. Thus Lie algebroids in the presence of higher cocycles cannot directly be associated with Jacobi algebroids.

THEOREM 5. (GRABOWSKI-MARMO [8]) *There is a one-to-one correspondence between Jacobi algebroids and Lie algebroids in the presence of an odd 1-cocycle.*

We must again remark that everything here is done in the category of supermanifolds and that we have both Grassmann even and odd cocycles. For the classical case where $E \rightarrow M$ is in the category of pure even classical manifolds 1-cocycles are necessarily odd. Thus the above propositions and theorem include the classical structures.

For clarity let us examine the association of a Lie algebroid in the presence of a 1-cocycle with a Jacobi algebroid in natural local coordinates. It is not hard to see that given \mathbb{D} and q we have

$$Q = \xi^\alpha Q_\alpha^A \frac{\partial}{\partial x^A} + \frac{1}{2} \left(\xi^\alpha \xi^\beta Q_{\beta\alpha}^\gamma + (-1)^{\tilde{\alpha}} 2\xi^\alpha Q_\alpha \xi^\gamma \right) \frac{\partial}{\partial \xi^\gamma}, \quad (3.2)$$

$$\phi = (-1)^{\tilde{\alpha}+1} \xi^\alpha Q_\alpha.$$

By careful symmetrisation we see that building the Lie algebroid structure on ΠE associated with a Jacobi algebroid is essentially described by the replacement

$$\mathbb{D} \longrightarrow Q,$$

viz

$$Q_{\beta\alpha}^\gamma \longrightarrow Q_{\beta\alpha}^\gamma - (-1)^{\tilde{\alpha}+\tilde{\beta}} \left(\delta_\alpha^\gamma Q_\beta + (-1)^{(\tilde{\alpha}+1)(\tilde{\beta}+1)} Q_\alpha \delta_\beta^\gamma \right).$$

One can then more-or-less read off the Lie bracket on the sections of E and the anchor map $a : \Gamma(E) \rightarrow \text{Vect}(M)$. Picking a basis of sections (s_α) for $\Gamma(E)$ and being intentionally slack with the signs we have

$$[s_\alpha, s_\beta] = \pm Q_{\alpha\beta}^\gamma s_\gamma \pm Q_\alpha s_\beta \pm s_\alpha Q_\beta,$$

$$a(s_\alpha) = \pm Q_\alpha^A \frac{\partial}{\partial x^A}.$$

Dual to this one can consider the associated Schouten structure which is given by

$$\bar{S} = (-1)^{\tilde{\alpha}} \pi^\alpha Q_\alpha^A p_A + \frac{1}{2} \left((-1)^{\tilde{\alpha}+\tilde{\beta}} \pi^\alpha \pi^\beta Q_{\beta\alpha}^\gamma + (-1)^{\tilde{\gamma}} 2\pi^\alpha Q_\alpha \pi^\gamma \right) \eta_\gamma. \quad (3.3)$$

Similarly, the 1-cocycle becomes $\bar{\phi} = -\pi^\alpha Q_\alpha$ and it is not hard to see that

$$\{\bar{S}, \bar{S}\}_{T^*(\Pi E^*)} = 0,$$

$$\{\bar{S}, \bar{\phi}\}_{T^*(\Pi E^*)} = 0.$$

COROLLARY 3. *If \mathfrak{g} is a Lie algebra with a distinguished odd 1-cocycle $\phi \in C^\infty(\Pi\mathfrak{g})$ then $\Pi\mathfrak{g}^*$ is a (formal) odd Jacobi manifold.*

In local coordinates we have $Q = \frac{1}{2}\xi^\alpha\xi^\beta Q_{\beta\alpha}^\gamma \frac{\partial}{\partial \xi^\gamma} \in \text{Vect}(\Pi\mathfrak{g})$ which encodes the Lie algebra structure on \mathfrak{g} . The 1-cocycle is given by $\phi = (-1)^{\tilde{\alpha}}\xi^\alpha Q_\alpha$, the sign is picked for convenience. Then the odd Jacobi structure on $\Pi\mathfrak{g}^*$ is given by

$$S = (-1)^{\tilde{\alpha}+\tilde{\beta}}\frac{1}{2}\pi^\alpha\pi^\beta Q_{\beta\alpha}^\gamma \eta_\gamma + (-1)^{\tilde{\gamma}}\pi^\alpha Q_\alpha \pi^\gamma \eta_\gamma,$$

$$\mathcal{Q} = \pi^\alpha Q_\alpha.$$

The associated odd Jacobi brackets should be thought of generalisation of the ‘‘Lie-Schouten’’ bracket on $\Pi\mathfrak{g}^*$ [25] in the presence of a 1-cocycle. Both these odd brackets are then considered as odd generalisations of the ‘‘Lie-Poisson-Berezin-Kirillov’’ bracket on \mathfrak{g}^* .

COROLLARY 4. *If $(\Pi E, Q)$ is a Lie algebroid, then $\Pi E^* \times \mathbb{R}^{0|1}$ is a Jacobi algebroid.*

Let us employ natural local coordinates on $T^*(\Pi E^* \times \mathbb{R}^{0|1})$ which we denote as $(x^A, \eta_\alpha, \tau, p_A, \pi^\alpha, \pi)$. The weight assigned to these extra coordinates is $w(\tau) = 1$ and $w(\pi) = -1$. In these local coordinates the weight minus one Jacobi structure is given by

$$S = (-1)^{\tilde{\alpha}}\pi^\alpha Q_\alpha^A p_A + (-1)^{\tilde{\alpha}+\tilde{\beta}}\frac{1}{2}\pi^\alpha\pi^\beta Q_{\beta\alpha}^\gamma \eta_\gamma + \pi\pi^\alpha \eta_\alpha,$$

$$\mathcal{Q} = -\pi.$$

STATEMENT: Extending the fibres of the vector bundle $E \rightarrow M$ underlying a Lie algebroid by \mathbb{R} allows one to directly construct a Jacobi algebroid structure on $\Pi E^* \times \mathbb{R}^{0|1}$.

Naturally the proceeding corollary includes Lie algebra as Lie algebroids over a point. Then, if \mathfrak{g} is a Lie algebra one can extend the vector space structure to $\mathfrak{g} \times \mathbb{R}$. Directly associated with this is the (formal) manifold $\Pi(\mathfrak{g}^* \times \mathbb{R})$ which comes with an odd Jacobi structure of weight minus one.

COROLLARY 5. *Let M be a manifold and \mathbb{A} be a closed, odd one-form (a flat Abelian connection). Then $\Pi T M$ can be made into quasi Q -manifold of weight one, or in other words, $\Pi T^* M$ can be considered as a Jacobi algebroid.*

In natural local coordinates (x^A, dx^A) on $\Pi T M$, the quasi Q-manifold structure is given by

$$\begin{aligned}\mathbb{D} &= d + \mathbb{A} \Xi = dx^A \frac{\partial}{\partial x^A} + dx^B \mathbb{A}_B dx^A \frac{\partial}{\partial dx^A}, \\ q &= \mathbb{A} = dx^B \mathbb{A}_B.\end{aligned}$$

The weight here is simply assigned as $w(x^A) = 0$ and $w(dx^A) = 1$. Picking natural local coordinates (x^A, x_A^*, p_A, p_A^*) on $T^*(\Pi T^* M)$ allows us to write the corresponding odd Jacobi structure on $\Pi T^* M$ as

$$\begin{aligned}S &= (-1)^{\tilde{A}} p_*^A p_A + (-1)^{\tilde{B}} p_*^B \mathbb{A}_B p_*^A x_A^*, \\ Q &= -p_*^A \mathbb{A}_A.\end{aligned}$$

Note that the first term of the almost Schouten structure is the canonical Schouten structure on the anticotangent bundle.

3.3. SCHOUTENIZATION AND LIE ALGEBROIDS

In this section we show that given arbitrary Jacobi algebroid one can extend the structure via the Schoutenisation process described earlier to construct a genuine Lie algebroid. Consider the manifold $T^*(\Pi E^* \times \mathbb{R})$ which we equip with local coordinates $(x^A, \eta_a, t, p_A, \pi^\alpha, p)$. The weight we assign as:

$w(x^A) = 0$	$w(p_A) = 0$
$w(\eta_a) = 1$	$w(\pi^\alpha) = -1$
$w(t) = 0$	$w(p) = 0$

PROPOSITION 6. *Let $(\Pi E^*, S, Q)$ be a Jacobi algebroid. Then $\Pi E^* \times \mathbb{R}$ is a weight minus one Schouten manifold where the Schouten structure is given by*

$$\bar{S} = e^{-t} (S - Qp). \quad (3.4)$$

Proof. Follows directly from Theorem 2. The assignment of the weight follows directly from the definition. ■

In natural local coordinates this Schouten structure is given by

$$\bar{S} = e^{-t} \left((-1)^{\tilde{\alpha}} \pi^\alpha Q_\alpha^A p_A + (-1)^{\tilde{\alpha}+\tilde{\beta}} \frac{1}{2} \pi^\alpha \pi^\beta Q_{\beta\alpha}^\gamma \eta_\gamma - \pi^\alpha Q_\alpha p \right). \quad (3.5)$$

We need to understand the vector bundle structure in order to really identify the Lie algebroid structure. Given the weight assigned to the coordinates on $\Pi E^* \times \mathbb{R}$ the associate underlying (dual) vector bundle structure is $\text{proj}^* E \longrightarrow M \times \mathbb{R}$. That is the pullback of $E \rightarrow M$ by $\text{proj} : M \times \mathbb{R} \rightarrow M$.

COROLLARY 6. *If ΠE^* has the structure of a Jacobi algebroid then $\text{proj}^* E \longrightarrow M \times \mathbb{R}$ is a Lie algebroid.*

STATEMENT: Given a Jacobi algebroid structure on ΠE^* , one can extend the base space M of the underlying vector bundle $E \longrightarrow M$ by \mathbb{R} to directly construct a Lie algebroid.

3.4. ODD CONTACT MANIFOLDS AND JACOBI ALGEBROIDS

In this section we show that the manifold $M := \Pi T^*N \times \mathbb{R}^{0|1}$ considered as an odd contact manifold provides a canonical example of a Jacobi algebroid than lends itself to the description in terms of odd Jacobi brackets.

Let N be a pure even classical manifold of dimension n . Consider the manifold $M := \Pi T^*N \times \mathbb{R}^{0|1}$ equipped with natural local coordinates (x^a, x_a^*, τ) . The coordinates x^a are even, while the other coordinates x_a^* and τ are odd. The dimension of M is $(n|n + 1)$. The manifold M comes equipped with an *odd contact one form*, which is the even one form

$$\alpha = d\tau - x_a^* dx^a. \tag{3.6}$$

It was shown in Example 4. that M is an odd Jacobi manifold with the odd Jacobi structure being

$$\begin{aligned} S &= p_*^a (p_a + x_a^* \pi), \\ Q &= -\pi, \end{aligned} \tag{3.7}$$

where we have employed natural coordinates $(x^a, x_a^*, \tau, p_a, p_*^a, \pi)$ on T^*M . Indeed this odd Jacobi structure is directly equivalent to the odd contact structure. Without details, both the odd contact and odd Jacobi structure on M can be considered as the “natural superisation” of the classical structures on \mathbb{R}^3 . Note that ΠT^*N comes equipped with a canonical Schouten (odd symplectic) structure, but $\Pi T^*N \times \mathbb{R}^{0|1}$ comes with a canonical odd Jacobi structure.

Let us attach the weight to the local coordinates on M as:

$w(x^a) = 0$	$w(p_a) = 0$
$w(x_a^*) = 1$	$w(p_*^a) = -1$
$w(\tau) = 1$	$w(\pi) = -1$

This weight is the “natural weight” with respect to the underlying vector bundle structure $T^*N \times \mathbb{R} \rightarrow N$. With respect to this weight it is clear that the odd Jacobi structure on $M := \Pi T^*N \times \mathbb{R}^{0|1}$ is of weight minus one and we thus have a Jacobi algebroid.

Now consider $M^* := \Pi TN \times \mathbb{R}^{0|1}$ equipped with natural local coordinates $(x^a, \xi^a, \eta, p_a, \pi_a, \theta)$. The canonical double vector bundle morphism $R : T^*M \rightarrow T^*M^*$ act on the coordinates as $R^*(\xi^a) = p_*^a$, $R^*(\eta) = \pi$, $R^*(\pi_a) = x_a^*$ and $R^*(\theta) = \tau$. Then we can pull-back the odd Jacobi structure to give

$$\hat{S} = \xi^a(p_a + \pi_a \eta), \quad \hat{Q} = -\eta,$$

both of which are now functions on the total space of M^* . Then we can “undo” the symbol (and after a little reordering) to produce

$$\mathbb{D} = \xi^a \frac{\partial}{\partial x^a} + \eta \xi^a \frac{\partial}{\partial \xi^a}, \quad q = \eta. \tag{3.8}$$

Direct calculation confirms that $M^* := \Pi TN \times \mathbb{R}^{0|1}$ is a quasi Q-manifold.

STATEMENT: In light of Proposition 4, $TN \times \mathbb{R} \rightarrow N$ is a Lie algebroid in the presence of a 1-cocycle. The de Rham differential on N is the associated homological vector field and the 1-cocycle is identified with the “odd time”.

Remark. As this work was being completed, Mehta [22] established a one-to-one correspondence between Jacobi manifolds and degree 1 contact NQ -manifolds. Mehta shows how to interpret the “Poissonisation” of a Jacobi manifold as the “symplectification” of the corresponding degree 1 contact NQ -manifold. There is no doubt that Mehta’s results can be slightly reformulated to sit comfortably with the conventions used here: one would consider “Shoutenization” and “symplectification” of odd contact structures. This generalises the correspondence between Poisson manifolds and degree 1 symplectic NQ -manifolds, as established by Roytenberg [23]. We also direct the reader to Grabowski [11] who studies degree 2 contact NQ -manifolds as a generalisation of Courant algebroids. The author conjectures that interest in contact structures on super and graded manifolds will continue to grow.

4. CONCLUDING REMARKS

In this paper we defined the notion of an odd Jacobi manifold and examined their basic properties. In particular, it was shown that on a supermanifold equipped with an odd Jacobi structure $J := (S, Q)$ the algebra of smooth functions over the supermanifold $C^\infty(M)$ comes with the structure of an odd Jacobi algebra. Furthermore such the homological vector field Q satisfies a derivation property over the odd Jacobi brackets.

However, it remains open as to if interesting or realistic gauge theories exist that require the use of odd Jacobi structures (with $Q \neq 0$) in a generalised Batalin-Vilkovisky formalism. The notion of a gauge system c.f. [18] in the context of odd Jacobi manifolds is straight forward.

In truth there appears no applications in physics of even Jacobi structures that cannot simply be restated in terms of Poisson geometry. That said, it is conceivable that odd contact structures could find quite direct application in theories with explicit dependency on gauge parameters.

The idea of “odd time” (see Example 4) has already been applied in the Batalin-Vilkovisky formalism to get at general and direct solutions of the master equation for a large class of gauge theories, see Dayi [6]. In essence one understands the BRST operator as the partial derivative with respect to the “odd time” and then one can formulate the BV formalism in a way akin to classical mechanics. It would be very desirable to properly understand the supergeometry of Dayi’s constructions and how this relates to the work here, in particular to odd contact manifolds. The notion of “odd time” is also essential when constructing flows of odd vector fields. It is certainly expected that odd contact structures are of wider interest than just their relation with odd Jacobi manifolds.

We defined Jacobi algebroids in terms of an odd Jacobi structure on ΠE^* of weight minus one. That is the “multivector fields” come equipped with an odd Jacobi bracket. For Lie algebroids the bracket between “multivector fields” is a Schouten bracket, i.e., satisfies a strict Leibniz rule.

This construction was then used to construct a weight one almost homological vector field. That is the “differential forms” come equipped with a kind of *deformed* de Rham differential. Importantly we no longer have a homological vector field as in the case of Lie algebroids, but rather the homological condition is weakened in a very specific way as to provide a quasi Q -manifold structure. As such Jacobi algebroids can be considered as very specific examples of *skew algebroids* [10], which are a kind of Lie algebroid in which the Jacobi identity is lost. If the corresponding anchor is a Lie al-

gebra morphism between sections of the vector bundle and vector fields over the base then we have the notion of an *almost Lie algebroid* [16]. Interest in algebroids without the Jacobi identity comes from nonholonomic mechanics, where skew algebroids provide a general geometric setting.

However, via a simple redefinition one can rephrase Jacobi algebroids in terms of Lie algebroids in the presence of a 1-cocycle, which are also known as generalised Lie algebroids. In doing so we recover, maybe up to conventions, the notion of a Jacobi algebroid in the sense of [8, 13].

The obvious areas of the present work that require further illumination include:

- Do Jacobi bialgebroids have an efficient description in terms of a compatible odd Jacobi structure and a quasi Q-structure?
- Can one define non-linear Jacobi algebroids in terms of odd Jacobi structures over non-negatively graded supermanifolds? Are these naturally related to Voronov's ([26]) non-linear Lie algebroids?
- Can one develop a theory of higher or homotopy odd Jacobi structures and the related L_∞ -algebras.

5. APPENDIX: CANONICAL DOUBLE VECTOR BUNDLE MORPHISMS

For completeness we present the canonical double vector bundle morphisms used in this work. In particular we prove that the morphisms are symplectomorphisms. We describe vector bundles in terms of coordinates on their total spaces and the associated vector bundle automorphisms. Specifically we have:

$E \longrightarrow M$	$E^* \longrightarrow M$
$(x^A, e^\alpha) \mapsto (x^A)$	$(x^A, e_\alpha) \mapsto (x^A)$
$\bar{x}^A = \bar{x}^A(x)$	$\bar{x}^A = \bar{x}^A(x)$
$\bar{e}^\alpha = e^\beta T_\beta^\alpha(x)$	$\bar{e}_\alpha = (T^{-1})_\alpha^\beta(x)$

Where $T_\beta^\gamma (T^{-1})_\gamma^\alpha = \delta_\beta^\alpha$, etc. We take $\widetilde{e}^\alpha = \widetilde{e}_\alpha = \widetilde{\alpha}$.

Let us employ natural local coordinates:

$T^*(\Pi E^*)$	$(x^A, \eta_\alpha, p_A, \pi^\alpha)$
$T^*(\Pi E)$	$(x^A, \xi^\alpha, p_A, \pi_\alpha)$

The Grassmann parities are given by $\tilde{x}^A = \tilde{p}_A = \tilde{A}$, $\tilde{\eta}_\alpha = \tilde{\pi}^\alpha = \tilde{\pi}_\alpha = \tilde{\xi}^\alpha = (\tilde{\alpha} + 1)$. The weights are assigned as $w(x^A) = 0$, $w(\eta_\alpha) = 1$, $w(p_A) = 0$, $w(\pi^\alpha) = -1$, $w(\xi^\alpha) = -1$, $w(\pi_\alpha) = 1$. The admissible changes of coordinates are:

$T^*(\Pi E^*)$	$\bar{x}^A = \bar{x}^A(x), \quad \bar{\eta}_\alpha = (T^{-1})_\alpha^\beta \eta_\beta$ $\bar{p}_A = \left(\frac{\partial x^B}{\partial \bar{x}^A} \right) p_B + (-1)^{\tilde{A}(\tilde{\gamma}+1)+\tilde{\delta}} \tilde{\pi}^\delta T_\delta^\gamma \left(\frac{\partial (T^{-1})_{\gamma^\alpha}}{\partial \bar{x}^A} \right) \eta_\alpha$ $\bar{\pi}^\alpha = (-1)^{\tilde{\alpha}+\tilde{\beta}} \tilde{\pi}^\beta T_\beta^\alpha$
$T^*(\Pi E)$	$\bar{x}^A = \bar{x}^A(x), \quad \bar{\xi}^\alpha = \xi^\beta T_\beta^\alpha$ $\bar{p}_A = \left(\frac{\partial x^B}{\partial \bar{x}^A} \right) p_B + (-1)^{\tilde{A}(\tilde{\gamma}+1)} \xi^\delta T_\delta^\gamma \left(\frac{\partial (T^{-1})_{\gamma^\alpha}}{\partial \bar{x}^A} \right) \pi_\alpha$ $\bar{\pi}_\alpha = (T^{-1})_\alpha^\beta \pi_\beta$

The canonical double vector bundle morphism $R : T^*(\Pi E^*) \rightarrow T^*(\Pi E)$ is given in local coordinates by

$$R^*(\pi_\alpha) = \eta_\alpha, \quad R^*(\xi^\alpha) = (-1)^{\tilde{\alpha}} \tilde{\pi}^\alpha.$$

LEMMA. *The canonical double vector bundle morphism*

$$R : T^*(\Pi E^*) \rightarrow T^*(\Pi E)$$

is a symplectomorphism.

Proof. The canonical even symplectic structure on $T^*(\Pi E^*)$ is given by $\omega_{T^*(\Pi E^*)} = dp_A dx^A + d\pi^\alpha d\eta_\alpha$ and on $T^*(\Pi E)$ is given by $\omega_{T^*(\Pi E)} = dp_A dx^A + d\pi_\alpha d\xi^\alpha$. Thus, $R^*\omega_{T^*(\Pi E)} = \omega_{T^*(\Pi E^*)}$ and we see that R is indeed a symplectomorphism. ■

ACKNOWLEDGEMENTS

The author would like to thank Janusz Grabowski, Giuseppe Marmo and Rajan Amit Mehta for their comments and suggestions on earlier drafts of this work. The author must also thank the anonymous referee for their valuable comments.

REFERENCES

- [1] M. ALEXANDROV, M. KONTSEVICH, A. SCHWARZ, O. ZABORONSKY, The Geometry of the master equation and topological quantum field theory, *Internat. J. Modern Phys. A* **12** (1997), 1405–1429.
- [2] P. ANTUNES, C. LAURENT-GENGOUX, Jacobi structures in supergeometric formalism, *J. Geom. Phys.* **61** (2011), 2254–2266.
- [3] V.I. ARNOLD, “Mathematical Methods of Classical Mechanics”, Translated from the 1974 Russian original by K. Vogtmann and A. Weinstein (corrected reprint of the second edition 1989), Graduate Texts in Mathematics, 60, Springer-Verlag, New York.
- [4] I.A. BATALIN, G.A. VILKOVISKY, Gauge algebra and quantization, *Phys. Lett. B* **102** (1981), 27–31.
- [5] I.A. BATALIN, G.A. VILKOVISKY, Quantization of gauge theories with linearly dependent generators, *Phys. Rev. D (3)* **28** (1983), 2567–2582.
- [6] Ö.F. DAYI, Odd time formulation of the Batalin-Vilkovisky method of quantization, *Modern Phys. Lett. A* **4** (1989) 361–365.
- [7] A. COSTE, P. DAZORD, A. WEINSTEIN, “Groupoïdes Symplectiques”, Publ. Dép. Math. Nouvelle Sér. A, 87-2, Univ. Claude-Bernard, Lyon, 1987.
- [8] J. GRABOWSKI, G. MARMO, Jacobi structures revisited, *J. Phys. A* **34** (2001), 10975–10990.
- [9] J. GRABOWSKI, G. MARMO, The graded Jacobi algebras and (co)homology, *J. Phys. A* **36** (2003), 161–181.
- [10] J. GRABOWSKI, P. URBAŃSKI, Algebroids—general differential calculi on vector bundles, *J. Geom. Phys.* **31** (1999), 111–141.
- [11] J. GRABOWSKI, Graded contact manifolds and principal Courant algebroids, [arXiv:1112.0759 \[math.DG\]](https://arxiv.org/abs/1112.0759), 2011.
- [12] J. GRABOWSKI, M. ROTKIEWICZ, Higher vector bundles and multi-graded symplectic manifolds, *J. Geom. Phys.* **59** (2009), 1285–1305.
- [13] D. IGLESIAS, J.C. MARRERO, Generalized Lie bialgebroids and Jacobi structures, *J. Geom. Phys.* **40** (2001), 176–199.
- [14] H.M. KHUDAVERDIAN, Semidensities on odd symplectic supermanifold, *Comm. Math. Phys.* **247** (2004), 353–390.
- [15] T. LADA, J. STASHEFF, Introduction to SH Lie algebras for physicists, *Internat. J. Theoret. Phys.* **32** (1993), 1087–1103.
- [16] M. DE LEÓN, J.C. MARRERO, D. MARTÍN DE DIEGO, Linear almost Poisson structures and Hamilton-Jacobi equation, Applications to nonholonomic mechanics, *J. Geom. Mech.* **2** (2010), 159–198.
- [17] A. LICHTNEROWICZ, Les variétés de Poisson et leurs algèbres de Lie associées, *J. Differential Geometry* **12** (1977), 253–300.
- [18] S.L. LYAKHOVICH, A.A. SHARAPOV, Characteristic classes of gauge systems, *Nuclear Phys. B* **703** (2004), 419–453.

- [19] S.L. LYAKHOVICH, E.A. MOSMAN, A.A. SHARAPOV, Characteristic classes of Q-manifolds: Classification and applications, *J. Geom. Phys.* **60** (2010), 729–759.
- [20] F. PETALIDOU, On a new relation between Jacobi and homogeneous Poisson manifolds, *J. Phys. A* **35** (2002), 2505–2518.
- [21] A.S. SCHWARZ, Geometry of Batalin-Vilkovisky quantization, *Comm. Math. Phys.* **155** (1993), 249–260.
- [22] R.A. MEHTA, Differential graded contact geometry and Jacobi structures, [arXiv:1111.4705v1](https://arxiv.org/abs/1111.4705v1) [math.SG], 2011.
- [23] D. ROYTENBERG, On the structure of graded symplectic supermanifolds and Courant algebroids, in “Quantization, Poisson Brackets and Beyond” (Manchester, 2001), *Contemp. Math.*, 315, Amer. Math. Soc., Providence, RI, 2002, 169–185.
- [24] A.YU. VAĬNTROB, Lie algebroids and homological vector fields (Russian), *Uspekhi Mat. Nauk* **52** (1997), 161–162; translation in *Russian Math. Surveys* **52** (1997), 428–429.
- [25] TH. VORONOV, Graded manifolds and Drinfeld doubles for Lie bialgebroids, in “Quantization, Poisson Brackets and Beyond” (Manchester, 2001), *Contemp. Math.*, 315, Amer. Math. Soc., Providence, RI, 2002, 131–168.
- [26] TH.TH. VORONOV, Q-manifolds and higher analogs of Lie algebroids, in “XXIX Workshop on Geometric Methods in Physics”, *AIP Conf. Proc.*, 1307, Amer. Inst. Phys., Melville, NY, 2010, 191–202; also available as [arXiv:1010.2503v1](https://arxiv.org/abs/1010.2503v1) [math.DG]