

1 **Table 1.** Types of poisoned baits (positives) based on material preparation, and presentation.

Type of bait	Baits: n (%)	Cases: n
Pieces of fresh meat	61 (30.9%)	43
Commercial chemical substances	32 (16.2%)	13
Sausages and dried cured meat	28 (14.2%)	15
Commercial baits (rodenticides)	21 (10.1%)	9
Impregnated objects (perches)	15 (7.6%)	12
Small animal carcasses	13 (6.6%)	5
Cooked human food	11 (5.6%)	7
Viscera (organs and placenta)	9 (4.6%)	6
Covered seeds	3 (1.5%)	3
Fish and seafood	3 (1.5%)	2
Vegetables	2 (1.0%)	1
TOTAL	197 (100%)	141

2

3

4 **Table 2.** Substances identified in the toxicological analysis of the baits submitted in the
5 present study.

Substance	Baits: n (%)	Cases: n
No toxic substance detected (negatives)	49 (19.9%)	42
Aldicarb	43 (17.5%)	31
Aldicarb (microgranules)	38 (15.4%)	26
Carbofuran	25 (10.2%)	22
Carbofuran (microgranules)	14 (5.7%)	9
Fenthion	15 (6.1%)	12
Brodifacoum	10 (4.1%)	1
Fenamiphos	7 (2.8%)	7
Bromadiolone	6 (2.4%)	4
Aldicarb + carbofuran	5 (2.0%)	3
Chlorpyrifos	4 (1.6%)	1
Malathion	4 (1.6%)	3
Bromadiolone + brodifacoum	3 (1.2%)	2

Chlorpyrifos + methomyl + endosulfan	3 (1.2%)	2
Triflumuron	2 (0.8%)	2
Phenobarbital	2 (0.8%)	1
Strychnine	2 (0.8%)	1
Chlorpyrifos (+ sulfotep)	2 (0.8%)	2
Methomyl	1 (0.4%)	1
Difenacoum	1 (0.4%)	1
Ethoxyquin	1 (0.4%)	1
Metaldehyde	1 (0.4%)	1
Oxamyl	1 (0.4%)	1
Monocrotophos	1 (0.4%)	1
Deltamethrin	1 (0.4%)	1
Aldicarb + lindane	1 (0.4%)	1
Fenamiphos + aldicarb	1 (0.4%)	1
Methiocarb + imidacloprid + metalaxyl + thiram + atrazine	1 (0.4%)	1
Brodifacoum + difenacoum	1 (0.4%)	1
Cypermethrin + chlorfenvinphos + chlorpyrifos + fenitrothion	1 (0.4%)	1
TOTAL	246 (100%)	183

6

7

8