

On Normal Stratified Pseudomanifolds[†]

G. PADILLA

*Universidad Central de Venezuela, Escuela de Matemáticas, Facultad de Ciencias,
Ciudad Universitaria, Caracas 1010, Venezuela*

e-mail: gabrielp@euler.ciens.ucv.ve

(Presented by Manuel de León)

AMS Subject Class. (2000): 55N33, 35S35

Received March 7, 2003

*Devoted to the victims of the natural tragedy in Vargas, Dec. 15/1999,
who died under the rage of Waraira Repano.*

FOREWORD

A stratified pseudomanifold is normal if its links are connected. A normalization of a stratified pseudomanifold X is a normal stratified pseudomanifold X^N together with a finite-to-one projection $\mathbf{n} : X^N \rightarrow X$ preserving the intersection homology. Recall that intersection homology is the suitable algebraic tool for the stratified point of view: it was first introduced by Goresky and MacPherson in the pl -category and later extended for any topological stratified pseudomanifold [2], [3]. Following Borel the map \mathbf{n} is usually required to satisfy the following property: For each $x \in X$ there is a distinguished neighborhood U such that the points of $\mathbf{n}^{-1}(x)$ are in correspondence with the connected components of the regular part of U . A normalization satisfying the above condition always exists for any pl -stratified pseudomanifold [4], [6]. In this article we study the main properties of the map \mathbf{n} . More precisely, we prove that \mathbf{n} can be required to satisfy a stronger condition: it is a locally trivial stratified morphism preserving the conical structure transverse to the strata. We make an explicit construction of such a normalization for any topological stratified pseudomanifold. Our construction is functorial, thus unique. We exhibit the relationship between the stratifications of X and X^N . Finally

[†]This paper was supported by the Euskal Herria Unibertsitatea's Math Department, the University of Artois' LaboGA and the Universidad Central de Venezuela's C.D.C.H.

we prove that the normalization preserves the intersection homology with the family of perversities given in [7], see also [5], [9]. This family of perversities is larger than the usual one. With little adjust our procedure holds also in the C^∞ category.

1. STRATIFIED PSEUDOMANIFOLDS

For a detailed treatment of the results contained in this section, see [8].

Manifolds considered in this paper will always be topological manifolds. A topological space is stratified if it can be written as a disjoint union of manifolds which are related by an incidence condition.

DEFINITION 1.1. Let X be a Hausdorff, paracompact, second countable space. A *stratification* of X is a locally finite partition \mathcal{S} of X in connected and locally closed subspaces called *strata*, satisfying

- (1) Each stratum is a manifold with the induced topology.
- (2) If a stratum S intersects the closure $\overline{S'}$ of another stratum S' then $S \subset \overline{S'}$ and we say S' lies on S .
- (3) There exist open strata, all of them having the same dimension.

If this situation we say X is a *stratified space*. A stratum is *regular* if it is open and *singular* if not; the *dimension* of X is the dimension of the open strata; which we write $\dim(X)$. The *regular part* (resp. *singular part*) is the union of regular (resp. singular) strata. We write the singular part with the symbol Σ . For any paracompact subspace $Y \subset X$ the *induced partition* is the family \mathcal{S}_Y whose elements are the connected component of $Y \cap S$, where S runs over the strata of X . If \mathcal{S}_Y is a stratification of Y then we say that Y is a *stratified subspace*.

EXAMPLES 1.2. Here there are some examples of stratified spaces:

- (1) For every manifold the canonical stratification is the family of its connected components. For every connected manifold M and every stratified space X the canonical stratification of the product $M \times X$ is the family of products $M \times S$ such that S is a stratum of X .
- (2) The canonical stratification of the n -simplex $\Delta \subset \mathbb{R}^{n+1}$ is the family of connected components of $\Delta_j - \Delta_{j-1}$, where $0 \leq j \leq n$ and Δ_j is the j -skeleton of Δ . Any face of Δ is a stratified subspace.
- (3) For every compact stratified space L the *cone of L* is the quotient space

$c(L) = L \times [0, \infty) / L \times \{0\}$. We write $[p, r]$ for the equivalence class of $(p, r) \in L \times [0, \infty)$, the symbol \star denotes the equivalence class $L \times \{0\}$ called the *vertex*. By convention $c(\emptyset) = \{\star\}$. The radius is the function $\rho : c(L) \rightarrow [0, \infty)$ defined by $\rho[p, r] = r$. For every $\varepsilon > 0$ we write $c_\varepsilon(L) = \rho^{-1}[0, \varepsilon)$. The canonical stratification of the cone is the family

$$\mathcal{S}_{c(L)} = \{\star\} \sqcup \{S \times \mathbb{R}^+ : S \text{ is a stratum of } L\}.$$

Every $c_\varepsilon(L)$ is a stratified subspace of $c(L)$. A *basic model* is a product $M \times c(L)$ of a contractible manifold M with the cone.

LEMMA 1.3. *If \mathcal{S} is a stratification of X then*

- (1) *The relationship $S \leq S' \Leftrightarrow S \subset \overline{S'}$ is a partial order on \mathcal{S} .*
- (2) *For any stratum S the family of strata lying on S is finite.*
- (3) *A stratum is maximal (resp. minimal) \Leftrightarrow it is open (resp. closed).*
- (4) *The regular part is a dense open subset.*
- (5) *If X is compact then it has a finite number of strata.*

By the above lemma, the union of closed strata is called the *minimal part* of X , which we write Σ_{min} .

Since the stratification is locally finite, strict order chains

$$S_0 < S_1 < \dots < S_p$$

are always finite. The *length* of X is the maximal p for which there exists such an order chain, we write it $\text{len}(X)$. For instance, $\text{len}(X) = 0$ if and only if X is a manifold with the canonical stratification.

A function $f : X \rightarrow X'$ between two stratified spaces is a *morphism* (resp. *isomorphism*) if

- (1) f is a continuous function (resp. an homeomorphism).
- (2) f preserves the regular part: $f(X - \Sigma) \subset (X' - \Sigma')$.
- (3) f sends smoothly (resp. diffeomorphically) strata into strata.

For instance the change of radius $f_\varepsilon : c(L) \rightarrow c_\varepsilon(L)$ defined by $f_\varepsilon[p, r] = [p, 2\varepsilon \cdot \arctan(r)/\pi]$ is an isomorphism. A morphism f is an *immersion* when $f(X)$ is a stratified subspace of X' and f is an isomorphism from X onto $f(X)$. An *embedding* is an immersion whose image is open.

A stratified pseudomanifold is a stratified space having a conical behavior near the singular part, transversally to the singular strata.

DEFINITION 1.4. Fix a stratified space X (and a stratification \mathcal{S}), take a stratum S and a point $x \in S$. A *chart* in x is the embedding of a basic model

$$\varphi : U \times c(L) \rightarrow X$$

such that $U \subset S$ is a contractible open neighborhood of x and $\varphi(u, \star) = u$ for each $u \in U$. The image $\text{Im}(\varphi)$ is a *distinguished neighborhood* of x . The compact stratified space L is the *link* of the chart. Notice that $\text{len}(L) < \text{len}(X)$. An *atlas* of S is a family of charts with the same link $\mathcal{A}_S = \{\alpha : U_\alpha \times c(L) \rightarrow X\}_\alpha$ such that $\{U_\alpha\}_\alpha$ is a covering of S . We say that X is a *stratified pseudomanifold* if every stratum has an atlas whose link is a stratified pseudomanifold itself. In that case, an *atlas* of X is the choice of an atlas for each stratum.

EXAMPLES 1.5. These are the three examples of stratified pseudomanifolds the most frequently used in this work:

- (1) Every manifold (with the canonical stratification) is a stratified pseudomanifold, the link of any stratum being the empty set. For every manifold M and every stratified pseudomanifold X the product $M \times X$ is a stratified pseudomanifold.
- (2) If L is a compact stratified pseudomanifold then the open cone $c(L)$ is a stratified pseudomanifold. The link of the vertex is L .
- (3) The canonical n -simplex $\Delta \subset \mathbb{R}^{n+1}$ is a compact stratified pseudomanifold, the links are faces of Δ .

LEMMA 1.6. *If X is a stratified pseudomanifold then*

- (1) *Any two strata $S < S'$ satisfy $\dim(S) < \dim(S')$.*
- (2) *The family of distinguished neighborhoods is a base of the topology of X .*
- (3) *Every open subset of X is itself a stratified pseudomanifold.*

2. NORMALIZATIONS AND NORMALIZERS

From now on we fix a stratified pseudomanifold X and an atlas of X . In this section we will provide a detailed construction of the normalization of X . A stratified pseudomanifold is *normal* if its links are connected. For instance the canonical n -simplex $\Delta \subset \mathbb{R}^{n+1}$ is a normal stratified pseudomanifold. If X is a normal stratified pseudomanifold then any open subset $A \subset X$ is also normal, and any link of X is normal. The following result is straightforward.

LEMMA 2.1. *Each connected normal stratified pseudomanifold has only one regular stratum.*

The normalization of X is made up by cutting along the singular strata.

DEFINITION 2.2. A *normalization* of a stratified pseudomanifold X is a proper surjective morphism

$$\mathbf{n} : X^N \rightarrow X$$

from a normal stratified pseudomanifold X^N to X , together with a family of normalizations of the links $\{\mathbf{n}_L : L^N \rightarrow L\}_L$ (of some fixed atlas) satisfying

- (1) The restriction $\mathbf{n} : (X^N - \Sigma) \rightarrow (X - \Sigma)$ is an isomorphism.
- (2) For each singular point z of X^N there is a commutative diagram

$$\begin{array}{ccc} U \times c(L)^N & \xrightarrow{\varphi^N} & X^N \\ \downarrow \mathbf{n}_0 & & \downarrow \mathbf{n} \\ U \times c(L) & \xrightarrow{\varphi} & X \end{array} \tag{1}$$

satisfying

- (a) φ is a chart of $\mathbf{n}(z)$.
- (b) $c(L)^N = \sqcup_j c(K_j)$ where K_1, \dots, K_m are the connected components of L^N .
- (c) φ^N is an embedding and $\text{Im}(\varphi^N) = \mathbf{n}^{-1}(\text{Im}(\varphi))$.
- (d) $\mathbf{n}_0(u, [p, r]_j) = (u, [\mathbf{n}_L(p), r])$ where $[p, r]_j \in c(K_j)$.

In the above situation we will say X^N is a *normalizer* of X .

EXAMPLES 2.3. These are three easy examples of normalizations:

- (1) For any normal stratified pseudomanifold Z the identity 1_Z is a normalization.
- (2) Left vertical arrow \mathbf{n}_0 of diagram (1) is a normalization.
- (3) Fix a normalization $\mathbf{n} : X^N \rightarrow X$. Then for every open subset A of X the restriction $\mathbf{n} : \mathbf{n}^{-1}(A) \rightarrow A$ is a normalization of A . If X' is another stratified pseudomanifold and $f : X \rightarrow X'$ is an isomorphism then the composition $f\mathbf{n}$ is a normalization of X' . Finally, for every manifold M the map

$$1_M \times \mathbf{n} : M \times X^N \rightarrow M \times X$$

is a normalization of $M \times X$.

The stratification of a normalizer can be written in terms of the starting stratified pseudomanifold.

PROPOSITION 2.4. *If $\mathfrak{n} : X^N \rightarrow X$ is a normalization then*

- (1) *For each stratum S the restriction $\mathfrak{n} : \mathfrak{n}^{-1}(S) \rightarrow S$ is a locally trivial finite covering.*
- (2) *Every stratum of X^N is a connected component of $\mathfrak{n}^{-1}(S)$ for some stratum S of X .*
- (3) *$\mathfrak{n}^{-1}(\Sigma_{min}) = \Sigma_{min}$ and $\text{len}(X^N) = \text{len}(X)$.*

Proof. (1) It follows directly from § 2.2. (2) Fix a stratum S of X . Since any two strata $R, R' \subset \mathfrak{n}^{-1}(S)$ have the same dimension of S , they cannot be compared. Consequently $\mathfrak{n}^{-1}(S) \cap \overline{R} = R$; so R is a closed, connected and codimensional submanifold of $\mathfrak{n}^{-1}(S)$. Hence it is a connected component.

(3) In order to prove the first equality we first notice that

(a) *If S is a stratum and $R \subset \mathfrak{n}^{-1}(S)$ is a stratum then $\mathfrak{n}(R) = S$:* By step (1) of this proof and the fact that \mathfrak{n} is proper; it follows that $\mathfrak{n}(R)$ is a closed, connected and codimensional submanifold of S .

(b) *Strata contained in $\mathfrak{n}^{-1}(\Sigma_{min})$ are not comparable:* It follows from step (2), and the fact that minimal strata in X are not comparable (they are disjoint closed subsets).

Next fix a stratum R of X^N . By § 1.3 it suffices to show that that R is closed $\Leftrightarrow S = \mathfrak{n}(R)$ is closed. Then \Leftarrow follows from step (2) and the continuity of \mathfrak{n} . By the other hand, since \mathfrak{n} is a proper map; the converse \Rightarrow follows from step (a). This proves the first equality. The second equality is straightforward, it can be deduce from the first one by an inductive argument. ■

In order to show the uniqueness of the normalization we will establish its functoriality.

PROPOSITION 2.5. (The lifting property) *Let $\mathfrak{n} : Y^N \rightarrow Y$ be another normalization, $f : X \rightarrow Y$ a continuous map preserving the regular part. Then there is a unique continuous function f^N making commutative the following diagram*

$$\begin{array}{ccc} X^N & \xrightarrow{f^N} & Y^N \\ \downarrow \mathfrak{n} & & \downarrow \mathfrak{n} \\ X & \xrightarrow{f} & Y \end{array}$$

Proof. If there exists such a map f^N , then in $X^N - \Sigma$ it satisfies

$$f^N = \mathfrak{n}^{-1} f \mathfrak{n}$$

So f^N is unique because $X^N - \Sigma$ is an open dense. The above equation shows that we only need to define f^N in the singular part. Fix a singular point z of X^N ; we will say that $f^N(z) = v$ if there exists a sequence $\{z_j\} \subset X^N - \Sigma$ converging to z , such that $\{f^N(z_j)\}_j$ converges to v . Such a v always exists because $X^N - \Sigma$ is an open dense and the normalizations are proper maps. Notice that, by definition

$$\mathfrak{n}(v) = \mathfrak{n} \left(\lim_j f^N(z_j) \right) = \lim_j \mathfrak{n} f^N(z_j) = \lim_j f \mathfrak{n}(z_j) = y \tag{2}$$

Consequently, $v \in \mathfrak{n}^{-1}(y)$. Then:

- *The lifting is well defined:* Since this is a local matter, we can suppose that X^N is connected and little. By §2.1 X^N has a unique regular stratum R (and so does X). Take a chart $\psi : V \times c(L) \rightarrow Y$ of $y = f \mathfrak{n}(z)$ as in diagram (1). Assume that $f(X) \subset \text{Im}(\psi)$, so we get a diagram

$$\begin{array}{ccccc} X^N & \xrightarrow{f^N} & Y^N & \xleftarrow{\psi^N} & V \times c(L)^N \\ \downarrow \mathfrak{n} & & \downarrow \mathfrak{n} & & \downarrow \mathfrak{n}^0 \\ X & \xrightarrow{f} & Y & \xleftarrow{\psi} & V \times c(L) \end{array}$$

where the dashed arrow f^N is well defined in the regular part. So the composition $(\psi^N)^{-1} f^N$ sends R in a regular stratum $U \times R' \times \mathbb{R}^+ \subset U \times c(L)^N$. Notice that $\overline{R'} = K_j$ is a connected component of L^N . So, for any sequence $\{z_i\}_i \subset R$ converging to z the sequence $\{(\psi^N)^{-1} f^N(z_i)\}_i$ is contained in $U \times c(K_j)$. Hence

$$\lim_i (\psi^N)^{-1} f^N(z_i) = (f \mathfrak{n}(z), \star_j) \in U \times \{\star_j\}$$

This implies that $f^N(z) = \psi^N(f \mathfrak{n}(z), \star_j)$ is well defined.

- *The lifting is continuous:* This is easily seen by taking limits. ■

THEOREM 2.6. (Functoriality) *In the same situation of § 2.5, if f is a morphism (resp. an embedding or an isomorphism) then so is f^N .*

Proof. We consider three cases:

(a) *f is a morphism:* Since f^N is continuous and $\mathfrak{n} f^N = f \mathfrak{n}$, by §2.4, it sends strata into strata. Hence f^N is a morphism.

(b) *f is an isomorphism*: The inverse morphism f^{-1} lifts to a unique morphism $g : Y^N \rightarrow X^N$ such that $\mathbf{n}g = f^{-1}\mathbf{n}$. Notice that

$$\mathbf{n}gf^N = f^{-1}\mathbf{n}f^N = f^{-1}f\mathbf{n} = 1_Y\mathbf{n}$$

so gf^N is a lifting of the identity 1_Y . Since the lifting is unique $gf^N = 1_Y^N$ is the identity of Y^N . With the same argument $f^Ng = 1_X^N$ is the identity of X^N , thus g is the inverse of f^N .

(c) *f is an embedding*: Notice that the restriction $\mathbf{n} : \mathbf{n}^{-1}(f(X)) \rightarrow f(X)$ is a normalization, because $f(X)$ is open. Now apply step (b) to the isomorphism $f : X \rightarrow f(X)$. ■

COROLLARY 2.7. *Normalizations and normalizers are unique up to isomorphisms.*

Now we will prove the existence of a normalization of X .

THEOREM 2.8. *Each stratified pseudomanifold has a normalization.*

Proof. Fix a stratified pseudomanifold X and an atlas.

• *Reduction to a local matter*: Proceed by induction on $p = \text{len}(X)$. For $p = 0$ it is trivial, assume the inductive hypothesis. Then each link of X has a normalization and for each chart $\varphi : U \times c(L) \rightarrow X$ the composition $\varphi\mathbf{n}_0$ is a normalization of $\text{Im}(\varphi)$, where \mathbf{n}_0 is the map given in § 2.2. The family

$$\mathcal{U} = \{A \subset X : A \text{ is open and has a normalizer}\}$$

is a basis of the topology of X (see § 1.6), and it is closed by finite intersections (see § 2.3) and by arbitrary disjoint unions. By a Mayer-Vietoris argument known as the *Bredon's Trick* [1], it suffices to consider the case when $X = A \cup B$ for two open subsets A, B having normalizations

$$\mathbf{n}_A : A^N \rightarrow A \quad \mathbf{n}_B : B^N \rightarrow B$$

By Theorem 2.6 there is an isomorphism ϕ such that the following diagram commutes

$$\begin{array}{ccc} \mathbf{n}_A^{-1}(A \cap B) & \xrightarrow{\phi} & \mathbf{n}_B^{-1}(A \cap B) \\ \downarrow \mathbf{n}_A & & \downarrow \mathbf{n}_B \\ A \cap B & \xrightarrow{1} & A \cap B \end{array} \quad \mathbf{n}_B\phi = \mathbf{n}_A \quad (3)$$

- *Construction of the normalizer:* Define X^N as the sum of A^N and B^N amalgamated by the isomorphism ϕ , i.e.;

$$X^N = A^N \sqcup B^N / \sim \quad z \sim \phi(z) \quad \forall z \in \mathbf{n}_A^{-1}(A \cap B) \quad (4)$$

We write $[z]$ for the equivalence class of $z \in A^N \sqcup B^N$. Endow X^N with the quotient topology induced by the canonical projection $q : z \mapsto [z]$. Since ϕ is an isomorphism it's easy to see that X^N is a stratified space and q is a surjective morphism.

- *Embedding of A^N and B^N in X^N :* The arrow $q : A^N \rightarrow X^N$ is continuous and injective by definition. For every subset $R \subset A^N$ we have

$$q^{-1}q(R) = R \sqcup \phi(R \cap \mathbf{n}_A^{-1}(B))$$

If R is open then so is $q(R)$, which implies $q|_{A^N}$ is open. By the other hand, if R is a stratum then q sends it into a stratum of X^N and $q : R \rightarrow q(R)$ is a homeomorphism. Hence, $q|_{A^N}$ is an embedding; the same holds for $q|_{B^N}$.

- *Definition of the normalization:* Define $\mathbf{n} : X^N \rightarrow X$ by the rule

$$\mathbf{n}[z] = \begin{cases} \mathbf{n}_A(z) & \text{if } z \in A^N \\ \mathbf{n}_B(z) & \text{if } z \in B^N \end{cases} \quad (5)$$

Then \mathbf{n} is well defined by diagram (3); besides it's continuous and surjective. We obtain a commutative diagram

$$\begin{array}{ccc} X^N & \xrightarrow{q} & A^N \sqcup B^N \\ \mathbf{n} \downarrow & & \nearrow \mathbf{n}_A \sqcup \mathbf{n}_B \\ X & & \end{array}$$

where q and $\mathbf{n}_A \sqcup \mathbf{n}_B$ are morphisms, then so is \mathbf{n} . It's easy to see that restricted to the regular part \mathbf{n} is a diffeomorphism, so we verify the other properties of § 2.2.

- *Local triviality:* For every point $z \in X^N$ its image $x = \mathbf{n}(z)$ lies in A or in B . Suppose $x \in A$ and let $\varphi : U \times c(L) \rightarrow A$ a chart of x . We get a commutative diagram

$$\begin{array}{ccccc} U \times c(L)^N & \xrightarrow{\varphi^N} & A^N & \xrightarrow{q} & X^N \\ \downarrow 1 \times \mathbf{n}_L & & \downarrow \mathbf{n}_A & & \mathbf{n} \\ U \times c(L) & \xrightarrow{\varphi} & A^N & \hookrightarrow & X \end{array}$$

By restricting $q\varphi^N$ to the connected component of $U \times c(L)^N$ which contains $(\varphi^N)^{-1}(z)$ we get a chart of z .

From the above discussion it is straightforward that \mathbf{n} is a proper map and X^N is a normal stratified pseudomanifold. ■

Remark 2.9. If we ask \mathbf{n} to be smooth on each stratum and a diffeomorphism in the regular part; then Theorems 1 and 2 still hold in the C^∞ category.

EXAMPLE 2.10. *Normalizing a tubular neighborhood:* A tubular neighborhood is a locally trivial fiber bundle $\xi = (T, \tau, S, c(L))$ where

- (1) T is a stratified pseudomanifold, $S \subset T$ is the unique minimal stratum. The fiber $c(L)$ is the cone of a compact stratified pseudomanifold.
- (2) $\tau : T \rightarrow S$ is a morphism, the restriction $\tau|_S$ is the identity of S .
- (3) The structure group of the fiber bundle is $G = Iso(L)$. In other words, there is a trivializing atlas

$$\mathcal{U} = \{\alpha : U_\alpha \times c(L) \rightarrow \tau^{-1}(U_\alpha)\}_\alpha$$

such that, if $U_\alpha \cap U_\beta \neq \emptyset$, then the change of charts satisfies

$$\beta\alpha^{-1} : U_\alpha \cap U_\beta \times c(L) \rightarrow U_\alpha \cap U_\beta \times c(L) \quad (x, [p, r]) \mapsto (x, [g_{\alpha\beta}(x, p), r])$$

Notice that $g_{\alpha\beta}(x, -)$ is an isomorphism and the global extension of the radius

$$\rho : T \rightarrow [0, \infty) \quad \rho\alpha(x, [p, r]) = r$$

is a stratified morphism which makes sense.

Fix two normalizations $\mathbf{n} : T^N \rightarrow T$ and $\mathbf{n}_L : L^N \rightarrow L$; take $\mathbf{n}_0 : c(L)^N \rightarrow c(L)$ as in § 2.2. We will show that there is a fiber bundle

$$\xi^N = (T^N, \tau^N, \mathbf{n}^{-1}(S), c(L)^N)$$

where each object is induced by the process of normalization and the structure group is $G = Iso(L^N)$. But this is immediate since, by Theorem 2.6, there is a unique lifting $\tau^N : T^N \rightarrow \mathbf{n}^{-1}(S)$ such that the following diagram commutes

$$\begin{array}{ccc} T^N & \xrightarrow{\tau^N} & \mathbf{n}^{-1}(S) \\ \mathbf{n} \downarrow & & \downarrow \mathbf{n} \\ T & \xrightarrow{\tau} & S \end{array}$$

Each trivialization α in the atlas \mathcal{U} lifts to a trivialization

$$\alpha^N : U_\alpha \times c(L)^N \rightarrow (\tau^N)^{-1}(\mathfrak{n}^{-1}(U))$$

By the existence and uniqueness of liftings, the family of cocycles $\{g_{\alpha\beta}\}$ lifts to a family $\{g_{\alpha\beta}^N\}$ satisfying $g_{\alpha\beta}^N g_{\beta\delta}^N = g_{\alpha\delta}^N$ for all α, β, δ ; thus it is again a family of cocycles. It is straightforward that the radium ρ lifts in a consistent way to a radium ρ^N .

3. INTERSECTION HOMOLOGY OF THE NORMALIZER

Intersection homology was first introduced by Goresky and MacPherson in the category of pl -stratified pseudomanifolds, and later extended for any stratified pseudomanifold [2], [3]. In order to show that the normalization preserves the intersection homology, we extend Borel’s procedure to the family of perversities given in [7]; see also [4], [5], [9]. Those perversities are more general than the usual ones.

DEFINITION 3.1. Fix a stratified pseudomanifold X . A *perversity* in X is a function $\bar{p} : \mathcal{S} \rightarrow \mathbb{Z}$ from the family of strata \mathcal{S} to the integers. A singular simplex $\sigma : \Delta \rightarrow X$ is \bar{p} -*admissible* if it satisfies the following properties:

- (1) σ sends the interior of Δ in $X - \Sigma$.
- (2) $\sigma^{-1}(S) \subset (\dim(\Delta) - \text{codim}(S) + \bar{p}(S))$ -skeleton of Δ , for each singular stratum S of X .

A singular chain $\xi = \sum_{j=1}^m r_j \sigma_j$ is \bar{p} -*admissible* if every σ_j is \bar{p} -admissible. We will say that ξ is a \bar{p} -*chain* if ξ and its boundary $\partial\xi$ are both \bar{p} -admissible. We write $SC_*^{\bar{p}}(X)$ for the complex of \bar{p} -chains. The \bar{p} -*intersection homology* of X is the homology $H_*^{\bar{p}}(X)$ of the complex $SC_*^{\bar{p}}(X)$.

Given a normalization $\mathfrak{n} : X^N \rightarrow X$, each perversity \bar{p} in X induces trivially a perversity in X^N which we write again \bar{p} by abuse of language.

THEOREM 3.2. $H_*^{\bar{p}}(X^N) = H_*^{\bar{p}}(X)$ for any perversity \bar{p} in X .

Proof. We claim that the usual chain morphism $\mathfrak{n}_* : SC_*^{\bar{p}}(X^N) \rightarrow SC_*^{\bar{p}}(X)$ is an isomorphism. It suffice to show it in the \bar{p} -simplexes.

- *The arrow is well defined:* We will proof that if σ is a simplex \bar{p} -admissible in X^N then $\mathfrak{n}\sigma$ is \bar{p} -admissible in X . Condition (1) is trivial, so we verify (2):

Fix a singular stratum S of X . Let S' be a stratum contained in $\mathbf{n}^{-1}(S)$. Then $\sigma^{-1}(S') \subset \dim(\Delta) - \text{codim}(S') + \bar{p}(S')$ -skeleton of Δ . Since $\dim(S) = \dim(S')$ then we conclude that $(\mathbf{n}\sigma)^{-1}(S) \subset (\dim(\Delta) - \text{codim}(S) + \bar{p}(S))$ -skeleton of Δ .

- *The arrow is injective:* Take two singular simplexes $\sigma, \sigma' : \Delta \rightarrow X^N$ and suppose $\mathbf{n}\sigma = \mathbf{n}\sigma'$; then σ and σ' coincide in the interior of Δ which implies that $\sigma = \sigma'$.

- *The arrow is surjective:* Recall that Δ is a normal stratified pseudomanifold, thus the identity $1_\Delta : \Delta \rightarrow \Delta$ is a normalization. Take a \bar{p} -admissible singular simplex $\sigma : \Delta \rightarrow X$. By §2.5; lifts to a unique singular simplex $\sigma^N : \Delta \rightarrow X^N$ satisfying $\mathbf{n}\sigma^N = \sigma$. Conditions (1) and (2) of §3.1 are easily verified for σ^N ; so it is \bar{p} -admissible. ■

ACKNOWLEDGEMENTS

I would like to thank M. Saralegi for encouraging me to look at this problem, and also R. Popper for some helpful conversations.

REFERENCES

- [1] BREDON, G.E., "Topology and Geometry", Graduate Texts in Mathematics 139, Springer-Verlag, New York, 1993.
- [2] GORESKY, M., MACPHERSON, R., Intersection homology theory, *Topology*, **19** (1980), 135–162.
- [3] GORESKY, M., MACPHERSON, R., Intersection homology II, *Invent. Math.*, **71** (1983), 77–129.
- [4] HAEFLIGER, A., Introduction to piecewise linear intersection homology, in "Intersection Cohomology" (Bern, 1983), *Progr. Math.*, **50**, 1–22, Birkhäuser, Boston, MA, 1984.
- [5] KING, H.C., Topology invariance of intersection homology without sheaves, *Topology Appl.*, **20** (2) (1985), 149–160.
- [6] MACCRORY, C., "Poincaré Duality in Spaces with Singularities", Ph.D. Thesis, Brandeis University, 1972.
- [7] MACPHERSON, R., "Intersection Homology and Perverse Sheaves", Colloquium Lectures, Annual Meeting of the AMS, San Francisco, June 1991.
- [8] PFLAUM, M.J., "Analytic and Geometric Study of Stratified Spaces", Lecture Notes in Mathematics 1768, Springer-Verlag, Berlin, 2001.
- [9] SARALEGI, M., Homological properties of stratified spaces, *Illinois J. Math.*, **38** (1994), 47–70.