

Ostrowski Type Fractional Integral Inequalities for Generalized (g, s, m, φ) -Preinvex Functions

ARTION KASHURI, ROZANA LIKO

*Department of Mathematics, Faculty of Technical Science,
University "Ismail Qemali", Vloza, Albania*

artionkashuri@gmail.com rozanaliko86@gmail.com

Presented by Horst Martini

Received September 9, 2016

Abstract: In the present paper, a new class of generalized (g, s, m, φ) -preinvex function is introduced and some new integral inequalities for the left hand side of Gauss-Jacobi type quadrature formula involving generalized (g, s, m, φ) -preinvex functions are given. Moreover, some generalizations of Ostrowski type inequalities for generalized (g, s, m, φ) -preinvex functions via Riemann-Liouville fractional integrals are established. At the end, some applications to special means are given.

Key words: Ostrowski type inequality, Hölder's inequality, power mean inequality, Riemann-Liouville fractional integral, s -convex function in the second sense, m -invex, P -function.

AMS Subject Class. (2010): 26A51, 26A33, 26D07, 26D10, 26D15.

1. INTRODUCTION AND PRELIMINARIES

The following notation is used throughout this paper. We use I to denote an interval on the real line $\mathbb{R} = (-\infty, +\infty)$ and I° to denote the interior of I . For any subset $K \subseteq \mathbb{R}^n$, K° is used to denote the interior of K . \mathbb{R}^n is used to denote a generic n -dimensional vector space. The nonnegative real numbers are denoted by $\mathbb{R}_0 = [0, +\infty)$. The set of integrable functions on the interval $[a, b]$ is denoted by $L_1[a, b]$.

The following result is known in the literature as the Ostrowski inequality (see [11]) and the references cited therein, which gives an upper bound for the approximation of the integral average $\frac{1}{b-a} \int_a^b f(t)dt$ by the value $f(x)$ at point $x \in [a, b]$.

THEOREM 1.1. *Let $f : I \rightarrow \mathbb{R}$, where $I \subseteq \mathbb{R}$ is an interval, be a mapping differentiable in the interior I° of I , and let $a, b \in I^\circ$ with $a < b$. If $|f'(x)| \leq M$ for all $x \in [a, b]$, then*

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq M(b-a) \left[\frac{1}{4} + \frac{\left(x - \frac{a+b}{2}\right)^2}{(b-a)^2} \right], \quad \forall x \in [a, b]. \quad (1.1)$$

For other recent results concerning Ostrowski type inequalities (see [11]) and the references cited therein, also (see [12]) and the references cited therein.

Fractional calculus (see [10]) and the references cited therein, was introduced at the end of the nineteenth century by Liouville and Riemann, the subject of which has become a rapidly growing area and has found applications in diverse fields ranging from physical sciences and engineering to biological sciences and economics.

DEFINITION 1.2. Let $f \in L_1[a, b]$. The Riemann-Liouville integrals $J_{a+}^\alpha f$ and $J_{b-}^\alpha f$ of order $\alpha > 0$ with $a \geq 0$ are defined by

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \quad x > a$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, \quad b > x,$$

where $\Gamma(\alpha) = \int_0^{+\infty} e^{-u} u^{\alpha-1} du$. Here $J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x)$.

In the case of $\alpha = 1$, the fractional integral reduces to the classical integral.

Due to the wide application of fractional integrals, some authors extended to study fractional Ostrowski type inequalities for functions of different classes (see [10]) and the references cited therein.

Now, let us recall some definitions of various convex functions.

DEFINITION 1.3. (see [2]) A nonnegative function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}_o$ is said to be P -function or P -convex, if

$$f(tx + (1-t)y) \leq f(x) + f(y), \quad \forall x, y \in I, t \in [0, 1].$$

DEFINITION 1.4. (see [3]) A function $f : \mathbb{R}_o \rightarrow \mathbb{R}$ is said to be s -convex in the second sense, if

$$f(\lambda x + (1-\lambda)y) \leq \lambda^s f(x) + (1-\lambda)^s f(y) \quad (1.2)$$

for all $x, y \in \mathbb{R}_o$, $\lambda \in [0, 1]$ and $s \in (0, 1]$.

It is clear that a 1-convex function must be convex on \mathbb{R}_o as usual. The s -convex functions in the second sense have been investigated in (see [3]).

DEFINITION 1.5. (see [4]) A set $K \subseteq \mathbb{R}^n$ is said to be invex with respect to the mapping $\eta : K \times K \rightarrow \mathbb{R}^n$, if $x + t\eta(y, x) \in K$ for every $x, y \in K$ and $t \in [0, 1]$.

Notice that every convex set is invex with respect to the mapping $\eta(y, x) = y - x$, but the converse is not necessarily true. For more details please see (see [4], [5]) and the references therein.

DEFINITION 1.6. (see [6]) The function f defined on the invex set $K \subseteq \mathbb{R}^n$ is said to be preinvex with respect η , if for every $x, y \in K$ and $t \in [0, 1]$, we have that

$$f(x + t\eta(y, x)) \leq (1 - t)f(x) + tf(y).$$

The concept of preinvexity is more general than convexity since every convex function is preinvex with respect to the mapping $\eta(y, x) = y - x$, but the converse is not true.

The Gauss-Jacobi type quadrature formula has the following

$$\int_a^b (x - a)^p (b - x)^q f(x) dx = \sum_{k=0}^{+\infty} B_{m,k} f(\gamma_k) + R_m^* |f|, \quad (1.3)$$

for certain $B_{m,k}, \gamma_k$ and rest $R_m^* |f|$ (see [7]).

Recently, Liu (see [8]) obtained several integral inequalities for the left hand side of (1.3) under the Definition 1.3 of P -function. Also in (see [9]), Özdemir et al. established several integral inequalities concerning the left-hand side of (1.3) via some kinds of convexity. Motivated by these results, in Section 2, the notion of generalized (g, s, m, φ) -preinvex function is introduced and some new integral inequalities for the left hand side of (1.3) involving generalized (g, s, m, φ) -preinvex functions are given. In Section 3, some generalizations of Ostrowski type inequalities for generalized (g, s, m, φ) -preinvex functions via fractional integrals are given. In Section 4, some applications to special means are given.

2. NEW INTEGRAL INEQUALITIES FOR GENERALIZED (g, s, m, φ) -PREINVEX FUNCTIONS

DEFINITION 2.1. (see [1]) A set $K \subseteq \mathbb{R}^n$ is said to be m -invex with respect to the mapping $\eta : K \times K \times (0, 1] \rightarrow \mathbb{R}^n$ for some fixed $m \in (0, 1]$, if

$m\eta(y, x, m) \in K$ holds for each $x, y \in K$ and any $t \in [0, 1]$.

Remark 2.2. In Definition 2.1, under certain conditions, the mapping $\eta(y, x, m)$ could reduce to $\eta(y, x)$.

We next give new definition, to be referred as generalized (g, s, m, φ) -preinvex function.

DEFINITION 2.3. Let $K \subseteq \mathbb{R}^n$ be an open m -invex set with respect to $\eta : K \times K \times (0, 1] \rightarrow \mathbb{R}^n$, $g : [0, 1] \rightarrow [0, 1]$ be a differentiable function and $\varphi : I \rightarrow \mathbb{R}$ is a continuous increasing function. For $f : K \rightarrow \mathbb{R}$ and any fixed $s, m \in (0, 1]$, if

$$f(m\varphi(x) + g(t)\eta(\varphi(y), \varphi(x), m)) \leq m(1 - g(t))^s f(\varphi(x)) + g^s(t)f(\varphi(y)) \quad (2.1)$$

is valid for all $x, y \in K, t \in [0, 1]$, then we say that f is a generalized (g, s, m, φ) -preinvex function with respect to η .

Remark 2.4. In Definition 2.3, it is worthwhile to note that the class of generalized (g, s, m, φ) -preinvex function is a generalization of the class of s -convex in the second sense function given in Definition 1.4. Also, for $g(t) = \lambda, \lambda \in [0, 1]$ and $\varphi(x) = x, \forall x \in K$, we get the notion of generalized (s, m) -preinvex function (see [1]).

EXAMPLE 2.5. Let $f(x) = -|x|, g(t) = t, \varphi(x) = x, s = 1$ and

$$\eta(y, x, m) = \begin{cases} y - mx, & \text{if } x \geq 0, y \geq 0; \\ y - mx, & \text{if } x \leq 0, y \leq 0; \\ mx - y, & \text{if } x \geq 0, y \leq 0; \\ mx - y, & \text{if } x \leq 0, y \geq 0. \end{cases}$$

Then $f(x)$ is a generalized $(t, 1, m, x)$ -preinvex function of with respect to $\eta : \mathbb{R} \times \mathbb{R} \times (0, 1] \rightarrow \mathbb{R}$ and any fixed $m \in (0, 1]$. However, it is obvious that $f(x) = -|x|$ is not a convex function on \mathbb{R} .

In this section, in order to prove our main results regarding some new integral inequalities involving generalized (g, s, m, φ) -preinvex functions, we need the following new lemma:

LEMMA 2.6. Let $\varphi : I \rightarrow \mathbb{R}$ be a continuous increasing function and $g : [0, 1] \rightarrow [0, 1]$ is a differentiable function. Assume that

$$f : K = [m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)] \rightarrow \mathbb{R}$$

is a continuous function on the interval of real numbers K° with respect to $\eta : K \times K \times (0, 1] \rightarrow \mathbb{R}$, for $\varphi(a), \varphi(b) \in K$, $a < b$ and $m\varphi(a) < m\varphi(a) + \eta(\varphi(b), \varphi(a), m)$. Then for any fixed $m \in (0, 1]$ and $p, q > 0$, we have

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b),\varphi(a),m)} (x - m\varphi(a))^p \left(m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x \right)^q f(x) dx \\ &= \eta(\varphi(b), \varphi(a), m)^{p+q+1} \int_0^1 g^p(t) (1 - g(t))^q \\ & \quad \times f\left(m\varphi(a) + g(t)\eta(\varphi(b), \varphi(a), m)\right) d[g(t)]. \end{aligned}$$

Proof. It is easy to observe that

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b),\varphi(a),m)} (x - m\varphi(a))^p \left(m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x \right)^q f(x) dx \\ &= \eta(\varphi(b), \varphi(a), m) \int_0^1 \left(m\varphi(a) + g(t)\eta(\varphi(b), \varphi(a), m) - m\varphi(a) \right)^p \\ & \quad \times \left(m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - m\varphi(a) - g(t)\eta(\varphi(b), \varphi(a), m) \right)^q \\ & \quad \times f\left(m\varphi(a) + g(t)\eta(\varphi(b), \varphi(a), m)\right) d[g(t)] \\ &= \eta(\varphi(b), \varphi(a), m)^{p+q+1} \int_0^1 g^p(t) (1 - g(t))^q \\ & \quad \times f\left(m\varphi(a) + g(t)\eta(\varphi(b), \varphi(a), m)\right) d[g(t)]. \end{aligned}$$

■

The following definition will be used in the sequel.

DEFINITION 2.7. The Euler Beta function is defined for $x, y > 0$ as

$$\beta(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}.$$

THEOREM 2.8. Let $\varphi : I \rightarrow \mathbb{R}$ be a continuous increasing function and $g : [0, 1] \rightarrow [0, 1]$ is a differentiable function. Assume that $f : K = [m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)] \rightarrow \mathbb{R}$ is a continuous function on the interval of real numbers K° with $\varphi(a), \varphi(b) \in K$, $a < b$ with $m\varphi(a) < m\varphi(a) + \eta(\varphi(b), \varphi(a), m)$. Let $k > 1$. If $|f|^{\frac{k}{k-1}}$ is a generalized (g, s, m, φ) -preinvex function on an open m -invex set K with respect to $\eta : K \times K \times (0, 1] \rightarrow \mathbb{R}$ for

any fixed $s, m \in (0, 1]$, then for any fixed $p, q > 0$,

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b),\varphi(a),m)} (x - m\varphi(a))^p \left(m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x \right)^q f(x) dx \\ & \leq \frac{|\eta(\varphi(b), \varphi(a), m)|^{p+q+1}}{(s+1)^{\frac{k-1}{k}}} B^{\frac{1}{k}}(g(t); k, p, q) \\ & \quad \times \left[m \left((1 - g(0))^{s+1} - (1 - g(1))^{s+1} \right) |f(\varphi(a))|^{\frac{k}{k-1}} \right. \\ & \quad \left. + (g^{s+1}(1) - g^{s+1}(0)) |f(\varphi(b))|^{\frac{k}{k-1}} \right]^{\frac{k-1}{k}}, \end{aligned}$$

where $B(g(t); k, p, q) = \int_0^1 g^{kp}(t)(1 - g(t))^{kq} d[g(t)]$.

Proof. Since $|f|^{\frac{k}{k-1}}$ is a generalized (g, s, m, φ) -preinvex function on K , combining with Lemma 2.6 and Hölder inequality for all $t \in [0, 1]$ and for any fixed $s, m \in (0, 1]$, we get

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b),\varphi(a),m)} (x - m\varphi(a))^p \left(m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x \right)^q f(x) dx \\ & \leq |\eta(\varphi(b), \varphi(a), m)|^{p+q+1} \left[\int_0^1 g^{kp}(t)(1 - g(t))^{kq} d[g(t)] \right]^{\frac{1}{k}} \\ & \quad \times \left[\int_0^1 |f(m\varphi(a) + g(t)\eta(\varphi(b), \varphi(a), m))|^{\frac{k}{k-1}} d[g(t)] \right]^{\frac{k-1}{k}} \\ & \leq |\eta(\varphi(b), \varphi(a), m)|^{p+q+1} B^{\frac{1}{k}}(g(t); k, p, q) \\ & \quad \times \left[\int_0^1 (m(1 - g(t))^s |f(\varphi(a))|^{\frac{k}{k-1}} + g^s(t) |f(\varphi(b))|^{\frac{k}{k-1}}) d[g(t)] \right]^{\frac{k-1}{k}} \\ & = \frac{|\eta(\varphi(b), \varphi(a), m)|^{p+q+1}}{(s+1)^{\frac{k-1}{k}}} B^{\frac{1}{k}}(g(t); k, p, q) \\ & \quad \times \left[m \left((1 - g(0))^{s+1} - (1 - g(1))^{s+1} \right) |f(\varphi(a))|^{\frac{k}{k-1}} \right. \\ & \quad \left. + (g^{s+1}(1) - g^{s+1}(0)) |f(\varphi(b))|^{\frac{k}{k-1}} \right]^{\frac{k-1}{k}}. \end{aligned}$$

■

COROLLARY 2.9. Under the conditions of Theorem 2.8 for $g(t) = t$, we get

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b),\varphi(a),m)} (x - m\varphi(a))^p \left(m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x\right)^q f(x) dx \\ & \leq \frac{|\eta(\varphi(b), \varphi(a), m)|^{p+q+1}}{(s+1)^{\frac{k-1}{k}}} \left[\beta(kp+1, kq+1)\right]^{\frac{1}{k}} \\ & \quad \times \left(m|f(\varphi(a))|^{\frac{k}{k-1}} + |f(\varphi(b))|^{\frac{k}{k-1}}\right)^{\frac{k-1}{k}}. \end{aligned}$$

THEOREM 2.10. Let $\varphi : I \rightarrow \mathbb{R}$ be a continuous increasing function and $g : [0, 1] \rightarrow [0, 1]$ is a differentiable function. Assume that $f : K = [m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)] \rightarrow \mathbb{R}$ is a continuous function on the interval of real numbers K° with $\varphi(a), \varphi(b) \in K$, $a < b$ with $m\varphi(a) < m\varphi(a) + \eta(\varphi(b), \varphi(a), m)$. Let $l \geq 1$. If $|f|^l$ is a generalized (g, s, m, φ) -preinvex function on an open m -invex set K with respect to $\eta : K \times K \times (0, 1] \rightarrow \mathbb{R}$ for any fixed $s, m \in (0, 1]$, then for any fixed $p, q > 0$,

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b),\varphi(a),m)} (x - m\varphi(a))^p \left(m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x\right)^q f(x) dx \\ & \leq |\eta(\varphi(b), \varphi(a), m)|^{p+q+1} B^{\frac{l-1}{l}}(g(t); p, q) \\ & \quad \times \left[m|f(\varphi(a))|^l B(g(t); p, q+s) + |f(\varphi(b))|^l B(g(t); p+s, q) \right]^{\frac{1}{l}}, \end{aligned}$$

where $B(g(t); p, q) = \int_0^1 g^p(t)(1-g(t))^q d[g(t)]$.

Proof. Since $|f|^l$ is a generalized (s, m, φ) -preinvex function on K , combining with Lemma 2.6 and the well-known power mean inequality for all $t \in [0, 1]$ and for any fixed $s, m \in (0, 1]$, we get

$$\begin{aligned} & \int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b),\varphi(a),m)} (x - m\varphi(a))^p \left(m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x\right)^q f(x) dx \\ & = \eta(\varphi(b), \varphi(a), m)^{p+q+1} \int_0^1 \left[g^p(t)(1-g(t))^q \right]^{\frac{l-1}{l}} \left[g^p(t)(1-g(t))^q \right]^{\frac{1}{l}} \\ & \quad \times f\left(m\varphi(a) + g(t)\eta(\varphi(b), \varphi(a), m)\right) d[g(t)] \end{aligned}$$

$$\begin{aligned}
&\leq |\eta(\varphi(b), \varphi(a), m)|^{p+q+1} \left[\int_0^1 g^p(t)(1-g(t))^q d[g(t)] \right]^{\frac{l-1}{l}} \\
&\quad \times \left[\int_0^1 g^p(t)(1-g(t))^q \left| f\left(m\varphi(a) + g(t)\eta(\varphi(b), \varphi(a), m)\right) \right|^l d[g(t)] \right]^{\frac{1}{l}} \\
&\leq |\eta(\varphi(b), \varphi(a), m)|^{p+q+1} B^{\frac{l-1}{l}}(g(t); p, q) \\
&\quad \times \left[\int_0^1 g^p(t)(1-g(t))^q \left(m(1-g(t))^s |f(\varphi(a))|^l + g^s(t) |f(\varphi(b))|^l \right) d[g(t)] \right]^{\frac{1}{l}} \\
&= |\eta(\varphi(b), \varphi(a), m)|^{p+q+1} B^{\frac{l-1}{l}}(g(t); p, q) \\
&\quad \times \left[m |f(\varphi(a))|^l B(g(t); p, q+s) + |f(\varphi(b))|^l B(g(t); p+s, q) \right]^{\frac{1}{l}}.
\end{aligned}$$

■

COROLLARY 2.11. Under the conditions of Theorem 2.10 for $g(t) = t$, we get

$$\begin{aligned}
&\int_{m\varphi(a)}^{m\varphi(a)+\eta(\varphi(b), \varphi(a), m)} (x - m\varphi(a))^p \left(m\varphi(a) + \eta(\varphi(b), \varphi(a), m) - x \right)^q f(x) dx \\
&\leq |\eta(\varphi(b), \varphi(a), m)|^{p+q+1} \left[\beta(p+1, q+1) \right]^{\frac{l-1}{l}} \\
&\quad \times \left[m |f(\varphi(a))|^l \beta(p+1, q+s+1) + |f(\varphi(b))|^l \beta(p+s+1, q+1) \right]^{\frac{1}{l}}.
\end{aligned}$$

3. OSTROWSKI TYPE FRACTIONAL INTEGRAL INEQUALITIES FOR GENERALIZED (g, s, m, φ) -PREINVEX FUNCTIONS

In this section, in order to prove our main results regarding some generalizations of Ostrowski type inequalities for generalized (g, s, m, φ) -preinvex functions via fractional integrals, we need the following new lemma:

LEMMA 3.1. Let $\varphi : I \rightarrow \mathbb{R}$ be a continuous increasing function and $g : [0, 1] \rightarrow [0, 1]$ is a differentiable function. Suppose $K \subseteq \mathbb{R}$ be an open m -invex subset with respect to $\eta : K \times K \times (0, 1] \rightarrow \mathbb{R}$ for any fixed $m \in (0, 1]$ and let $\varphi(a), \varphi(b) \in K$, $a < b$ with $m\varphi(a) < m\varphi(a) + \eta(\varphi(b), \varphi(a), m)$. Assume

that $f : K \rightarrow \mathbb{R}$ is a differentiable function on K° and f' is integrable on $[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$. Then for $\alpha > 0$, we have

$$\begin{aligned} & \frac{\eta(\varphi(x), \varphi(a), m)^\alpha}{\eta(\varphi(b), \varphi(a), m)} \left[g^\alpha(1) f\left(m\varphi(a) + g(1)\eta(\varphi(x), \varphi(a), m)\right) \right. \\ & \quad \left. - g^\alpha(0) f\left(m\varphi(a) + g(0)\eta(\varphi(x), \varphi(a), m)\right) \right] \\ & - \frac{\eta(\varphi(x), \varphi(b), m)^\alpha}{\eta(\varphi(b), \varphi(a), m)} \left[g^\alpha(1) f\left(m\varphi(b) + g(1)\eta(\varphi(x), \varphi(b), m)\right) \right. \\ & \quad \left. - g^\alpha(0) f\left(m\varphi(b) + g(0)\eta(\varphi(x), \varphi(b), m)\right) \right] \tag{3.1} \\ & - \frac{\alpha}{\eta(\varphi(b), \varphi(a), m)} \left[\int_{m\varphi(a)+g(0)\eta(\varphi(x), \varphi(a), m)}^{m\varphi(a)+g(1)\eta(\varphi(x), \varphi(a), m)} (t - m\varphi(a))^{\alpha-1} f(t) dt \right. \\ & \quad \left. - \int_{m\varphi(b)+g(0)\eta(\varphi(x), \varphi(b), m)}^{m\varphi(b)+g(1)\eta(\varphi(x), \varphi(b), m)} (t - m\varphi(b))^{\alpha-1} f(t) dt \right] \\ & = \frac{\eta(\varphi(x), \varphi(a), m)^{\alpha+1}}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 g^\alpha(t) f'\left(m\varphi(a) + g(t)\eta(\varphi(x), \varphi(a), m)\right) d[g(t)] \\ & \quad - \frac{\eta(\varphi(x), \varphi(b), m)^{\alpha+1}}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 g^\alpha(t) f'\left(m\varphi(b) + g(t)\eta(\varphi(x), \varphi(b), m)\right) d[g(t)]. \end{aligned}$$

Proof. A simple proof of the equality can be done by performing an integration by parts in the integrals from the right side and changing the variable. The details are left to the interested reader. ■

Remark 3.2. Clearly, if we choose $m = 1$, $g(t) = t$, $\eta(\varphi(y), \varphi(x), m) = \varphi(y) - m\varphi(x)$ and $\varphi(x) = x$, $\forall x, y \in K$ in Lemma 3.1, we get Lemma 1 in [11].

Let denote

$$\begin{aligned} & S_{f,g,\eta,\varphi}(x; \alpha, m, a, b) \\ & = \frac{\eta(\varphi(x), \varphi(a), m)^{\alpha+1}}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 g^\alpha(t) f'\left(m\varphi(a) + g(t)\eta(\varphi(x), \varphi(a), m)\right) d[g(t)] \\ & \quad - \frac{\eta(\varphi(x), \varphi(b), m)^{\alpha+1}}{\eta(\varphi(b), \varphi(a), m)} \int_0^1 g^\alpha(t) f'\left(m\varphi(b) + g(t)\eta(\varphi(x), \varphi(b), m)\right) d[g(t)]. \end{aligned}$$

By using Lemma 3.1, one can extend to the following results.

THEOREM 3.3. *Let $\varphi : I \rightarrow \mathbb{R}$ be a continuous increasing function and $g : [0, 1] \rightarrow [0, 1]$ is a differentiable function. Suppose $A \subseteq \mathbb{R}$ be an open m -invex subset with respect to $\eta : A \times A \times (0, 1] \rightarrow \mathbb{R}$ for some fixed $s, m \in (0, 1]$ and let $\varphi(a), \varphi(b) \in A$, $a < b$ with $m\varphi(a) < m\varphi(a) + \eta(\varphi(b), \varphi(a), m)$. Assume that $f : A \rightarrow \mathbb{R}$ is a differentiable function on A° . If $|f'|^q$ is a generalized (g, s, m, φ) -preinvex function on $[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$, $q > 1$, $p^{-1} + q^{-1} = 1$, then for $\alpha > 0$, we have*

$$\begin{aligned}
 & |S_{f,g,\eta,\varphi}(x; \alpha, m, a, b)| \tag{3.2} \\
 & \leq \frac{1}{(s+1)^{1/q}} \left(\frac{g^{p\alpha+1}(1) - g^{p\alpha+1}(0)}{p\alpha+1} \right)^{\frac{1}{p}} \frac{1}{|\eta(\varphi(b), \varphi(a), m)|} \\
 & \times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{\alpha+1} \left[m((1-g(0))^{s+1} - (1-g(1))^{s+1}) |f'(\varphi(a))|^q \right. \right. \\
 & \quad \left. \left. + (g^{s+1}(1) - g^{s+1}(0)) |f'(\varphi(x))|^q \right]^{\frac{1}{q}} \right. \\
 & \quad \left. + |\eta(\varphi(x), \varphi(b), m)|^{\alpha+1} \left[m((1-g(0))^{s+1} - (1-g(1))^{s+1}) |f'(\varphi(b))|^q \right. \right. \\
 & \quad \left. \left. + (g^{s+1}(1) - g^{s+1}(0)) |f'(\varphi(x))|^q \right]^{\frac{1}{q}} \right\}.
 \end{aligned}$$

Proof. Suppose that $q > 1$. Using Lemma 3.1, generalized (g, s, m, φ) -preinvexity of $|f'|^q$, Hölder inequality and taking the modulus, we have

$$\begin{aligned}
 & |S_{f,g,\eta,\varphi}(x; \alpha, m, a, b)| \\
 & \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \int_0^1 g^\alpha(t) \left| f' \left(m\varphi(a) + g(t)\eta(\varphi(x), \varphi(a), m) \right) \right| d[g(t)] \\
 & \quad + \frac{|\eta(\varphi(x), \varphi(b), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \int_0^1 g^\alpha(t) \left| f' \left(m\varphi(b) + g(t)\eta(\varphi(x), \varphi(b), m) \right) \right| d[g(t)] \\
 & \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \left(\int_0^1 g^{p\alpha}(t) d[g(t)] \right)^{\frac{1}{p}} \\
 & \quad \times \left(\int_0^1 \left| f' \left(m\varphi(a) + g(t)\eta(\varphi(x), \varphi(a), m) \right) \right|^q d[g(t)] \right)^{\frac{1}{q}}
 \end{aligned}$$

$$\begin{aligned}
 & + \frac{|\eta(\varphi(x), \varphi(b), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \left(\int_0^1 g^{p\alpha}(t) d[g(t)] \right)^{\frac{1}{p}} \\
 & \quad \times \left(\int_0^1 \left| f'(m\varphi(b) + g(t)\eta(\varphi(x), \varphi(b), m)) \right|^q d[g(t)] \right)^{\frac{1}{q}} \\
 \leq & \frac{|\eta(\varphi(x), \varphi(a), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \left(\int_0^1 g^{p\alpha}(t) d[g(t)] \right)^{\frac{1}{p}} \\
 & \quad \times \left[\int_0^1 \left(m(1-g(t))^s |f'(\varphi(a))|^q + g^s(t) |f'(\varphi(x))|^q \right) d[g(t)] \right]^{\frac{1}{q}} \\
 & + \frac{|\eta(\varphi(x), \varphi(b), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \left(\int_0^1 g^{p\alpha}(t) d[g(t)] \right)^{\frac{1}{p}} \\
 & \quad \times \left[\int_0^1 \left(m(1-g(t))^s |f'(\varphi(b))|^q + g^s(t) |f'(\varphi(x))|^q \right) d[g(t)] \right]^{\frac{1}{q}} \\
 = & \frac{1}{(s+1)^{1/q}} \left(\frac{g^{p\alpha+1}(1) - g^{p\alpha+1}(0)}{p\alpha+1} \right)^{\frac{1}{p}} \frac{1}{|\eta(\varphi(b), \varphi(a), m)|} \\
 & \quad \times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{\alpha+1} \left[m \left((1-g(0))^{s+1} - (1-g(1))^{s+1} \right) |f'(\varphi(a))|^q \right. \right. \\
 & \quad \quad \left. \left. + (g^{s+1}(1) - g^{s+1}(0)) |f'(\varphi(x))|^q \right]^{\frac{1}{q}} \right. \\
 & \quad \left. + |\eta(\varphi(x), \varphi(b), m)|^{\alpha+1} \left[m \left((1-g(0))^{s+1} - (1-g(1))^{s+1} \right) |f'(\varphi(b))|^q \right. \right. \\
 & \quad \quad \left. \left. + (g^{s+1}(1) - g^{s+1}(0)) |f'(\varphi(x))|^q \right]^{\frac{1}{q}} \right\}.
 \end{aligned}$$

■

COROLLARY 3.4. Under the conditions of Theorem 3.3 for $g(t) = t$ and $|f'| \leq K$, we get

$$\begin{aligned} & \frac{1}{|\eta(\varphi(b), \varphi(a), m)|} \left| \eta(\varphi(x), \varphi(a), m)^\alpha f(m\varphi(a) + \eta(\varphi(x), \varphi(a), m)) \right. \\ & \quad - \eta(\varphi(x), \varphi(b), m)^\alpha f(m\varphi(b) + \eta(\varphi(x), \varphi(b), m)) \\ & \quad - \Gamma(\alpha + 1) \left[J_{(m\varphi(a) + \eta(\varphi(x), \varphi(a), m))^-}^\alpha f(m\varphi(a)) \right. \\ & \quad \quad \left. - J_{(m\varphi(b) + \eta(\varphi(x), \varphi(b), m))^-}^\alpha f(m\varphi(b)) \right] \left| \right. \\ & \leq \frac{K}{(p\alpha + 1)^{1/p}} \left(\frac{m + 1}{s + 1} \right)^{\frac{1}{q}} \left[\frac{|\eta(\varphi(x), \varphi(a), m)|^{\alpha+1} + |\eta(\varphi(x), \varphi(b), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \right]. \end{aligned}$$

THEOREM 3.5. *Let $\varphi : I \rightarrow \mathbb{R}$ be a continuous increasing function and $g : [0, 1] \rightarrow [0, 1]$ is a differentiable function. Suppose $A \subseteq \mathbb{R}$ be an open m -invex subset with respect to $\eta : A \times A \times (0, 1] \rightarrow \mathbb{R}$ for some fixed $s, m \in (0, 1]$ and let $\varphi(a), \varphi(b) \in A$, $a < b$ with $m\varphi(a) < m\varphi(a) + \eta(\varphi(b), \varphi(a), m)$. Assume that $f : A \rightarrow \mathbb{R}$ is a differentiable function on A° . If $|f'|^q$ is a generalized (g, s, m, φ) -preinvex function on $[m\varphi(a), m\varphi(a) + \eta(\varphi(b), \varphi(a), m)]$, $q \geq 1$, then for $\alpha > 0$, we have*

$$\begin{aligned} |S_{f,g,\eta,\varphi}(x; \alpha, m, a, b)| & \leq \left(\frac{g^{\alpha+1}(1) - g^{\alpha+1}(0)}{\alpha + 1} \right)^{1 - \frac{1}{q}} \frac{1}{|\eta(\varphi(b), \varphi(a), m)|} \quad (3.3) \\ & \quad \times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{\alpha+1} \left[m |f'(\varphi(a))|^q B(g(t); \alpha, s) \right. \right. \\ & \quad \quad \left. \left. + \left(\frac{g^{\alpha+s+1}(1) - g^{\alpha+s+1}(0)}{\alpha + s + 1} \right) |f'(\varphi(x))|^q \right]^{\frac{1}{q}} \right. \\ & \quad \quad \left. + |\eta(\varphi(x), \varphi(b), m)|^{\alpha+1} \left[m |f'(\varphi(b))|^q B(g(t); \alpha, s) \right. \right. \\ & \quad \quad \left. \left. + \left(\frac{g^{\alpha+s+1}(1) - g^{\alpha+s+1}(0)}{\alpha + s + 1} \right) |f'(\varphi(x))|^q \right]^{\frac{1}{q}} \right\}, \end{aligned}$$

where $B(g(t); \alpha, s) = \int_0^1 g^\alpha(t) (1 - g(t))^s d[g(t)]$.

Proof. Suppose that $q \geq 1$. Using Lemma 3.1, generalized (g, s, m, φ) -preinvexity of $|f'|^q$, the well-known power mean inequality and taking the modulus, we have

$$\begin{aligned}
& |S_{f,g,\eta,\varphi}(x; \alpha, m, a, b)| \\
& \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \int_0^1 g^\alpha(t) \left| f' \left(m\varphi(a) + g(t)\eta(\varphi(x), \varphi(a), m) \right) \right| d[g(t)] \\
& \quad + \frac{|\eta(\varphi(x), \varphi(b), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \int_0^1 g^\alpha(t) \left| f' \left(m\varphi(b) + g(t)\eta(\varphi(x), \varphi(b), m) \right) \right| d[g(t)] \\
& \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \left(\int_0^1 g^\alpha(t) d[g(t)] \right)^{1-\frac{1}{q}} \\
& \quad \times \left(\int_0^1 g^\alpha(t) \left| f' \left(m\varphi(a) + g(t)\eta(\varphi(x), \varphi(a), m) \right) \right|^q d[g(t)] \right)^{\frac{1}{q}} \\
& \quad + \frac{|\eta(\varphi(x), \varphi(b), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \left(\int_0^1 g^\alpha(t) d[g(t)] \right)^{1-\frac{1}{q}} \\
& \quad \times \left(\int_0^1 g^\alpha(t) \left| f' \left(m\varphi(b) + g(t)\eta(\varphi(x), \varphi(b), m) \right) \right|^q d[g(t)] \right)^{\frac{1}{q}} \\
& \leq \frac{|\eta(\varphi(x), \varphi(a), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \left(\int_0^1 g^\alpha(t) d[g(t)] \right)^{1-\frac{1}{q}} \\
& \quad \times \left[\int_0^1 g^\alpha(t) \left(m(1-g(t))^s |f'(\varphi(a))|^q + g^s(t) |f'(\varphi(x))|^q \right) d[g(t)] \right]^{\frac{1}{q}} \\
& \quad + \frac{|\eta(\varphi(x), \varphi(b), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \left(\int_0^1 g^\alpha(t) d[g(t)] \right)^{1-\frac{1}{q}} \\
& \quad \times \left[\int_0^1 g^\alpha(t) \left(m(1-g(t))^s |f'(\varphi(b))|^q + g^s(t) |f'(\varphi(x))|^q \right) d[g(t)] \right]^{\frac{1}{q}}
\end{aligned}$$

$$\begin{aligned}
&= \left(\frac{g^{\alpha+1}(1) - g^{\alpha+1}(0)}{\alpha + 1} \right)^{1-\frac{1}{q}} \frac{1}{|\eta(\varphi(b), \varphi(a), m)|} \\
&\times \left\{ |\eta(\varphi(x), \varphi(a), m)|^{\alpha+1} \left[m |f'(\varphi(a))|^q B(g(t); \alpha, s) \right. \right. \\
&\quad \left. \left. + \left(\frac{g^{\alpha+s+1}(1) - g^{\alpha+s+1}(0)}{\alpha + s + 1} \right) |f'(\varphi(x))|^q \right]^{\frac{1}{q}} \right. \\
&+ |\eta(\varphi(x), \varphi(b), m)|^{\alpha+1} \left[m |f'(\varphi(b))|^q B(g(t); \alpha, s) \right. \\
&\quad \left. \left. + \left(\frac{g^{\alpha+s+1}(1) - g^{\alpha+s+1}(0)}{\alpha + s + 1} \right) |f'(\varphi(x))|^q \right]^{\frac{1}{q}} \right\}. \blacksquare
\end{aligned}$$

COROLLARY 3.6. Under the conditions of Theorem 3.5 for $g(t) = t$ and $|f'| \leq K$, we get

$$\begin{aligned}
&\frac{1}{|\eta(\varphi(b), \varphi(a), m)|} \left| \eta(\varphi(x), \varphi(a), m)^\alpha f(m\varphi(a) + \eta(\varphi(x), \varphi(a), m)) \right. \\
&\quad \left. - \eta(\varphi(x), \varphi(b), m)^\alpha f(m\varphi(b) + \eta(\varphi(x), \varphi(b), m)) \right. \\
&\quad \left. - \Gamma(\alpha + 1) \left[J_{(m\varphi(a) + \eta(\varphi(x), \varphi(a), m))}^\alpha - f(m\varphi(a)) \right. \right. \\
&\quad \quad \left. \left. - J_{(m\varphi(b) + \eta(\varphi(x), \varphi(b), m))}^\alpha - f(m\varphi(b)) \right] \right| \\
&\leq \frac{K}{(1 + \alpha)^{1-\frac{1}{q}}} \left(m\beta(\alpha + 1, s + 1) + \frac{1}{\alpha + s + 1} \right)^{\frac{1}{q}} \\
&\quad \times \left[\frac{|\eta(\varphi(x), \varphi(a), m)|^{\alpha+1} + |\eta(\varphi(x), \varphi(b), m)|^{\alpha+1}}{|\eta(\varphi(b), \varphi(a), m)|} \right].
\end{aligned}$$

Remark 3.7. For a particular choices of a differentiable function $g : [0, 1] \rightarrow [0, 1]$, for example: e^{-t} , $\ln(t + 1)$, $\sin\left(\frac{\pi t}{2}\right)$, $\cos\left(\frac{\pi t}{2}\right)$, etc., by our theorems mentioned in this paper we can get some special kinds of Ostrowski type fractional inequalities.

4. APPLICATIONS TO SPECIAL MEANS

In the following we give certain generalizations of some notions for a positive valued function of a positive variable.

DEFINITION 4.1. (see [13]) A function $M : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$, is called a Mean function if it has the following properties:

1. Homogeneity: $M(ax, ay) = aM(x, y)$, for all $a > 0$,
2. Symmetry: $M(x, y) = M(y, x)$,
3. Reflexivity: $M(x, x) = x$,
4. Monotonicity: If $x \leq x'$ and $y \leq y'$, then $M(x, y) \leq M(x', y')$,
5. Internality: $\min\{x, y\} \leq M(x, y) \leq \max\{x, y\}$.

We consider some means for arbitrary positive real numbers α, β ($\alpha \neq \beta$).

1. The arithmetic mean:

$$A := A(\alpha, \beta) = \frac{\alpha + \beta}{2}$$

2. The geometric mean:

$$G := G(\alpha, \beta) = \sqrt{\alpha\beta}$$

3. The harmonic mean:

$$H := H(\alpha, \beta) = \frac{2}{\frac{1}{\alpha} + \frac{1}{\beta}}$$

4. The power mean:

$$P_r := P_r(\alpha, \beta) = \left(\frac{\alpha^r + \beta^r}{2} \right)^{\frac{1}{r}}, \quad r \geq 1.$$

5. The identric mean:

$$I := I(\alpha, \beta) = \begin{cases} \frac{1}{e} \left(\frac{\beta^\beta}{\alpha^\alpha} \right), & \alpha \neq \beta; \\ \alpha, & \alpha = \beta. \end{cases}$$

6. The logarithmic mean:

$$L := L(\alpha, \beta) = \frac{\beta - \alpha}{\ln |\beta| - \ln |\alpha|}; \quad |\alpha| \neq |\beta|, \quad \alpha\beta \neq 0.$$

7. The generalized log-mean:

$$L_p := L_p(\alpha, \beta) = \left[\frac{\beta^{p+1} - \alpha^{p+1}}{(p+1)(\beta - \alpha)} \right]^{\frac{1}{p}}; \quad p \in \mathbb{R} \setminus \{-1, 0\}, \quad \alpha \neq \beta.$$

8. The weighted p -power mean:

$$M_p \left(\begin{array}{cccc} \alpha_1 & \alpha_2 & \cdots & \alpha_n \\ u_1 & u_2 & \cdots & u_n \end{array} \right) = \left(\sum_{i=1}^n \alpha_i u_i^p \right)^{\frac{1}{p}}$$

where $0 \leq \alpha_i \leq 1$, $u_i > 0$ ($i = 1, 2, \dots, n$) with $\sum_{i=1}^n \alpha_i = 1$.

It is well known that L_p is monotonic nondecreasing over $p \in \mathbb{R}$ with $L_{-1} := L$ and $L_0 := I$. In particular, we have the following inequality $H \leq G \leq L \leq I \leq A$. Now, let a and b be positive real numbers such that $a < b$. Consider the function $M := M(\varphi(a), \varphi(b)) : [\varphi(a), \varphi(a) + \eta(\varphi(b), \varphi(a))] \times [\varphi(a), \varphi(a) + \eta(\varphi(b), \varphi(a))] \rightarrow \mathbb{R}_+$, which is one of the above mentioned means, $\varphi : I \rightarrow \mathbb{R}$ be a continuous increasing function and $g : [0, 1] \rightarrow [0, 1]$ is a differentiable function. Therefore one can obtain various inequalities using the results of Section 3 for these means as follows:

Replace $\eta(\varphi(y), \varphi(x), m)$ with $\eta(\varphi(y), \varphi(x))$ and setting $\eta(\varphi(a), \varphi(b)) = M(\varphi(a), \varphi(b))$ for $m = 1$ in (3.2) and (3.3), one can obtain the following

interesting inequalities involving means:

$$\begin{aligned}
 & \frac{1}{|M(\varphi(a), \varphi(b))|} \left| M(\varphi(a), \varphi(x))^\alpha \left[g^\alpha(1)f(\varphi(a) + g(1)M(\varphi(a), \varphi(x))) \right. \right. \\
 & \qquad \qquad \qquad \left. \left. - g^\alpha(0)f(\varphi(a) + g(0)M(\varphi(a), \varphi(x))) \right] \right. \\
 & \qquad \qquad \qquad \left. - M(\varphi(b), \varphi(x))^\alpha \left[g^\alpha(1)f(\varphi(b) + g(1)M(\varphi(b), \varphi(x))) \right. \right. \\
 & \qquad \qquad \qquad \left. \left. - g^\alpha(0)f(\varphi(b) + g(0)M(\varphi(b), \varphi(x))) \right] \right. \\
 & \qquad \qquad \qquad \left. - \alpha \left[\int_{\varphi(a)+g(0)M(\varphi(a), \varphi(x))}^{\varphi(a)+g(1)M(\varphi(a), \varphi(x))} (t - \varphi(a))^{\alpha-1} f(t) dt \right. \right. \\
 & \qquad \qquad \qquad \left. \left. - \int_{\varphi(b)+g(0)M(\varphi(b), \varphi(x))}^{\varphi(b)+g(1)M(\varphi(b), \varphi(x))} (t - \varphi(b))^{\alpha-1} f(t) dt \right] \right| \\
 & \leq \frac{1}{(s+1)^{1/q}} \left(\frac{g^{p\alpha+1}(1) - g^{p\alpha+1}(0)}{p\alpha + 1} \right)^{\frac{1}{p}} \frac{1}{M(\varphi(a), \varphi(b))} \tag{4.1} \\
 & \times \left\{ M(\varphi(a), \varphi(x))^{\alpha+1} \left[\left((1 - g(0))^{s+1} - (1 - g(1))^{s+1} \right) |f'(\varphi(a))|^q \right. \right. \\
 & \qquad \qquad \qquad \left. \left. + (g^{s+1}(1) - g^{s+1}(0)) |f'(\varphi(x))|^q \right]^{\frac{1}{q}} \right. \\
 & \qquad \qquad \qquad \left. + M(\varphi(b), \varphi(x))^{\alpha+1} \left[\left((1 - g(0))^{s+1} - (1 - g(1))^{s+1} \right) |f'(\varphi(b))|^q \right. \right. \\
 & \qquad \qquad \qquad \left. \left. + (g^{s+1}(1) - g^{s+1}(0)) |f'(\varphi(x))|^q \right]^{\frac{1}{q}} \right\},
 \end{aligned}$$

$$\begin{aligned}
 & \frac{1}{M(\varphi(a), \varphi(b))} \left| M(\varphi(a), \varphi(x))^\alpha \left[g^\alpha(1)f(\varphi(a) + g(1)M(\varphi(a), \varphi(x))) \right. \right. \\
 & \qquad \qquad \qquad \left. \left. - g^\alpha(0)f(\varphi(a) + g(0)M(\varphi(a), \varphi(x))) \right] \right. \\
 & \qquad \qquad \qquad \left. - M(\varphi(b), \varphi(x))^\alpha \left[g^\alpha(1)f(\varphi(b) + g(1)M(\varphi(b), \varphi(x))) \right. \right. \\
 & \qquad \qquad \qquad \left. \left. - g^\alpha(0)f(\varphi(b) + g(0)M(\varphi(b), \varphi(x))) \right] \right. \\
 & \qquad \qquad \qquad \left. - \alpha \left[\int_{\varphi(a)+g(0)M(\varphi(a), \varphi(x))}^{\varphi(a)+g(1)M(\varphi(a), \varphi(x))} (t - \varphi(a))^{\alpha-1} f(t) dt \right. \right.
 \end{aligned}$$

$$\begin{aligned}
& \left| - \int_{\varphi(b)+g(0)M(\varphi(b),\varphi(x))}^{\varphi(b)+g(1)M(\varphi(b),\varphi(x))} (t - \varphi(b))^{\alpha-1} f(t) dt \right| \\
\leq & \left(\frac{g^{\alpha+1}(1) - g^{\alpha+1}(0)}{\alpha + 1} \right)^{1-\frac{1}{q}} \frac{1}{M(\varphi(a), \varphi(b))} \\
& \times \left\{ M(\varphi(a), \varphi(x))^{\alpha+1} \left[|f'(\varphi(a))|^q B(g(t); \alpha, s) \right. \right. \\
& \quad \left. \left. + \left(\frac{g^{\alpha+s+1}(1) - g^{\alpha+s+1}(0)}{\alpha + s + 1} \right) |f'(\varphi(x))|^q \right]^{\frac{1}{q}} \right. \\
& \quad \left. + M(\varphi(b), \varphi(x))^{\alpha+1} \left[|f'(\varphi(b))|^q B(g(t); \alpha, s) \right. \right. \\
& \quad \left. \left. + \left(\frac{g^{\alpha+s+1}(1) - g^{\alpha+s+1}(0)}{\alpha + s + 1} \right) |f'(\varphi(x))|^q \right]^{\frac{1}{q}} \right\}. \tag{4.2}
\end{aligned}$$

Letting $M(\varphi(a), \varphi(b)) = A, G, H, P_r, I, L, L_p, M_p$ in (4.1) and (4.2), we get the inequalities involving means for a particular choice of a differentiable generalized $(g, s, 1, \varphi)$ -preinvex functions f . The details are left to the interested reader.

REFERENCES

- [1] T. S. DU, J. G. LIAO, Y. J. LI, Properties and integral inequalities of Hadamard-Simpson type for the generalized (s, m) -preinvex functions, *J. Nonlinear Sci. Appl.* **9** (5) (2016), 3112–3126.
- [2] S. S. DRAGOMIR, J. PEČARIĆ, L. E. PERSSON, Some inequalities of Hadamard type, *Soochow J. Math.* **21** (3) (1995), 335–341.
- [3] H. HUDZIK, L. MALIGRANDA, Some remarks on s -convex functions, *Aequationes Math.* **48** (1) (1994), 100–111.
- [4] T. ANTCZAK, Mean value in invexity analysis, *Nonlinear Anal.* **60** (8) (2005), 1473–1484.
- [5] X. M. YANG, X. Q. YANG, K. L. TEO, Generalized invexity and generalized invariant monotonicity, *J. Optim. Theory Appl.* **117** (3) (2003), 607–625.
- [6] R. PINI, Invexity and generalized convexity, *Optimization.* **22** (4) (1991), 513–525.

- [7] D. D. STANCU, G. COMAN, P. BLAGA, “Analiză Numerică și Teoria Aproximării”, Vol. II, Presa Universitară Clujeană, Cluj-Napoca, 2002.
- [8] W. LIU, New integral inequalities involving beta function via P -convexity, *Miskolc Math Notes*. **15** (2) (2014), 585–591.
- [9] M. E. ÖZDEMİR, E. SET, M. ALOMARI, Integral inequalities via several kinds of convexity, *Creat. Math. Inform.* **20** (1) (2011), 62–73.
- [10] W. LIU, W. WEN, J. PARK, Hermite-Hadamard type inequalities for MT-convex functions via classical integrals and fractional integrals, *J. Nonlinear Sci. Appl.* **9** (3) (2016), 766–777.
- [11] W. LIU, Ostrowski type fractional integral inequalities for MT-convex functions, *Miskolc Mathematical Notes*. **16** (1) (2015), 249–256.
- [12] M. TUNÇ, Ostrowski type inequalities for functions whose derivatives are MT-convex, *J. Comput. Anal. Appl.* **17** (4) (2014), 691–696.
- [13] P. S. BULLEN, “Handbook of Means and Their Inequalities”, *Kluwer Academic Publishers Group*, Dordrecht, 2003.