

## Addendum to “Interpolation of Banach Spaces by the $\gamma$ -Method”

JESÚS SUÁREZ, LUTZ WEIS

*Escuela Politécnica, Universidad de Extremadura, Avenida de la Universidad s/n,  
10071 Cáceres, Spain, [jesus@unex.es](mailto:jesus@unex.es)*

*Mathematisches Institut I, Universität Karlsruhe,  
76128 Karlsruhe, Germany, [lutz.weis@math.uni-karlsruhe.de](mailto:lutz.weis@math.uni-karlsruhe.de)*

Received September 19, 2009

*Abstract:* In this note we correct and simplify an interpolation theorem for radonifying operators in [7].

*Key words:* Interpolation of spaces of vector valued functions,  $\gamma$ -radonifying operators.

*AMS Subject Class. (2000):* 46B70, 47B07, 46E40.

### 1. INTRODUCTION

In [7] it was claimed that under B-convexity the radonifying operators  $\gamma(H, X)$  form an interpolation scale for the complex method. However the proof given there used the implicit hypothesis of a compact embedding  $H_0 \rightarrow H_1$ . In this note we give a simpler proof which also eliminates the compactness condition and works in some important situation where  $H_0$  is not embedded into  $H_1$ . Among examples are scales of Sobolev spaces and domains of selfadjoints operators. However the case of general interpolation pairs of Hilbert spaces remains open.

### 2. INTERPOLATION OF $\gamma(H, X)$

Given an interpolation couple  $(H_0, H_1)$  of two Hilbert spaces  $H_0, H_1$  continuously embedded in some locally convex space  $H$  one considers the Hilbert space  $H = H_0 \cap H_1$  with the norm  $\|f\|_H^2 = \|f\|_{H_0}^2 + \|f\|_{H_1}^2$ . Since  $(x, y)_{H_j} \leq \|x\|_H \|y\|_H$  for  $j = 0, 1$  the Lax-Milgram theorem gives us positive, selfadjoint operators  $B_j$  on  $H$  of norm 1 so that

$$(x, y)_{H_j} = (x, B_j y)_H.$$

The interpolation couple  $(H_0, H_1)$  of Hilbert spaces is called *commutative* if the two bounded operators  $B_j$  constructed above commute. The point of this definition is the following consequence of the spectral theorem for commuting selfadjoint operators which allows us to replace abstract Hilbert spaces by  $L_2$ -spaces with densities.

LEMMA 2.1. *The interpolation couple  $(H_0, H_1)$  is commutative if and only if there is a  $\sigma$ -finite measure space  $(\Omega, \mu)$ , a unitary map  $U : H_0 \cap H_1 \rightarrow L_2(\Omega, \mu)$  and densities  $0 < g_j \leq 1$  so that  $U$  extends to isometries  $U_j : H_j \rightarrow L_2(\Omega, g_j d\mu)$ .*

*Proof.* The spectral theorem (see e.g. [6, p. 246, Problem 4]) gives  $(\Omega, \mu)$ ,  $U$  and  $g_j$  so that for  $x, y \in H_0 \cap H_1$

$$\begin{aligned} (x, y)_{H_j} &= (x, B_j y)_H = (Ux, (UB_j U^{-1})Uy) \\ &= \int_{\Omega} \overline{f(\omega)} h(\omega) g_j(\omega) d\mu(\omega) = (f, h)_{L_2(g_j d\mu)} \end{aligned}$$

with  $f = U(x)$ ,  $h = U(y)$  from a dense subset of  $L_2(\Omega, g_j d\mu)$ . Since the operators  $B_j$  have norm one and are positive and injective we obtain  $0 < g_j \leq 1$ . ■

There are some important interpolation couples covered by this situation.

EXAMPLES. (1) If  $H_0, H_1$  with a continuous dense embedding  $H_0 \cap H_1 = H_0$  and  $B_0 = Id$ , so that the pair  $(H_0, H_1)$  is commutative.

(2) Let  $A$  be a positive injective and selfadjoint operator on a Hilbert space  $G$  and  $H_j = D(A^{\alpha_j})$  with  $\|h\|_j = \|A^{\alpha_j} x\|$  for  $\alpha_j \in \mathbb{R}$ ,  $j = 0, 1$ . Then the pair  $(H_0, H_1)$  is commutative. Indeed it is easy to check that  $B_j = A^{2\alpha_j} (A^{2\alpha_0} + A^{2\alpha_1})^{-1}$ .

(3) As a particular case of (2) we obtain for  $G = L_2(\mathbb{R}^n)$  and  $A = \Delta$  the pairs of Sobolev spaces  $H_2^{\alpha_0}(\mathbb{R}^n)$ ,  $H_2^{\alpha_1}(\mathbb{R}^n)$  for  $\alpha_j \in \mathbb{R}$ .

(4) An interpolation pair which is not commutative can be obtained as follows:

Let  $H_0 = L_2(\mathbb{R}, \langle t \rangle^{-2} dt)$  with  $\langle t \rangle = (1 + t^2)^{1/2}$  and  $H_1 = H^2(\mathbb{R})$ . Then  $L_2(\mathbb{R}) \subset H_0 \cap H_1$ . However the multiplication operator  $B_0 f(t) = \langle t \rangle^{-2} f(t)$  and  $B_2 = (I - A)^{-1}$  do not commute.

THEOREM 2.1. *Let  $H_0, H_1$  be separable Hilbert spaces forming a commutative interpolation couple. If  $X_0, X_1$  are  $B$ -convex, then*

$$(\gamma(H_0, X_0), \gamma(H_1, X_1))_{[\theta]} = \gamma((H_0, H_1)_{[\theta]}, (X_0, X_1)_{[\theta]})$$

with equivalent norms.

*Proof.* Since it is well known already that for  $H = (H_0, H_1)_{[\theta]}$  we have, see [4],

$$(\gamma(H_0, X_0), \gamma(H_1, X_1))_{[\theta]} = \gamma(H, (X_0, X_1)_{[\theta]})$$

it only remains to show that

$$(\gamma(H_0, X_0), \gamma(H_1, X_1))_{[\theta]} = (\gamma(H, X_0), \gamma(H, X_1))_{[\theta]}.$$

By assumption and the lemma we may assume that  $H_0 \cap H_1 = L_2(\Omega, \mu)$  and  $H_j = L_2(\Omega, g_j d\mu)$ ,  $j = 0, 1$ , for a  $\sigma$ -finite measure space  $(\Omega, \mu)$  and densities  $0 < g_j \leq 1$  on  $\Omega$ . For convenience let us use the abbreviation  $\gamma_j(h)$  for  $L_2(\Omega, g_j d\mu)$ . Note the following continuous inclusion

$$\gamma(L_2(\Omega, d\mu), X_0 \cap X_1) \subset \gamma_j(g_\theta) \subset \gamma(L_2(\Omega, g_0 \wedge g_1 d\mu), X_0 + X_1)$$

where  $g_\theta = g_0^{1-\theta} g_1^\theta$  for  $\theta \in [0, 1]$ . Hence  $(\gamma_0(g_0), \gamma_1(g_1))$  and  $(\gamma_0(g_\theta), \gamma_1(g_\theta))$  are interpolation couples and, going back to the definition of the complex interpolation method, we can consider the following [1] Banach space  $\mathcal{F}$  of all continuous

$$f : S = \{z : 0 \leq \operatorname{Re} z \leq 1\} \longrightarrow \gamma_0(g_0) + \gamma_1(g_1)$$

which are all continuous on  $S$  and analytic in its interior such that  $f(j + it) \in \gamma_j(g_j)$ ,  $j = 0, 1$ , with  $\|f(j + it)\| \rightarrow 0$  for  $|t| \rightarrow \infty$  and

$$\|f\|_{\mathcal{F}} = \sup \left\{ \|f(j + it)\|_{\gamma_j(g_j)} : t \in \mathbb{R} \right\}.$$

Then  $(\gamma_0(g_0), \gamma_1(g_1))_{[\theta]} = \{f(\theta) : f \in \mathcal{F}\}$  endowed with the quotient norm

$$\|T\| = \inf \{ \|f\| : f(\theta) = T \}.$$

In a similar way one defines  $\mathcal{F}_\theta$  for the pair  $(\gamma_0(g_\theta), \gamma_1(g_\theta))$ . For the moment we assume in addition that

$$\delta < g_j \quad \text{on } \Omega \quad \text{for } j = 0, 1 \quad \text{and some fixed } \delta > 0. \quad (*)$$

To prove the inclusion

$$(\gamma_0(g_0), \gamma_1(g_1))_{[\theta]} \subset (\gamma_0(g_\theta), \gamma_1(g_\theta))_{[\theta]},$$

we fix  $\theta \in [0, 1]$  and choose  $f \in \mathcal{F}$  with  $\|f\|_{\mathcal{F}} \leq 2\|f(\theta)\|_{(\gamma_0(g_0), \gamma_1(g_1))_{[\theta]}}$ . Next we define a family of operators

$$T(z) : L_2(\Omega, g_\theta d\mu) \longrightarrow L_2(\Omega, (g_0 \wedge g_1) d\mu)$$

with  $z \in S$  as the multiplication operators

$$T(z)f := \left( \frac{g_\theta}{g_0^{1-z} g_1^z} \right)^{1/2} f.$$

By our assumption we have bounded and invertible operators  $T(z)$  for all  $z$  in the interior of  $S$ . However for  $z = j + it$  these operators extend to isometries by [3]

$$T(j + it) : \gamma_j(g_\theta) \longrightarrow \gamma_j(g_j)$$

and  $T(\theta) = Id$ . It follows that  $f \circ T$  belongs to  $\mathcal{F}_\theta$  and  $\|f \circ T\| \leq \|f\|$ . Therefore

$$\|f \circ T(\theta)\| \leq \|f \circ T\| \leq 2\|f(\theta)\|$$

and the inclusion  $\subset$  is continuous. To show the reverse we argue similarly now replacing  $T(z)$  by  $S(z) = T(z)^{-1}$ .

It remains to remove the assumption (\*). Note that the assumption  $\delta < g_j$  was only used to bound the operators  $T(z)$  and  $T(z)^{-1}$  for  $z$  in the interior of  $S$ . For  $z \in \partial S$  and  $z = \theta$  the  $T(z)$  were isometries independent of the condition  $\delta < g_j$ , which therefore does not enter the calculation of the norms  $\|f\|$  or  $\|f \circ T\|$ . Now, for any  $\delta > 0$ , if we consider functions  $f$  which are zero on  $\Omega_\delta = \{g_0 \leq \delta\} \cup \{g_1 \leq \delta\}$  the above norm estimate apply with constants independent of  $\delta$ . However such functions  $f$  are dense in  $\mathcal{F}$  or  $\mathcal{F}_\theta$ . ■

#### REFERENCES

- [1] J. BERGH, J. LÖFSTRÖM, "Interpolation spaces. An introduction", Grundlehren der Mathematischen Wissenschaften, No. 223, Springer-Verlag, Berlin-New York, 1976.
- [2] N.J. KALTON, L. WEIS, The  $H^\infty$ -calculus and sums of closed operators, *Math. Ann.* **321** (2) (2001), 319–345.
- [3] N.J. KALTON, L. WEIS, Euclidean structures and their application to spectral theory, in preparation.

- [4] N. KALTON, P. KUNSTMANN, L. WEIS, Perturbation and interpolation theorems for the  $H^\infty$ -calculus with applications to differential operators, *Math. Ann.* **336** (4) (2006), 747–801.
- [5] P. KUNSTMANN, L. WEIS, Maximal  $L_p$ -regularity for parabolic equations, Fourier multiplier theorems and  $H^\infty$ -functional calculus, in “Functional Analytic Methods for Evolution Equations”, Lecture Notes in Math., No. 1855, Springer, Berlin, 2004, 65–311.
- [6] M. REED, B. SIMON, “Methods of Modern Mathematical Physics. I. Functional Analysis”, Academic Press, New York-London, 1972.
- [7] J. SUÁREZ, L. WEIS, Interpolation of Banach spaces by the  $\gamma$ -method, in “Methods in Banach Space Theory”, London Math. Soc. Lecture Note Ser., No. 337, Cambridge Univ. Press, Cambridge, 2006, 293–306.
- [8] H. TRIEBEL, “Interpolation Theory, Function Spaces, Differential Operators”, North-Holland Mathematical Library, No. 18, North-Holland Publishing Co., Amsterdam-New York, 1978.