

On Three-Dimensional Trans-Sasakian Manifolds

U. C. DE, AVIJIT SARKAR

*Department of Mathematics, University of Kalyani, Kalyani 741235,
West Bengal, India, e-mail: uc_de@yahoo.com*

*Department of Mathematics, University of Burdwan, Burdwan 713104,
West Bengal, India, e-mail: avjaj@yahoo.co.in*

Presented by Oscar García Prada

Received February 23, 2008

Abstract: The object of the present paper is to study 3-dimensional trans-Sasakian manifolds which are locally ϕ -symmetric and have η -parallel Ricci tensor. Also 3-dimensional trans-Sasakian manifolds of constant curvature have been considered. An example of a three-dimensional locally ϕ -symmetric trans-Sasakian manifold is given.

Key words: trans-Sasakian manifold, scalar curvature, locally ϕ -symmetric, η -parallel Ricci tensor, constant curvature.

AMS *Subject Class.* (2000): 53C25.

1. INTRODUCTION

Trans-Sasakian manifolds arose in a natural way from the classification of almost contact metric structures by D. Chinea and C. Gonzales [3], and they appear as a natural generalization of both Sasakian and Kenmotsu manifolds. Again in the Gray-Hervella classification of almost Hermite manifolds [7], there appears a class W_4 of Hermitian manifolds which are closely related to locally conformally Kähler manifolds. An almost contact metric structure on a manifold M is called a trans-Sasakian structure [13] if the product manifold $M \times R$ belongs to the class W_4 . The class $C_6 \oplus C_5$ ([10], [11]) coincides with the class of trans-Sasakian structures of type (α, β) . In [11], the local nature of the two subclasses C_5 and C_6 of trans-Sasakian structures is characterized completely. In [4], some curvature identities and sectional curvatures for C_5 , C_6 and trans-Sasakian manifolds are obtained. It is known that ([8]) trans-Sasakian structures of type $(0,0)$, $(0,\beta)$ and $(\alpha,0)$ are cosymplectic, β -Kenmotsu and α -Sasakian respectively. In [15], it is proved that trans-Sasakian structures are generalized quasi-Sasakian structures [12]. Thus, trans-Sasakian structures also provide a large class of generalized quasi-Sasakian structures.

The local structure of trans-Sasakian manifolds of dimension $n \geq 5$ has been completely characterized by J. C. Marrero [10]. He proved that a trans-

Sasakian manifold of dimension $n \geq 5$ is either cosymplectic or α -Sasakian or β -Kenmotsu manifold. But so far, it is not too much known about the 3-dimensional case.

This paper deals just on 3-dimensional connected trans-Sasakian manifolds. In Section 2 some preliminary results are recalled and explicit formulae for Ricci tensor and curvature tensor [6] of 3-dimensional trans Sasakian manifolds are given. In Section 3 we characterize 3-dimensional locally ϕ -symmetric trans-Sasakian manifolds and prove that a 3-dimensional connected trans-Sasakian manifold of type (α, β) is locally ϕ -symmetric if and only if the scalar curvature of the manifold is constant where α and β are constants. This result is an extension of an analogous result concerning Kenmotsu manifolds obtained by the first author [5]. Section 4 of our paper deals with a 3-dimensional trans-Sasakian manifold with η -parallel Ricci tensor. In this section we also show that a 3-dimensional connected trans-Sasakian manifold of type (α, β) has η -parallel Ricci tensor if and only if the scalar curvature of the manifold is constant where α and β are constants. In Section 5, we show that a 3-dimensional compact connected trans-Sasakian manifold of constant curvature is either α -Sasakian or β -Kenmotsu. This is the most important result obtained in this paper. Finally in the last section we construct an example of a three-dimensional locally ϕ -symmetric trans-Sasakian manifold.

2. PRELIMINARIES

Let M be a connected almost contact metric manifold with an almost contact metric structure (ϕ, ξ, η, g) , that is, ϕ is an $(1, 1)$ tensor field, ξ is a vector field, η is an 1-form and g is compatible Riemannian metric such that

$$\phi^2(X) = -X + \eta(X)\xi, \quad \eta(\xi) = 1, \quad \phi\xi = 0, \quad \eta\phi = 0, \quad (2.1)$$

$$g(\phi X, \phi Y) = g(X, Y) - \eta(X)\eta(Y), \quad (2.2)$$

$$g(X, \phi Y) = -g(\phi X, Y), \quad g(X, \xi) = \eta(X), \quad (2.3)$$

for all $X, Y \in T(M)$ [1]. The fundamental 2-form Φ of the manifold is defined by

$$\Phi(X, Y) = g(X, \phi Y), \quad (2.4)$$

for $X, Y \in T(M)$.

An almost contact metric structure (ϕ, ξ, η, g) on a connected manifold M is called trans-Sasakian structure [13] if $(M \times R, J, G)$ belongs to the class

W_4 [7], where J is the almost complex structure on $M \times R$ defined by

$$J(X, fd/df) = (\phi X - f\xi, \eta(X)d/dt),$$

for all vector fields X on M , a smooth function f on $M \times R$ and the product metric G on $M \times R$. This may be expressed by the condition [2]

$$(\nabla_X \phi)Y = \alpha(g(X, Y)\xi - \eta(Y)X) + \beta(g(\phi X, Y)\xi - \eta(Y)\phi X), \tag{2.5}$$

for smooth functions α and β on M . Here we say that the trans-Sasakian structure is of type (α, β) . From (2.5) it follows that

$$\nabla_X \xi = -\alpha\phi X + \beta(X - \eta(X)\xi), \tag{2.6}$$

$$(\nabla_X \eta)Y = -\alpha g(\phi X, Y) + \beta g(\phi X, \phi Y). \tag{2.7}$$

An explicit example of 3-dimensional proper trans-Sasakian manifold is constructed in [10]. In [6], the Ricci operator, Ricci tensor and curvature tensor for 3-dimensional trans-Sasakian manifolds are studied and their explicit formulae are given.

From [6] we know that for a 3-dimensional trans-Sasakian manifold

$$2\alpha\beta + \xi\alpha = 0, \tag{2.8}$$

$$S(X, \xi) = (2(\alpha^2 - \beta^2) - \xi\beta)\eta(X) - X\beta - (\phi X)\alpha, \tag{2.9}$$

$$\begin{aligned} S(X, Y) = & \left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2)\right)g(X, Y) \\ & - \left(\frac{r}{2} + \xi\beta - 3(\alpha^2 - \beta^2)\right)\eta(X)\eta(Y) \\ & - (Y\beta + (\phi Y)\alpha)\eta(X) - (X\beta + (\phi X)\alpha)\eta(Y), \end{aligned} \tag{2.10}$$

and

$$\begin{aligned} R(X, Y)Z = & \left(\frac{r}{2} + 2\xi\beta - 2(\alpha^2 - \beta^2)\right)(g(Y, Z)X - g(X, Z)Y) \\ & - g(Y, Z) \left[\left(\frac{r}{2} + \xi\beta - 3(\alpha^2 - \beta^2)\right)\eta(X)\xi \right. \\ & \quad \left. - \eta(X)(\phi \text{grad}\alpha - \text{grad}\beta) + (X\beta + (\phi X)\alpha)\xi \right] \\ & + g(X, Z) \left[\left(\frac{r}{2} + \xi\beta - 3(\alpha^2 - \beta^2)\right)\eta(Y)\xi \right. \\ & \quad \left. - \eta(Y)(\phi \text{grad}\alpha - \text{grad}\beta) + (Y\beta + (\phi Y)\alpha)\xi \right] \end{aligned} \tag{2.11}$$

$$\begin{aligned}
& - \left[(Z\beta + (\phi Z)\alpha)\eta(Y) + (Y\beta + (\phi Y)\alpha)\eta(Z) \right. \\
& \quad \left. + \left(\frac{r}{2} + \xi\beta - 3(\alpha^2 - \beta^2) \right) \eta(Y)\eta(Z) \right] X \\
& + \left[(Z\beta + (\phi Z)\alpha)\eta(X) + (X\beta + (\phi X)\alpha)\eta(Z) \right. \\
& \quad \left. + \left(\frac{r}{2} + \xi\beta - 3(\alpha^2 - \beta^2) \right) \eta(X)\eta(Z) \right] Y,
\end{aligned}$$

where S is the Ricci tensor of type $(0, 2)$, R is the curvature tensor of type $(1, 3)$ and r is the scalar curvature of the manifold M .

3. LOCALLY ϕ -SYMMETRIC THREE-DIMENSIONAL TRANS-SASAKIAN MANIFOLDS

DEFINITION 3.1. A trans-Sasakian manifold is said to be locally ϕ -symmetric if

$$\phi^2(\nabla_W R)(X, Y)Z = 0,$$

for all vector fields W, X, Y, Z orthogonal to ξ .

This notion was introduced for Sasakian manifolds by Takahashi [14].

Let M be a 3-dimensional connected trans-Sasakian manifold. Then its curvature tensor is given by (2.11). Differentiating (2.11) we get

$$\begin{aligned}
& (\nabla_W R)(X, Y)Z \\
& = \left[\frac{dr(W)}{2} + 2(\nabla_W(\xi\beta)) - 4(d\alpha(W) - d\beta(W)) \right] [g(Y, Z)X - g(X, Z)Y] \\
& \quad - g(Y, Z) \left[\left(\frac{dr(W)}{2} + (\nabla_W(\xi\beta)) - 6(d\alpha(W) - d\beta(W)) \right) \eta(X)\xi \right. \\
& \quad \quad + \left(\frac{r}{2} + \xi\beta - 3(\alpha^2 - \beta^2) \right) ((\nabla_W\eta)(X)\xi + \eta(X)(\nabla_W\xi)) \\
& \quad \quad - (\nabla_W\eta)(X)(\phi(\text{grad}\alpha) - \text{grad}\beta) - \eta(X)(\nabla_W(\phi(\text{grad}\alpha) - \text{grad}\beta)) \\
& \quad \quad \left. + (\nabla_W(X\beta + (\phi X)\alpha))\xi + (X\beta + (\phi X)\alpha)\nabla_W\xi \right] \\
& + g(X, Z) \left[\left(\frac{dr(W)}{2} + (\nabla_W(\xi\beta)) - 6(d\alpha(W) - d\beta(W)) \right) \eta(Y)\xi \right. \\
& \quad + \left(\frac{r}{2} + \xi\beta - 3(\alpha^2 - \beta^2) \right) ((\nabla_W\eta)(Y)\xi + \eta(Y)(\nabla_W\xi)) \\
& \quad - (\nabla_W\eta)(Y)(\phi(\text{grad}\alpha) - \text{grad}\beta) - \eta(Y)(\nabla_W(\phi(\text{grad}\alpha) - \text{grad}\beta)) \\
& \quad \left. + (\nabla_W(Y\beta + (\phi Y)\alpha))\xi + (Y\beta + (\phi Y)\alpha)\nabla_W\xi \right] \tag{3.1}
\end{aligned}$$

$$\begin{aligned}
 & - \left[(\nabla_W(Z\beta + (\phi Z)\alpha))\eta(Y) + (Z\beta + (\phi Z)\alpha)(\nabla_W\eta)Y \right. \\
 & + (\nabla_W(Y\beta + (\phi Y)\alpha))\eta(Z) + (Y\beta + (\phi Y)\alpha)(\nabla_W\eta)Z \\
 & + \left(\frac{dr(W)}{2} + (\nabla_W(\xi\beta)) - 6(d\alpha(W) - d\beta(W)) \right) \eta(Y)\eta(Z) \\
 & + \left(\frac{r}{2} + \xi\beta - 3(\alpha^2 - \beta^2) \right) ((\nabla_W\eta)Y\eta(Z) + \eta(Y)(\nabla_W\eta)Z) \Big] X \\
 & + \left[(\nabla_W(Z\beta + (\phi Z)\alpha))\eta(X) + (Z\beta + (\phi Z)\alpha)(\nabla_W\eta)X \right. \\
 & + (\nabla_W(X\beta + (\phi X)\alpha))\eta(Z) + (X\beta + (\phi X)\alpha)(\nabla_W\eta)Z \\
 & + \left(\frac{dr(W)}{2} + (\nabla_W(\xi\beta)) - 6(d\alpha(W) - d\beta(W)) \right) \eta(X)\eta(Z) \\
 & + \left(\frac{r}{2} + \xi\beta - 3(\alpha^2 - \beta^2) \right) ((\nabla_W\eta)X\eta(Z) + \eta(X)(\nabla_W\eta)Z) \Big] Y.
 \end{aligned}$$

Suppose that α and β are constants and X, Y, Z, W are orthogonal to ξ . Then using $\phi\xi = 0$ and (3.1), we get

$$\phi^2(\nabla_W R)(X, Y)Z = \left(\frac{dr(W)}{2} \right) (g(Y, Z)X - g(X, Z)Y). \tag{3.2}$$

Hence from (3.2) we get $\phi^2(\nabla_W R)(X, Y)Z = 0$ if and only if the scalar curvature r is constant. Thus we can state the following:

THEOREM 3.1. *A 3-dimensional connected trans-Sasakian manifold of type (α, β) is locally ϕ -symmetric if and only if the scalar curvature is constant provided α and β are constants.*

The above theorem is just an extension of an analogous result concerning Kenmotsu manifolds obtained by the first author in the paper [5].

4. η -PARALLEL RICCI TENSOR

DEFINITION 4.1. The Ricci tensor S of a trans-Sasakian manifold is said to be η -parallel if it satisfies

$$(\nabla_X S)(\phi Y, \phi Z) = 0, \tag{4.1}$$

for all vector fields X, Y and Z .

This notion was introduced in the context of Sasakian manifolds by Kon [9].

Let M be a 3-dimensional connected trans-Sasakian manifold. Then its Ricci tensor is given by (2.10)

In (2.10) replacing X by ϕX , Y by ϕY and using (2.1) we get for a trans-Sasakian manifold of dimension three

$$S(\phi X, \phi Y) = \left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2) \right) (g(X, Y) - \eta(X)\eta(Y)). \quad (4.2)$$

Now we see that

$$\begin{aligned} (\nabla_Z S)(\phi X, \phi Y) &= \nabla_Z S(\phi X, \phi Y) - S(\nabla_Z \phi X, \phi Y) - S(\phi X, \nabla_Z \phi Y) \\ &= \nabla_Z S(\phi X, \phi Y) - S((\nabla_Z \phi)X, \phi Y) - S(\phi \nabla_Z X, \phi Y) \\ &\quad - S(\phi X, (\nabla_Z \phi)Y) - S(\phi X, \phi \nabla_Z Y). \end{aligned} \quad (4.3)$$

Using (2.5), (2.10) and (4.2) in (4.3) we have

$$\begin{aligned} &(\nabla_Z S)(\phi X, \phi Y) \\ &= \left(\frac{1}{2} dr(Z) + \nabla_Z(\xi\beta) - 2\alpha d\alpha(Z) + 2\beta d\beta(Z) \right) (g(X, Y) - \eta(X)\eta(Y)) \\ &\quad + \left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2) \right) \left(\nabla_Z g(X, Y) - (\nabla_Z \eta(X))\eta(Y) \right. \\ &\quad \left. - \eta(X)(\nabla_Z \eta(Y)) \right) \\ &\quad - S\left(\alpha(g(Z, X)\xi - \eta(X)Z) + \beta(g(\phi Z, X)\xi - \eta(X)\phi Z), \phi Y \right) \\ &\quad - \left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2) \right) (g(\nabla_Z X, Y) - \eta(\nabla_Z X)\eta(Y)) \\ &\quad - S\left(\phi X, \alpha(g(Z, Y)\xi - \eta(Y)Z) + \beta(g(\phi Z, Y)\xi - \eta(Y)\phi Z) \right) \\ &\quad - \left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2) \right) (g(X, \nabla_Z Y) - \eta(X)\eta(\nabla_Z Y)). \end{aligned} \quad (4.4)$$

By virtue of (2.9) and (2.10) we obtain from (4.4)

$$\begin{aligned} &(\nabla_Z S)(\phi X, \phi Y) \\ &= \left(\frac{1}{2} dr(Z) + \nabla_Z(\xi\beta) - 2\alpha d\alpha(Z) + 2\beta d\beta(Z) \right) (g(X, Y) - \eta(X)\eta(Y)) \\ &\quad + \left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2) \right) \left(\nabla_Z g(X, Y) - (\nabla_Z \eta(X))\eta(Y) - \eta(X)(\nabla_Z \eta(Y)) \right) \\ &\quad + \alpha g(Z, X)((\phi Y)\beta + (\phi^2 Y)\alpha) \end{aligned} \quad (4.5)$$

$$\begin{aligned}
 & + \alpha\eta(X) \left(\left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2) \right) g(Z, \phi Y) - ((\phi Y)\beta + (\phi^2 Y)\alpha)\eta(Z) \right) \\
 & + \beta g(\phi Z, X)((\phi Y)\beta + (\phi^2 Y)\alpha) \\
 & + \beta\eta(X) \left(\left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2) \right) (g(Z, Y) - \eta(Z)\eta(Y)) \right) \\
 & - \left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2) \right) (g(\nabla_Z X, Y) - \eta(\nabla_Z X)\eta(Y)) \\
 & + \alpha g(Z, Y)((\phi X)\beta + (\phi^2 X)\alpha) \\
 & + \alpha\eta(Y) \left(\left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2) \right) g(Z, \phi X) - ((\phi X)\beta + (\phi^2 X)\alpha)\eta(Z) \right) \\
 & + \beta g(\phi Z, Y)((\phi X)\beta + (\phi^2 X)\alpha) \\
 & + \beta\eta(Y) \left(\left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2) \right) (g(Z, X) - \eta(Z)\eta(X)) \right) \\
 & - \left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2) \right) (g(X, \nabla_Z Y) - \eta(X)\eta(\nabla_Z Y)).
 \end{aligned}$$

The above relation can be written as

$$\begin{aligned}
 & (\nabla_Z S)(\phi X, \phi Y) \\
 & = \left(\frac{1}{2} dr(Z) + \nabla_Z(\xi\beta) - 2\alpha d\alpha(Z) + 2\beta d\beta(Z) \right) (g(X, Y) - \eta(X)\eta(Y)) \\
 & + \left(\frac{r}{2} + \xi\beta - (\alpha^2 - \beta^2) \right) \left(\nabla_Z g(X, Y) - (\nabla_Z \eta(X))\eta(Y) - \eta(X)(\nabla_Z \eta(Y)) \right. \\
 & + \alpha\eta(X)g(Z, \phi Y) + \beta\eta(X)(g(Z, Y) - \eta(Z)\eta(Y)) - g(\nabla_Z X, Y) \\
 & + \eta(\nabla_Z X)\eta(Y) + \alpha\eta(Y)g(Z, \phi X) + \beta\eta(Y)(g(Z, X) - \eta(Z)\eta(X)) \\
 & \left. - g(X, \nabla_Z Y) + \eta(X)\eta(\nabla_Z Y) \right) \tag{4.6} \\
 & + ((\phi Y)\beta + (\phi^2 Y)\alpha)(\alpha g(Z, X) + \beta g(\phi Z, X) - \alpha\eta(X)\eta(Z)) \\
 & + ((\phi X)\beta + (\phi^2 X)\alpha)(\alpha g(Z, Y) + \beta g(\phi Z, Y) - \alpha\eta(Y)\eta(Z)).
 \end{aligned}$$

Suppose that α and β are constants. Then using (2.7) in (4.6), we obtain

$$(\nabla_Z S)(\phi X, \phi Y) = \frac{1}{2} dr(Z)(g(X, Y) - \eta(X)\eta(Y)). \tag{4.7}$$

Hence, from (4.7) we can state the following:

THEOREM 4.1. *A 3-dimensional connected trans-Sasakian manifold of type (α, β) has η -parallel Ricci tensor if and only if the scalar curvature of the manifold is constant provided α and β are constants.*

From Theorem 3.1 and Theorem 4.1 we can state the following:

COROLLARY 4.1. *A 3-dimensional connected trans-Sasakian manifold of type (α, β) has η -parallel Ricci tensor if and only if it is locally ϕ -symmetric provided α and β are constants.*

5. THREE-DIMENSIONAL TRANS-SASAKIAN MANIFOLD WITH CONSTANT CURVATURE

Let M be a 3-dimensional compact connected trans-Sasakian manifold. If the manifold is of constant curvature then the Ricci tensor of type $(0, 2)$ of the manifold is given by

$$S(X, Y) = 2\lambda g(X, Y), \quad (5.1)$$

where λ is a constant. Putting $Y = \xi$ in (5.1) and using (2.9), we get

$$X\beta + (\phi X)\alpha + [2(\lambda - \alpha^2 + \beta^2) + \xi\beta]\eta(X) = 0. \quad (5.2)$$

For $X = \xi$, (5.2) yields

$$\xi\beta = -(\lambda - \alpha^2 + \beta^2). \quad (5.3)$$

By virtue of (5.2) and (5.3) it follows that

$$X\beta + (\phi X)\alpha + (\lambda - \alpha^2 + \beta^2)\eta(X) = 0. \quad (5.4)$$

The gradient of the function β is related to the exterior derivative $d\beta$ by the formula

$$d\beta(X) = g(\text{grad}\beta, X). \quad (5.5)$$

Using (5.5) in (5.4) we obtain

$$d\beta(X) + g(\text{grad}\alpha, \phi X) + (\lambda - \alpha^2 + \beta^2)\eta(X) = 0. \quad (5.6)$$

Differentiating (5.6) covariantly with respect to Y we get

$$\begin{aligned} (\nabla_Y d\beta)(X) + g(\nabla_Y \text{grad}\alpha, \phi X) + g(\text{grad}\alpha, (\nabla_Y \phi)X) \\ + Y(\beta^2 - \alpha^2)\eta(X) + (\lambda - \alpha^2 + \beta^2)(\nabla_Y \eta)(X) = 0. \end{aligned} \quad (5.7)$$

Interchanging X and Y in (5.7), we get

$$\begin{aligned}
 (\nabla_X d\beta)(Y) + g(\nabla_X \text{grad}\alpha, \phi Y) + g(\text{grad}\alpha, (\nabla_X \phi)Y) \\
 + X(\beta^2 - \alpha^2)\eta(Y) + (\lambda - \alpha^2 + \beta^2)(\nabla_X \eta)(Y) = 0.
 \end{aligned}
 \tag{5.8}$$

Subtracting (5.7) from (5.8) we get

$$\begin{aligned}
 g(\nabla_X \text{grad}\alpha, \phi Y) - g(\nabla_Y \text{grad}\alpha, \phi X) + ((\nabla_X \phi)Y - (\nabla_Y \phi)X)\alpha \\
 + [X(\beta^2 - \alpha^2)\eta(Y) - Y(\beta^2 - \alpha^2)\eta(X)] \\
 + (\lambda - \alpha^2 + \beta^2)((\nabla_X \eta)(Y) - (\nabla_Y \eta)(X)) = 0.
 \end{aligned}
 \tag{5.9}$$

From (2.7) and (2.4) we get

$$(\nabla_X \eta)(Y) - (\nabla_Y \eta)(X) = \alpha(\Phi(X, Y) - \Phi(Y, X)) = 2\alpha\Phi(X, Y).
 \tag{5.10}$$

Using (5.10) in (5.9) we have

$$\begin{aligned}
 g(\nabla_X \text{grad}\alpha, \phi Y) - g(\nabla_Y \text{grad}\alpha, \phi X) + ((\nabla_X \phi)Y - (\nabla_Y \phi)X)\alpha \\
 + [X(\beta^2 - \alpha^2)\eta(Y) - Y(\beta^2 - \alpha^2)\eta(X)] \\
 + 2(\lambda - \alpha^2 + \beta^2)\alpha\Phi(X, Y) = 0.
 \end{aligned}
 \tag{5.11}$$

Let $\{E_0, E_1, E_2\}$ be a local ϕ -basis, that is, an orthonormal frame such that $E_0 = \xi$ and $E_2 = \phi E_1$. In (2.5) putting $X = E_1, Y = E_2$ we get

$$\begin{aligned}
 (\nabla_{E_1} \phi)E_2 &= \alpha(g(E_1, E_2)\xi - \eta(E_2)E_1) + \beta(g(\phi E_1, E_2)\xi - \eta(E_2)\phi E_1) \\
 &= \beta g(\phi E_1, E_2)\xi = \beta\xi.
 \end{aligned}
 \tag{5.12}$$

Similarly,

$$(\nabla_{E_2} \phi)E_1 = -\beta\xi.
 \tag{5.13}$$

Now,

$$\Phi(E_1, E_2) = g(E_1, \phi E_2) = g(E_1, \phi^2 E_1) = -1.
 \tag{5.14}$$

In (5.11) putting $X = E_1$ and $Y = E_2$ and using (5.12), (5.13) and (5.14) we obtain

$$g(\nabla_{E_1} \text{grad}\alpha, E_1) + g(\nabla_{E_2} \text{grad}\alpha, E_2) = 2\beta\xi\alpha - 2\alpha(\lambda - \alpha^2 + \beta^2).
 \tag{5.15}$$

Also (2.8) can be written as

$$g(\text{grad}\alpha, \xi) = -2\alpha\beta.
 \tag{5.16}$$

Differentiating (5.16) covariantly with respect to ξ we get

$$g(\nabla_{\xi} \text{grad} \alpha, \xi) + g(\text{grad} \alpha, \nabla_{\xi} \xi) = -2\beta \xi \alpha - 2\alpha \xi \beta. \quad (5.17)$$

In view of (5.3) we can write the above relation as

$$g(\nabla_{\xi} \text{grad} \alpha, \xi) = -2\beta \xi \alpha + 2\alpha(\lambda - \alpha^2 + \beta^2). \quad (5.18)$$

From (5.15) and (5.18) we get $\Delta \alpha = 0$, where Δ is the Laplacian defined by

$$\Delta \alpha = \sum_{i=0}^2 g(\nabla_{E_i} \text{grad} \alpha, E_i).$$

Since M is compact we get α is constant.

Now let us consider the following two cases:

CASE-I: In this case we suppose that α is non-zero constant then by (2.8), $\beta = 0$ every where on M .

CASE-II: In this case let $\alpha = 0$. Then from (5.4)

$$X\beta + (\lambda + \beta^2)\eta(X) = 0,$$

that is,

$$g(\text{grad} \beta, X) + (\lambda + \beta^2)g(X, \xi) = 0.$$

Therefore,

$$\text{grad} \beta + (\lambda + \beta^2)\xi = 0. \quad (5.19)$$

Differentiating (5.19) covariantly with respect to X we have

$$\nabla_X \text{grad} \beta + (X\beta^2)\xi + (\lambda + \beta^2)\nabla_X \xi = 0.$$

Using (2.6) we get from above

$$\nabla_X \text{grad} \beta + (X\beta^2)\xi + (\lambda + \beta^2)(-\alpha \phi X + \beta(X - \eta(X)\xi)) = 0.$$

Now taking inner product of the above equation with X , we have

$$\begin{aligned} g(\nabla_X \text{grad} \beta, X) &= -g((X\beta^2)\xi, X) \\ &\quad - (\lambda + \beta^2)(g(-\alpha \phi X, X) + \beta g(X - \eta(X)\xi, X)). \end{aligned} \quad (5.20)$$

Therefore putting $X = E_i$ and taking summation over $i, i = 0, 1, 2$, we get from above

$$\Delta\beta = -2\beta(\xi\beta + \lambda + \beta^2). \tag{5.21}$$

For $\alpha = 0$, (5.3) yields $\xi\beta = -(\lambda + \beta^2)$, which in view of (5.21) gives $\Delta\beta = 0$. Hence $\beta = \text{constant}$, M being compact. This leads to the following:

THEOREM 5.1. *If a 3-dimensional compact connected trans-Sasakian manifold is of constant curvature then it is either α -Sasakian or β -Kenmotsu.*

6. EXAMPLE OF A LOCALLY ϕ -SYMMETRIC THREE-DIMENSIONAL TRANS-SASAKIAN MANIFOLD

We consider the three-dimensional manifold $M = \{(x, y, z) \in \mathbb{R}^3, z \neq 0\}$, where (x, y, z) are the standard coordinates in \mathbb{R}^3 . The vector fields

$$e_1 = z \frac{\partial}{\partial x}, \quad e_2 = z \frac{\partial}{\partial y}, \quad e_3 = z \frac{\partial}{\partial z}$$

are linearly independent at each point of M . Let g be the Riemannian metric defined by

$$g(e_1, e_3) = g(e_2, e_3) = g(e_1, e_2) = 0, \quad g(e_1, e_1) = g(e_2, e_2) = g(e_3, e_3) = 1.$$

Let η be the 1-form defined by $\eta(Z) = g(Z, e_3)$ for any $Z \in \chi(M)$. Let ϕ be the (1,1) tensor field defined by $\phi(e_1) = -e_2, \phi(e_2) = e_1, \phi(e_3) = 0$. Then using the linearity of ϕ and g we have

$$\eta(e_3) = 1, \quad \phi^2 Z = -Z + \eta(Z)e_3, \quad g(\phi Z, \phi W) = g(Z, W) - \eta(Z)\eta(W),$$

for any $Z, W \in \chi(M)$. Thus for $e_3 = \xi, (\phi, \xi, \eta, g)$ defines an almost contact metric structure on M . Now, by direct computations we obtain

$$[e_1, e_2] = 0, \quad [e_2, e_3] = -e_2, \quad [e_1, e_3] = -e_1.$$

The Riemannian connection ∇ of the metric g is given by the Koszul's formula which is

$$2g(\nabla_X Y, Z) = Xg(Y, Z) + Yg(Z, X) - Zg(X, Y) - g(X, [Y, Z]) - g(Y, [X, Z]) + g(Z, [X, Y]). \tag{6.1}$$

Using (6.1) we have

$$\begin{aligned} 2g(\nabla_{e_1} e_3, e_1) &= 2g(-e_1, e_1), \\ 2g(\nabla_{e_1} e_3, e_2) &= 0 = 2g(-e_1, e_2), \\ 2g(\nabla_{e_1} e_3, e_3) &= 0 = 2g(-e_1, e_3). \end{aligned}$$

Hence, $\nabla_{e_1} e_3 = -e_1$. Similarly, $\nabla_{e_2} e_3 = -e_2$ and $\nabla_{e_3} e_3 = 0$.
(6.1) further yields

$$\begin{aligned} \nabla_{e_1} e_2 &= 0, & \nabla_{e_1} e_1 &= e_3, \\ \nabla_{e_2} e_2 &= e_3, & \nabla_{e_2} e_1 &= 0, \\ \nabla_{e_3} e_2 &= 0, & \nabla_{e_3} e_1 &= 0. \end{aligned}$$

We see that

$$\begin{aligned} (\nabla_{e_1} \phi)e_1 &= \nabla_{e_1} \phi e_1 - \phi \nabla_{e_1} e_1 = -\nabla_{e_1} e_2 - \phi e_3 = -\nabla_{e_1} e_2 = 0 \\ &= 0(g(e_1, e_1)e_3 - \eta(e_1)e_1) - 1(g(\phi e_1, e_1)e_3 - \eta(e_1)\phi e_1). \end{aligned} \quad (6.2)$$

$$\begin{aligned} (\nabla_{e_1} \phi)e_2 &= \nabla_{e_1} \phi e_2 - \phi \nabla_{e_1} e_2 = \nabla_{e_1} e_1 - 0 = e_3 \\ &= 0(g(e_1, e_2)e_3 - \eta(e_2)e_1) - 1(g(\phi e_1, e_2)e_3 - \eta(e_2)\phi e_1). \end{aligned} \quad (6.3)$$

$$\begin{aligned} (\nabla_{e_1} \phi)e_3 &= \nabla_{e_1} \phi e_3 - \phi \nabla_{e_1} e_3 = 0 + \phi e_1 = -e_2 \\ &= 0(g(e_1, e_3)e_3 - \eta(e_3)e_1) - 1(g(\phi e_1, e_3)e_3 - \eta(e_3)\phi e_1). \end{aligned} \quad (6.4)$$

By (6.2), (6.3) and (6.4) we see that the manifold satisfies (2.5) for $X = e_1$, $\alpha = 0$, $\beta = -1$, and $e_3 = \xi$. Similarly it can be shown that for $X = e_2$ and $X = e_3$ the manifold also satisfies (2.5) for $\alpha = 0$, $\beta = -1$, and $e_3 = \xi$. Hence the manifold is a trans-Sasakian manifold of type $(0, -1)$. With the help of the above results it can be verified that

$$\begin{aligned} R(e_1, e_2)e_3 &= 0, & R(e_2, e_3)e_3 &= -e_2, & R(e_1, e_3)e_3 &= -e_1, \\ R(e_1, e_2)e_2 &= -e_1, & R(e_2, e_3)e_2 &= e_3, & R(e_1, e_3)e_2 &= 0, \\ R(e_1, e_2)e_1 &= e_2, & R(e_2, e_3)e_1 &= 0, & R(e_1, e_3)e_1 &= e_3. \end{aligned}$$

From which it follows that $\phi^2(\nabla_W R)(X, Y)Z = 0$. Hence the 3-dimensional trans-Sasakian manifold is locally ϕ -symmetric.

Also from the above expressions of the curvature tensor we obtain the scalar curvature $r = -3$. Hence we note that here α , β and r all are constants. Hence from Theorem 3.1 it follows that the manifold under consideration is locally ϕ -symmetric.

ACKNOWLEDGEMENTS

The authors are thankful to the referee for his valuable suggestions and remarks in the improvement of the paper.

REFERENCES

- [1] D. E. BLAIR, "Contact Manifolds in Riemannian Geometry", Lecture Notes in Mathematics, Vol. 509, Springer-Verlag, Berlin-New York, 1976.
- [2] D. E. BLAIR, J. A. OUBIÑA, Conformal and related changes of metric on the product of two almost contact metric manifolds, *Publ. Mat.* **34** (1) (1990), 199–207.
- [3] D. CHINEA, C. GONZALES, A classification of almost contact metric manifolds, *Ann. Mat. Pura Appl. (4)* **156** (1990), 15–36.
- [4] D. CHINEA, C. GONZALES, Curvature relations in trans-Sasakian manifolds, (Spanish), in "Proceedings of the XIIth Portuguese-Spanish Conference on Mathematics, Vol. II, (Portuguese), Braga, 1987," Univ. Minho, Braga, (1987), 564–571.
- [5] U. C. DE, On ϕ -symmetric Kenmotsu manifolds, *Int. Electron. J. Geom.* **1** (1) (2008), 33–38.
- [6] U. C. DE, M. M. TRIPATHI, Ricci tensor in 3-dimensional trans-Sasakian manifolds, *Kyungpook Math. J.* **43** (2) (2003), 247–255.
- [7] A. GRAY, L. M. HERVELLA, The sixteen classes of almost Hermitian manifolds and their linear invariants, *Ann. Mat. Pura Appl. (4)* **123** (1980), 35–58.
- [8] D. JANSSENS, L. VANHECKE, Almost contact structures and curvature tensors, *Kodai Math. J.* **4** (1) (1981), 1–27.
- [9] M. KON, Invariant submanifolds in Sasakian manifolds, *Math. Ann.* **219** (3) (1976), 277–290.
- [10] J. C. MARRERO, The local structure of trans-Sasakian manifolds, *Ann. Mat. Pura Appl. (4)* **162** (1992), 77–86.
- [11] J. C. MARRERO, D. CHINEA, On trans-Sasakian manifolds, (Spanish), in "Proceedings of the XIVth Spanish-Portuguese Conference on Mathematics, Vol. I-III, (Spanish), Puerto de la Cruz, 1989", Univ. La Laguna, La Laguna, 1990, 655–659.
- [12] R. S. MISHRA, Almost contact metric manifolds, Monograph 1, Tensor Society of India, Lucknow, 1991.
- [13] J. A. OUBIÑA, New classes of almost contact metric structures, *Publ. Math. Debrecen* **32** (3-4) (1985), 187–193.
- [14] T. TAKAHASHI, Sasakian ϕ -symmetric spaces, *Tohoku Math. J. (2)* **29** (1) (1977), 91–113.
- [15] M. M. TRIPATHI, Trans-Sasakian manifolds are generalized quasi-Sasakian, *Nepali Math. Sci. Rep.* **18** (1-2) (1999/2000), 11–14.