



Advances in flow cytometry in basic and applied equine andrology

Fernando J. Peña^{1,3}, Patricia Martín-Muñoz¹, Cristina Ortega-Ferrusola²

¹Laboratory of Equine Reproduction and Equine Spermatology, Veterinary Teaching Hospital, University of Extremadura, Cáceres, Spain.

²Reproduction and Obstetrics Department of Animal Medicine and Surgery, University of León, Spain.

Abstract

The aim of this review is to present the current probes available that assess different compartments and functions of stallion spermatozoa, including assays to investigate the functionality of the membranes, nucleus and mitochondria, and to study cell signaling in this particular cell. New multi-parametric protocols for the assessment of stallion sperm, recently developed in the laboratory of the authors, will also be presented. The potential clinical applicability of diagnostic tests based on flow cytometry will also be discussed.

Keywords: fluorescent probes, sperm, stallion.

Introduction

Particularly the last decade of the current century, has been witness to intensive research in sperm biology. Consequently, a better understanding of sperm function in relation to clinical andrology and sperm biotechnologies (Peña *et al.*, 2011, 2015) has arisen. Relevant advances in stallion sperm biology include, among others, the following: the understanding of osmotic shock (Ball and Vo, 2001; Pommer *et al.*, 2002; Ball, 2008) and its implications on cryopreservation, which promoted the development of new protocols based on more permeant cryoprotectants (Oldenhof *et al.*, 2010, 2012, 2013; Hoffmann *et al.*, 2011; Pukazhenthil *et al.*, 2014); advances in understanding the role of reactive oxygen species (ROS; Gibb *et al.*, 2014, 2015; Varner *et al.*, 2015; Gibb and Aitken, 2016); and the development of practical methods for stallion sperm separation and selection through colloidal centrifugation (Waite *et al.*, 2008; Johannisson *et al.*, 2009; Morrell *et al.*, 2009a, b; Edmond *et al.*, 2012; Crespo *et al.*, 2013; Ponthier *et al.*, 2013). More recently, the understanding of sperm bioenergetics and mitochondrial functionality have become two hot topics in stallion andrology. Beside these advances, better tools for sperm assessment have been developed, in which flow cytometry has played a major role. The aim of this review is to present a rapid summary of the current probes available to assess stallion sperm and describe new protocols for the assessment of stallion sperm, including those recently developed in the laboratory of the authors. The potential clinical applicability of a

diagnostic test based on flow cytometry will also be discussed. Interestingly, these assays have been recently supported with field fertility data (Barrier Battut *et al.*, 2016).

Basic principles of flow cytometry applied to sperm analysis

Flow cytometry measures multiple parameters of cells that rapidly flow in a stream through a system of photonic receptors. The properties measured include the size of the spermatozoa in the forward scatter detector (FSC), the complexity in the side scatter detector (SSC) and the relative fluorescence intensity in fluorescence detectors (FL). These characteristics are detected using a fluidic and optical to electronic coupling system that records how each individual spermatozoon or other particle presents in the sample, scatters incident laser light, and emits fluorescence. In the flow cytometer, spermatozoa are carried to the laser interrogation point in a fluid stream (sheath fluid). When they pass through the laser intercept, they scatter laser light, and any fluorescent molecules present are excited and emit light in different wavelengths. Appropriately positioned lenses collect the scatter and fluorescent light. A combination of beam splitters and filters steer fluorescence to detectors that produce electronic signals proportional to the optical signals striking them. List mode data are collected on every single spermatozoon and stored in the computer; these data are analyzed and provide information about subpopulations in the sample and are displayed graphically in histograms and dot plots. Fluorescent compounds are used to study stallion sperm functionality. A fluorescent compound absorbs light energy over a range of wavelengths characteristic for each. This absorption of light causes an electron in the fluorescent compound to be raised to a higher energy level; the electron quickly returns to the ground state, releasing the excess energy as a photon. This transition of the energy is termed fluorescence. The range of wavelengths in which a fluorescent compound can be excited is called absorption spectrum, while the range of wavelengths emitted is called emission spectrum. Ideally, the light produced by emission should be different from the light used for excitation, and this difference is known as the *Stokes Shift*. The wavelength of emission is longer than the wavelength of

³Corresponding author: fjuanpega@unex.es
Phone: +34(927)25-7167; Fax: +34(927)25-7102
Received: August 30, 2016
Accepted: September 19, 2016



excitation because typically more energy is used to excite the electrons of the fluorochrome than the energy released (as light) when the electrons return to the resting state. For example, a commonly used fluorochrome, fluorescein isothiocyanate (FICT), absorbs light in the range 400-550 nm, with a peak or maximum excitation at 490 nm (the laser is used to excite a particular dye, the blue laser (488 nm) in this case), and emits in the range 475-700 nm, with a peak at 525 nm (green spectrum). This range of wavelengths determines the filters and the channels (fluorescence channels FL) of detection to be used. Combining different fluorochromes with multiple wavelengths of excitation and emission allows multiple and simultaneous measurements, however compensation for spectral overlap has to be considered and carefully managed. When two or more dyes are used simultaneously, there is a chance that their emission profiles will coincide, making measurement of the true fluorescence for each one difficult. This outcome can be avoided by using dyes at distant positions in the spectrum; for example, a dye that is excited with the violet laser (405 nm) and a dye excited with the red laser (647 nm). However, using dyes at distant positions is not always possible, and a process called fluorescence compensation is applied. This process calculates how much interference, as a percentage, a fluorochrome will have in a channel that was not assigned specifically to measure it. The design of an experiment in the flow cytometer implies careful selection of probes suitable for each particular cytometer, identification of potential spectral overlap among probes, use of proper controls for positive and negative populations (unstained sample), and controls for compensation (single stained samples in which there is a stained and unstained population for each dye to be used in the experiment). Depending on each particular experiment, other controls can be necessary, including fluorescence minus one (FMO) controls, isotype controls or secondary antibody only controls.

The sperm membrane: integrity, permeability, fluidity, and functionality

Traditional assessment of the sperm membrane has focused on the physical integrity using dye exclusion tests. Classical combinations of fluorescent probes for this purpose include the combination of SYBR-14 and propidium iodide (PI). This combination of probes requires the blue laser for excitation (488 nm) and provides two wavelengths of emission; green for live sperm (521 nm SYBR-14) and red for dead sperm (635 nm PI). This combination of probes allows the rapid discrimination of debris (because both are DNA binding probes), and both probes are excited with the blue laser (488 nm). Spillover between emission wavelengths of both probes can occur (521 nm for SYBR-14 and 635 nm for PI), and proper fluorescence

compensation has to be established in order to use this probe pair because SYBR-14 still has, on average, a 6% emission at 635 nm. Furthermore, staining with SYBR-14/PI discriminates only between live and dead sperm and does not expose initial states of membrane damage. Additionally, doublets have to be identified to correctly interpret this assay. Alternatively, Hoechst 33342 and PI (Plaza Davila *et al.*, 2015) also allow the rapid discrimination of debris and have the advantage that spillover is unlikely due to the distinct excitation and emission spectra of these probes. However, H33342 needs a violet or ultraviolet laser for excitation. Detection of more subtle changes in the sperm membrane requires the use of other probes. Fluidity of sperm membranes can be assessed with merocyanine 540 (da Silva *et al.*, 2011), and subtle increases in the permeability of the plasma membrane can be detected with YoPro-1 (Gallardo Bolanos *et al.*, 2012, 2014a). Yo Pro-1 is routinely used in the authors' laboratory in combination with PI; H33342 is also incorporated to sort debris. These combinations allow the detection of changes in sperm membranes at much earlier stages than SYBR-14 and correlate better with motility and sperm velocities (Gallardo Bolanos *et al.*, 2012).

Recently, new fixable fluorescent dyes have become available in multiple colors, which facilitate experiments with multiple spectra in fixed sperm. These probes are based upon the reaction of fluorescent reactive dye with cellular amines. These are proprietary dyes that can permeate the compromised membranes of necrotic cells and react with free amines both in the cytoplasm and on the cell surface, resulting in intense fluorescent staining. In contrast, only the cell surface amines of intact cells are available to react with the dye, resulting in relatively dim staining. The discrimination is maintained following formalin-fixation of the sample under conditions that inactivate pathogens. Moreover, these assays use only one channel of the flow cytometer, leaving the other channels available for multicolor panels. The potential advantage of these dyes is the ability to process and stain the samples at locations remote to the flow cytometer.

The evaluation of the sperm's ability to undergo the acrosome reaction in response to an agonist challenge, is useful in cases of infertility in certain thoroughbred lines. This assay, the Acrosomal Responsiveness Assay (ARA; Johnson *et al.*, 2008; Vaner, 2008), evaluates the ability of the acrosome to react when challenged with the Ca^{2+} ionophore, A23197. Common probes to assess acrosomal integrity, either in a basal status after a challenge, are those which recognize targets inside the acrosome, including specific lectins (*Pisum sativum* agglutinin PSA, and *Arachis hypogea* agglutinin PNA) that bind to glucosidic residues in different parts of the acrosomal membrane. The acrosome has also been monitored in human sperm with anti-CD46 antibodies (Carver-Ward *et al.*, 1994; Grunewald *et al.*, 2008).



Mitochondria and stallion sperm functionality

The mitochondria of spermatozoa are increasingly studied in both basic and applied andrology (Gibb *et al.*, 2014; Peña *et al.*, 2015; Plaza Davila *et al.*, 2015). Stallion spermatozoa are highly dependent on mitochondrial production of ATP, and mitochondrial malfunction leads rapidly to sperm senescence and death. Stallion spermatozoa have particularly active mitochondria, and as a result, they generate large amounts of reactive oxygen species (ROS; Gibb *et al.*, 2014; Plaza Davila *et al.*, 2015). Sperm mitochondria are sensitive indicators of sperm stress during processes such as cooling and cryopreservation (Ortega-Ferrusola *et al.*, 2008, 2009a). Two common probes are used to assess stallion mitochondrial function by flow cytometry. The probe 5,5', 6,6'-tetrachloro-1,1', 3,3' tetraethylbenzimidazolyl carbocyanine iodide (JC-1) forms multimeric aggregates in mitochondria with high membrane potential (active mitochondria). These aggregates emit the high orange wavelength of 590 nm when excited at 488 nm. In mitochondria with low membrane potential (inactive mitochondria), JC-1 forms monomers that emit in the green wavelength (525 to 530 nm) when excited at 488 nm (Garner and Thomas, 1999; Gravance *et al.*, 2000). Recently, mitotracker dyes (Gallon *et al.*, 2006; Sousa *et al.*, 2011) have become available in multiple colors and provide colorful alternatives to be used in multicolor experiments. Both JC-1 and Mitotracker deep red have been recently used in our laboratory (Gallardo Bolanos *et al.*, 2014a). These probes measure different aspects of mitochondrial function; JC-1 reflects mitochondrial membrane potential, while mitotracker deep red passively diffuses across membranes and binds to thiols in active mitochondria (Peña *et al.*, 2016).

The sperm DNA

The sperm chromatin structure assay (SCSA) has been extensively used. In stallion andrology, this assay has successfully discriminated between stallions of low, high, and average fertility (Love and Kenney, 1998; Love, 2005). In spite of the importance of DNA, the origin of damage to sperm DNA is still largely ignored. In human andrology, it is becoming clear that two major factors are associated with damage to sperm DNA: oxidative stress and protamination of the spermatozoa. These two features are strongly linked because defective protamination renders spermatozoa more susceptible to oxidative damage (Aitken and De Iuliis, 2010; Aitken *et al.*, 2013, 2014; Gavriliouk and Aitken, 2015). Recent research from our laboratory shows evidence indicating that DNA damage in stallion spermatozoa is oxidative as well (Balao da Silva *et al.*, 2014). Oxidative stress can be assessed using specific antibodies against the oxidized form of guanine; 8-oxoguanine in fixed, permeabilized samples (Balao da

Silva *et al.*, 2016).

The stallion sperm: a redox regulated cell

Reactive oxygen species (ROS) as by-products of various metabolic processes may have detrimental effects, but ROS may also be important regulators of cellular functions (Stowe and Camara, 2009). These are chemical species formed after incomplete reduction of oxygen and include the superoxide anion ($O_2^{\bullet-}$), hydrogen peroxide (H_2O_2), and the hydroxyl radical (OH^{\bullet}). Superoxide anion ($O_2^{\bullet-}$) can be generated at different points within the mitochondrial electron transport chain (ETC) by univalent reduction of oxygen. Most superoxide is converted to H_2O_2 by superoxide dismutase inside and outside of the mitochondrial matrix, and superoxide in low and controlled amounts exerts important regulatory cellular functions. Excessive H_2O_2 can combine with Fe^{2+} to form reactive hydroxyl radical (OH^{\bullet} ; Shen *et al.*, 1992). Superoxide is short lived ($t_{1/2} = 1$ ms) and cell impermeable, while H_2O_2 is more stable and cell permeable. In the presence of nitric oxide (NO^{\bullet}), $O_2^{\bullet-}$ forms the reactant peroxynitrite ($ONOO^{\bullet}$), and $ONOOH$ induced nitrosylation of proteins, DNA, and lipids can modify their structure and function (Stowe and Camara, 2009). NO^{\bullet} is synthesized through the conversion of L-arginine to L-citrulline by nitric oxide synthase (NOS). These enzymes are present in stallion spermatozoa, possibly as sperm specific isoforms (Ortega-Ferrusola *et al.*, 2009b). Numerous studies indicate that ROS are important regulators of sperm function (Zini *et al.*, 1995; Aitken *et al.*, 1997; de Lamirande and Gagnon, 2002, 2003; De Lamirande and Lamothe, 2009), and ROS become detrimental only if homeostasis is lost (Peña *et al.*, 2015). Moreover, recent evidence suggests that stallion sperm mitochondria produce significant amounts of NO (Ortega-Ferrusola *et al.*, 2009b). Nitric oxide has a relatively long half-life (1 s) and is more reactive than $O_2^{\bullet-}$. Controlled ROS production occurs during capacitation in spermatozoa (Agarwal *et al.*, 2014). This controlled production triggers signaling pathways initiated by an increase in cyclic adenosine 3'-5' monophosphate (cAMP). Increased cAMP activates protein kinase A (PKA), and the subsequent phosphorylation of extracellular regulated -kinase-like proteins and finally tyrosine phosphorylation of proteins in the fibrous sheath of the spermatozoa, leading to sperm hyperactivation. Numerous assays have been developed to assess oxidative stress and production of ROS in the stallion spermatozoa (Baumber *et al.*, 2002; Sabeur and Ball, 2006; Burnaugh *et al.*, 2007; Gibb *et al.*, 2014; Plaza Davila *et al.*, 2015). Flow cytometry can be used to detect specific reactive oxygen species (ROS), reactive nitrogen species (RNS) and the consequences of perturbed ROS homeostasis, such as lipid peroxidation, DNA oxidation, increased membrane permeability and protein oxidation. Reactive oxygen species can be



detected using different probes. The superoxide indicator dihydroethidium, also called hydroethidine, exhibits blue-fluorescence in the cytosol until oxidized, where it intercalates within the cell's DNA, staining its nucleus a bright fluorescent red. Mitosox Red™ is used to specifically detect mitochondrial $O_2^{\bullet-}$. Hydrogen peroxide can be detected using dichlorodihydrofluorescein diacetate, although this probe is not highly specific for H_2O_2 . Recently, molecules such as aryl boronate have been described and appear highly specific for the detection of H_2O_2 in spermatozoa (Purdey *et al.*, 2015). Other probes have been recently introduced. The cellrox sensors™ are available in different colors. These may be fixed after staining, facilitating their use in multicolor panels. In our laboratory, the CellROX deep red™ has been used in multicolor experiments, especially to detect mitochondrial $O_2^{\bullet-}$ and the hydroxyl radical (OH^{\bullet} ; Gallardo Bolanos *et al.*, 2014b; Plaza Davila *et al.*, 2015). It is extremely important to consider the conditions in which the assay is performed. High Cell ROX deep red™ fluorescence may indicate either mitochondrial activity or real oxidative stress. The particular dependence of stallion spermatozoa on oxidative phosphorylation to generate ATP (Plaza Davila *et al.*, 2015) may reflect the apparent paradoxical relationship between ROS and sperm functionality due to increased electron leakage and thus increased $O_2^{\bullet-}$ production (Ortega-Ferrusola *et al.*, 2010; Gibb *et al.*, 2014; Yeste *et al.*, 2015).

The oxidation of the plasma membrane leads to increased membrane permeability (Christova *et al.*, 2004) that can be monitored with YoPro-1 (Ortega-Ferrusola *et al.*, 2008; da Silva *et al.*, 2011; Gallardo Bolanos *et al.*, 2012; Garcia *et al.*, 2012; Gibb *et al.*, 2014). Oxidation of DNA can be monitored using antibodies against the oxidized form of guanine. Peroxidation of sperm membranes can be detected with the probe BODIPY® 581/591 C11. This probe emits orange-red fluorescence in the non-oxidized state, shifting to green fluorescence when peroxidized (Ball and Vo, 2002; Ortega-Ferrusola *et al.*, 2009c). Lipid peroxidation is also monitored by detection of 4-hydroxynonenal (4-HNE) using specific antibodies, a product from the oxidation of sperm-membrane lipids (Aitken *et al.*, 2012; Gibb *et al.*, 2014). This assay is considered a reliable indicator of ROS imbalance, and specific protocols for stallion spermatozoa have been recently published (Martin Munoz *et al.*, 2015).

Sperm senescence

Senescent spermatozoa express active caspase 3 (Amann, 2010; Aitken and Baker, 2013; Aitken *et al.*, 2015). Depending of the presence of pro-survival factors, caspase 3 remains inactive due to the phosphorylation of protein kinase B (PKB or Akt;

Gallardo Bolanos *et al.*, 2014a). If pro-survival factors are lost or oxidative stress reaches a threshold, caspase 3 is activated and sperm senescence and death are triggered (Gallardo Bolanos *et al.*, 2014b). Cryopreservation triggers this phenomenon, and surviving spermatozoa experience accelerated senescence (Thomas *et al.*, 2006; Ortega-Ferrusola *et al.*, 2008, 2009a). Active caspase 3 can be detected using CellEvent™ Caspase-3 Green Detection Reagent, which consists of a four-amino-acid peptide (DEVD) conjugated to a nucleic acid-binding dye. This cell-permeant substrate is intrinsically non-fluorescent because the DEVD peptide inhibits the ability of the dye to bind to DNA. After activation of caspase-3 in apoptotic cells, the DEVD peptide is cleaved, enabling the dye to bind to DNA and produce a bright, fluorogenic response with an absorption/emission maximum of ~502/530 nm.

Concluding remarks

Flow cytometry is a powerful tool in andrology, allowing the rapid and simultaneous assessment of multiple sperm compartments and functions in thousands of spermatozoa in a few seconds. Recent data also suggests that the data generated are powerful forecasts of field fertility and an extremely important tool for quality control in stallion stations.

Acknowledgments

The authors received financial support for their studies from the Ministerio de Economía y Competitividad-FEDER, Madrid, Spain, grant AGL2013-43211-R, Junta de Extremadura-FEDER (GR 15029). PMM is supported by a pre-doctoral grant from the Ministerio de Educación, Cultura y Deporte, Madrid, Spain, FPU13/03991. COF is supported by a post-doctoral grant from the Ministerio de Economía y Competitividad “Juan de la Cierva” IJCI-2014-21671.

References

- Agarwal A, Virk G, Ong C, du Plessis SS. 2014. Effect of oxidative stress on male reproduction. *World J Mens Health*, 32:1-17.
- Aitken RJ, Fisher HM, Fulton N, Gomez E, Knox W, Lewis B, Irvine S. 1997. Reactive oxygen species generation by human spermatozoa is induced by exogenous NADPH and inhibited by the flavoprotein inhibitors diphenylene iodonium and quinacrine. *Mol Reprod Dev*, 47:468-482.
- Aitken RJ, De Iuliis GN. 2010. On the possible origins of DNA damage in human spermatozoa. *Mol Hum Reprod*, 16:3-13.
- Aitken RJ, Whiting S, De Iuliis GN, McClymont S, Mitchell LA, Baker MA. 2012. Electrophilic aldehydes generated by sperm metabolism activate mitochondrial



- reactive oxygen species generation and apoptosis by targeting succinate dehydrogenase. *J Biol Chem*, 287:33048-33060.
- Aitken RJ, Baker MA.** 2013. Causes and consequences of apoptosis in spermatozoa; contributions to infertility and impacts on development. *Int J Dev Biol*, 57:265-272.
- Aitken RJ, Bronson R, Smith TB, De Iuliis GN.** 2013. The source and significance of DNA damage in human spermatozoa; a commentary on diagnostic strategies and straw man fallacies. *Mol Hum Reprod*, 19:475-485.
- Aitken RJ, Smith TB, Jobling MS, Baker MA, De Iuliis GN.** 2014. Oxidative stress and male reproductive health. *Asian J Androl*, 16:31-38.
- Aitken RJ, Baker MA, Nixon B.** 2015. Are sperm capacitation and apoptosis the opposite ends of a continuum driven by oxidative stress? *Asian J Androl*, 17:633-639.
- Amann RP.** 2010. Tests to measure the quality of spermatozoa at spermiation. *Asian J Androl*, 12:71-78.
- Balao da Silva CM, Ortega Ferrusola C, Gallardo Bolanos JM, Plaza Davila M, Martin-Munoz P, Morrell JM, Rodriguez-Martinez H, Pena FJ.** 2014. Effect of overnight staining on the quality of flow cytometric sorted stallion sperm: comparison with traditional protocols. *Reprod Domest Anim*, 49:1021-1027.
- Balao da Silva CM, Ortega-Ferrusola C, Morrell JM, Rodriguez Martinez H, Pena FJ.** 2016. Flow cytometric chromosomal sex sorting of stallion spermatozoa induces oxidative stress on mitochondria and genomic DNA. *Reprod Domest Anim*, 51:18-25.
- Ball BA, Vo A.** 2001. Osmotic tolerance of equine spermatozoa and the effects of soluble cryoprotectants on equine sperm motility, viability, and mitochondrial membrane potential. *J Androl*, 22:1061-1069.
- Ball BA, Vo A.** 2002. Detection of lipid peroxidation in equine spermatozoa based upon the lipophilic fluorescent dye C11-BODIPY581/591. *J Androl*, 23:259-269.
- Ball BA.** 2008. Oxidative stress, osmotic stress and apoptosis: impacts on sperm function and preservation in the horse. *Anim Reprod Sci*, 107:257-267.
- Barrier Battut I, Kempfer A, Becker J, Lebailly L, Camugli S, Chevrier L.** 2016. Development of a new fertility prediction model for stallion semen, including flow cytometry. *Theriogenology*, 86:1111-1131.
- Baumber J, Vo A, Sabeur K, Ball BA.** 2002. Generation of reactive oxygen species by equine neutrophils and their effect on motility of equine spermatozoa. *Theriogenology*, 57:1025-1033.
- Burnaugh L, Sabeur K, Ball BA.** 2007. Generation of superoxide anion by equine spermatozoa as detected by dihydroethidium. *Theriogenology*, 67:580-589.
- Carver-Ward JA, Jaroudi KA, Einspinner M, Parhar RS, al-Sedairy ST, Sheth KV.** 1994. Pentoxifylline potentiates ionophore (A23187) mediated acrosome reaction in human sperm: flow cytometric analysis using CD46 antibody. *Hum Reprod*, 9:71-76.
- Christova Y, James PS, Jones R.** 2004. Lipid diffusion in sperm plasma membranes exposed to peroxidative injury from oxygen free radicals. *Mol Reprod Dev*, 68:365-372.
- Crespo F, Gosalvez J, Gutierrez-Cepeda L, Serres C, Johnston SD.** 2013. Colloidal centrifugation of stallion semen results in a reduced rate of sperm DNA fragmentation. *Reprod Domest Anim*, 48:e23-25.
- da Silva CM, Macias-Garcia B, Miro-Moran A, Gonzalez-Fernandez L, Morillo-Rodriguez A, Ortega-Ferrusola C, Gallardo-Bolaños JM, Stilwell G, Tapia JA, Peña FJ.** 2011. Melatonin reduces lipid peroxidation and apoptotic-like changes in stallion spermatozoa. *J Pineal Res*, 51:172-179.
- de Lamirande E, Gagnon C.** 2002. The extracellular signal-regulated kinase (ERK) pathway is involved in human sperm function and modulated by the superoxide anion. *Mol Hum Reprod*, 8:124-135.
- de Lamirande E, Gagnon C.** 2003. Redox control of changes in protein sulfhydryl levels during human sperm capacitation. *Free Radic Biol Med*, 35:1271-1285.
- de Lamirande E, Lamothe G.** 2009. Reactive oxygen-induced reactive oxygen formation during human sperm capacitation. *Free Radic Biol Med*, 46:502-510.
- Edmond AJ, Brinsko SP, Love CC, Blanchard TL, Teague SR, Varner DD.** 2012. Effect of centrifugal fractionation protocols on quality and recovery rate of equine sperm. *Theriogenology*, 77:959-966.
- Gallardo Bolanos JM, Miro Moran A, Balao da Silva CM, Morillo Rodriguez A, Plaza Davila M, Aparicio IM, Tapia JA, Ortega Ferrusola C, Peña FJ.** 2012. Autophagy and apoptosis have a role in the survival or death of stallion spermatozoa during conservation in refrigeration. *PLoS One*, 7:e30688.
- Gallardo Bolanos JM, Balao da Silva CM, Martin Munoz P, Morillo Rodriguez A, Plaza Davila M, Rodriguez-Martinez H, Aparicio IM, Tapia JA, Ortega Ferrusola C, Peña FJ.** 2014a. Phosphorylated AKT preserves stallion sperm viability and motility by inhibiting caspases 3 and 7. *Reproduction*, 148:221-235.
- Gallardo Bolanos JM, Balao da Silva C, Martin Munoz P, Plaza Davila M, Ezquerro J, Aparicio IM, Tapia JA, Ortega Ferrusola C, Peña FJ.** 2014b. Caspase activation, hydrogen peroxide production and Akt dephosphorylation occur during stallion sperm senescence. *Reprod Domest Anim*, 49:657-664.
- Gallon F, Marchetti C, Jouy N, Marchetti P.** 2006. The functionality of mitochondria differentiates human spermatozoa with high and low fertilizing capability. *Fertil Steril*, 86:1526-1530.
- Garcia BM, Moran AM, Fernandez LG, Ferrusola CO, Rodriguez AM, Bolanos JM, da Silva CM, Martínez HR, Tapia JA, Peña FJ.** 2012. The mitochondria of stallion spermatozoa are more sensitive than the plasmalemma to osmotic-induced stress: role of c-Jun N-terminal kinase (JNK) pathway. *J Androl*,



33:105-113.

Garner DL, Thomas CA. 1999. Organelle-specific probe JC-1 identifies membrane potential differences in the mitochondrial function of bovine sperm. *Mol Reprod Dev*, 53:222-229.

Gavrieliouk D, Aitken RJ. 2015. Damage to sperm DNA mediated by reactive oxygen species: its impact on human reproduction and the health trajectory of offspring. *Adv Exp Med Biol*, 868:23-47.

Gibb Z, Lambourne SR, Aitken RJ. 2014. The paradoxical relationship between stallion fertility and oxidative stress. *Biol Reprod*, 91:77. doi: 10.1095/biolreprod.114.118539.

Gibb Z, Lambourne SR, Quadrelli J, Smith ND, Aitken RJ. 2015. L-carnitine and pyruvate are pro-survival factors during the storage of stallion spermatozoa at room temperature. *Biol Reprod*, 93:104. doi: 10.1095/biolreprod.115.131326.

Gibb Z, Aitken RJ. 2016. The impact of sperm metabolism during in vitro storage: the stallion as a model. *BioMed Res Int*, 2016:9380609. doi.org/10.1155/2016/9380609.

Gravance CG, Garner DL, Baumber J, Ball BA. 2000. Assessment of equine sperm mitochondrial function using JC-1. *Theriogenology*, 53:1691-1703.

Grunewald S, Rasch M, Reinhardt M, Baumann T, Paasch U, Glander HJ. 2008. Stability of fluorochrome based assays to measure subcellular sperm functions. *Asian J Androl*, 10:455-459.

Hoffmann N, Oldenhof H, Morandini C, Rohn K, Sieme H. 2011. Optimal concentrations of cryoprotective agents for semen from stallions that are classified 'good' or 'poor' for freezing. *Anim Reprod Sci*, 125:112-118.

Johannisson A, Morrell JM, Thoren J, Jonsson M, Dalin AM, Rodriguez-Martinez H. 2009. Colloidal centrifugation with Androcoll-E prolongs stallion sperm motility, viability and chromatin integrity. *Anim Reprod Sci*, 116:119-128.

Johnson L, Thompson DL Jr, Varner DD. 2008. Role of Sertoli cell number and function on regulation of spermatogenesis. *Anim Reprod Sci*, 105:23-51.

Love CC, Kenney RM. 1998. The relationship of increased susceptibility of sperm DNA to denaturation and fertility in the stallion. *Theriogenology*, 50:955-972.

Love CC. 2005. The sperm chromatin structure assay: a review of clinical applications. *Anim Reprod Sci*, 89:39-45.

Martin Munoz P, Ortega Ferrusola C, Vizuete G, Plaz Davila M, Rodriguez Martinez H, Pena Vega FJ. 2015. Depletion of intracellular thiols and increased production of 4-hydroxynonenal that occur during cryopreservation of stallion spermatozoa leads to caspase activation, loss of motility, and cell death. *Biol Reprod*, 93:143. doi: 10.1095/biolreprod.115.132878.

Morrell JM, Dalin AM, Rodriguez-Martinez H. 2009a. Comparison of density gradient and single layer centrifugation of stallion spermatozoa: yield, motility and survival. *Equine Vet J*, 41:53-58.

Morrell JM, Johannisson A, Dalin AM, Rodriguez-Martinez H. 2009b. Morphology and chromatin integrity of stallion spermatozoa prepared by density gradient and single layer centrifugation through silica colloids. *Reprod Domest Anim*, 44:512-517.

Oldenhof H, Friedel K, Sieme H, Glasmacher B, Wolkers WF. 2010. Membrane permeability parameters for freezing of stallion sperm as determined by Fourier transform infrared spectroscopy. *Cryobiology*, 61:115-122.

Oldenhof H, Friedel K, Akhoondi M, Gojowsky M, Wolkers WF, Sieme H. 2012. Membrane phase behavior during cooling of stallion sperm and its correlation with freezability. *Mol Membr Biol*, 29:95-106.

Oldenhof H, Gojowsky M, Wang S, Henke S, Yu C, Rohn K, Wolkers WF, Sieme H. 2013. Osmotic stress and membrane phase changes during freezing of stallion sperm: mode of action of cryoprotective agents. *Biol Reprod*, 88:68. doi: 10.1095/biolreprod.112.104661.

Ortega-Ferrusola C, Sotillo-Galan Y, Varela-Fernandez E, Gallardo-Bolanos JM, Muriel A, Gonzalez-Fernandez L, Tapia JA, Peña FJ. 2008. Detection of "apoptosis-like" changes during the cryopreservation process in equine sperm. *J Androl*, 29:213-221.

Ortega-Ferrusola C, Garcia BM, Gallardo-Bolanos JM, Gonzalez-Fernandez L, Rodriguez-Martinez H, Tapia JA, Peña FJ. 2009a. Apoptotic markers can be used to forecast the freezability of stallion spermatozoa. *Anim Reprod Sci*, 114:393-403.

Ortega Ferrusola C, Gonzalez Fernandez L, Macias Garcia B, Salazar-Sandoval C, Morillo Rodriguez A, Rodriguez Martinez H, Tapia JA, Peña FJ. 2009b. Effect of cryopreservation on nitric oxide production by stallion spermatozoa. *Biol Reprod*, 81:1106-1111.

Ortega Ferrusola C, Gonzalez Fernandez L, Morrell JM, Salazar Sandoval C, Macias Garcia B, Rodriguez-Martinez H, Tapia JA, Peña FJ. 2009c. Lipid peroxidation, assessed with BODIPY-C11, increases after cryopreservation of stallion spermatozoa, is stallion-dependent and is related to apoptotic-like changes. *Reproduction*, 138:55-63.

Ortega Ferrusola C, Gonzalez Fernandez L, Salazar Sandoval C, Macias Garcia B, Rodriguez Martinez H, Tapia JA, Peña FJ. 2010. Inhibition of the mitochondrial permeability transition pore reduces "apoptosis like" changes during cryopreservation of stallion spermatozoa. *Theriogenology*, 74:458-465.

Peña FJ, Garcia BM, Samper JC, Aparicio IM, Tapia JA, Ferrusola CO. 2011. Dissecting the molecular damage to stallion spermatozoa: the way to improve current cryopreservation protocols? *Theriogenology*, 76:1177-1186.

Peña FJ, Plaza Davila M, Ball BA, Squires EL, Martin Munoz P, Ortega Ferrusola C, Balao da Silva C. 2015. The impact of reproductive technologies on stallion mitochondrial function. *Reprod Domest Anim*,



50:529-537.

Peña FJ, Ortega Ferrusola C, Martin Munoz P. 2016. New flow cytometry approaches in equine andrology. *Theriogenology*, 86:366-372.

Plaza Davila M, Martin Munoz P, Tapia JA, Ortega Ferrusola C, Balao da Silva CC, Pena FJ. 2015. Inhibition of mitochondrial complex I leads to decreased motility and membrane integrity related to increased hydrogen peroxide and reduced ATP production, while the inhibition of glycolysis has less impact on sperm motility. *PLoS One*, 10:e0138777.

Pommer AC, Rutlant J, Meyers SA. 2002. The role of osmotic resistance on equine spermatozoal function. *Theriogenology*, 58:1373-1384.

Ponthier J, Teague SR, Franck TY, de la Rebiere G, Serteyn DD, Brinsko SP, Love CC, Blanchard TL, Varner DD, Deleuze SC. 2013. Effect of non-sperm cells removal with single-layer colloidal centrifugation on myeloperoxidase concentration in post-thaw equine semen. *Theriogenology*, 80:1082-1087.

Pukazhenthil BS, Johnson A, Guthrie HD, Songsasen N, Padilla LR, Wolfe BA, Coutinho da Silva M, Alvarenga MA, Wildt DE. 2014. Improved sperm cryosurvival in diluents containing amides versus glycerol in the Przewalski's horse (*Equus ferus przewalskii*). *Cryobiology*, 68:205-214.

Purdey MS, Connaughton HS, Whiting S, Schartner EP, Monro TM, Thompson JG, Aitken RJ, Abell AD. 2015. Boronate probes for the detection of hydrogen peroxide release from human spermatozoa. *Free Radic Biol Med*, 81:69-76.

Sabeur K, Ball BA. 2006. Detection of superoxide anion generation by equine spermatozoa. *Am J Vet Res*, 67:701-706.

Shen X, Tian J, Li J, Li X, Chen Y. 1992. Formation of the excited ferryl species following Fenton reaction. *Free Radic Biol Med*, 13:585-592.

Sousa AP, Amaral A, Baptista M, Tavares R, Caballero Campo P, Caballero Peregrin P, Freitas A, Paiva A, Almeida-Santos T, Ramalho-Santos J. 2011. Not all sperm are equal: functional mitochondria characterize a subpopulation of human sperm with better fertilization potential. *PLoS One*, 6:e18112.

Stowe DF, Camara AK. 2009. Mitochondrial reactive oxygen species production in excitable cells: modulators of mitochondrial and cell function. *Antioxid Redox Signal*, 11:1373-1414.

Thomas AD, Meyers SA, Ball BA. 2006. Capacitation-like changes in equine spermatozoa following cryopreservation. *Theriogenology*, 65:1531-1550.

Varner DD. 2008. Developments in stallion semen evaluation. *Theriogenology*, 70:448-462.

Varner DD, Gibb Z, Aitken RJ. 2015. Stallion fertility: a focus on the spermatozoon. *Equine Vet J*, 47:16-24.

Waite JA, Love CC, Brinsko SP, Teague SR, Salazar JL Jr, Mancill SS, Varner DD. 2008. Factors impacting equine sperm recovery rate and quality following cushioned centrifugation. *Theriogenology*, 70:704-714.

Yeste M, Estrada E, Rocha LG, Marin H, Rodriguez-Gil JE, Miro J. 2015. Cryotolerance of stallion spermatozoa is related to ROS production and mitochondrial membrane potential rather than to the integrity of sperm nucleus. *Andrology*, 3:395-407.

Zini A, De Lamirande E, Gagnon C. 1995. Low levels of nitric oxide promote human sperm capacitation in vitro. *J Androl*, 16:424-431.