



Non-additive Lie centralizer of strictly upper triangular matrices

DRISS AIAT HADJ AHMED

Centre Régional des Metiers d'Education et de Formation (CRMEF)
Tangier, Morocco
ait_hadj@yahoo.com

Received December 31, 2018
Accepted February 4, 2019

Presented by Consuelo Martínez

Abstract: Let \mathcal{F} be a field of zero characteristic, let $N_n(\mathcal{F})$ denote the algebra of $n \times n$ strictly upper triangular matrices with entries in \mathcal{F} , and let $f : N_n(\mathcal{F}) \rightarrow N_n(\mathcal{F})$ be a non-additive Lie centralizer of $N_n(\mathcal{F})$, that is, a map satisfying that $f([X, Y]) = [f(X), Y]$ for all $X, Y \in N_n(\mathcal{F})$. We prove that $f(X) = \lambda X + \eta(X)$ where $\lambda \in \mathcal{F}$ and η is a map from $N_n(\mathcal{F})$ into its center $\mathcal{Z}(N_n(\mathcal{F}))$ satisfying that $\eta([X, Y]) = 0$ for every X, Y in $N_n(\mathcal{F})$.

Key words: Lie centralizer, strictly upper triangular matrices, commuting map.

AMS *Subject Class.* (2010): 16S50, 15A27, 16U80, 15B99, 47B47, 16R60.

1. INTRODUCTION

Consider a ring R . An additive mapping $T : R \rightarrow R$ is called a left (respectively right) centralizer if $T(ab) = T(a)b$ (respectively $T(ab) = aT(b)$) for all $a, b \in R$. The map T is called a centralizer if it is a left and a right centralizer. The characterization of centralizers on algebras or rings has been a widely discussed subject in various areas of mathematics.

In [13] Zalar proved the following interesting result: if R is a 2-torsion free semiprime ring and T is an additive mapping such that $T(a^2) = T(a)a$ (or $T(a^2) = aT(a)$), then T is a centralizer. Vukman [12] considered additive maps satisfying similar conditions, namely $2T(a^2) = T(a)a + aT(a)$ for any $a \in R$, and showed that if R is a 2-torsion free semiprime ring then T is also a centralizer. Since then, the centralizers have been intensively investigated by many mathematicians (see, e.g., [3, 4, 5, 6, 8]).

Let R be a ring. An additive map $f : R \rightarrow R$, is called a Lie centralizer of R if

$$f([x, y]) = [f(x), y] \quad \text{for all } x, y \in R, \quad (1.1)$$

where $[x, y]$ is the Lie product of x and y .



Recently, Ghomanjani and Bahmani [9] dealt with the structure of Lie centralizers of trivial extension algebras, whereas Fošner and Jing [7] studied Lie centralizers of triangular rings.

The inspiration of this paper comes from the articles [1, 5, 7] in which the authors deal with the Lie centralizer maps of triangular algebras and rings. In this note we will consider non-additive Lie centralizers on strictly upper triangular matrices over a field of zero characteristic.

Throughout this article, \mathcal{F} is a field of zero characteristic. Let $M_n(\mathcal{F})$ and $N_n(\mathcal{F})$ denote the algebra of all $n \times n$ matrices and the algebra of all $n \times n$ strictly upper triangular matrices over \mathcal{F} , respectively. We use $\text{diag}(a_1, a_2, \dots, a_n)$ to represent a diagonal matrix with diagonal (a_1, a_2, \dots, a_n) where $a_i \in \mathcal{F}$. The set of all $n \times n$ diagonal matrices over \mathcal{F} is denoted by $D_n(\mathcal{F})$. Let I_n be the identity in $M_n(\mathcal{F})$, $J = \sum_{i=1}^{n-1} E_{i,i+1}$ and $\{E_{ij} : 1 \leq i, j \leq n\}$ the canonical basis of $M_n(\mathcal{F})$, where E_{ij} is the matrix with 1 in the (i, j) position and zeros elsewhere. By $C_{N_n(\mathcal{F})}(X)$ we will denote the centralizer of the element X in the ring $N_n(\mathcal{F})$.

The notation $f : N_n(\mathcal{F}) \rightarrow N_n(\mathcal{F})$ means a non-additive map satisfying $f([X, Y]) = [f(X), Y]$ for all $X, Y \in N_n(\mathcal{F})$.

Notice that it is easy to check that $\mathcal{Z}(N_n(\mathcal{F})) = \mathcal{F}E_{1n}$.

The main result in this paper is the following:

THEOREM 1.1. *Let \mathcal{F} be a field of zero characteristic. If $f : N_n(\mathcal{F}) \rightarrow N_n(\mathcal{F})$ is a non-additive Lie centralizer then there exists $\lambda \in \mathcal{F}$ and a map $\eta : N_n(\mathcal{F}) \rightarrow \mathcal{Z}(N_n(\mathcal{F}))$ satisfying $\eta([X, Y]) = 0$ for every X, Y in $N_n(\mathcal{F})$ such that $f(X) = \lambda X + \eta(X)$ for all X in $N_n(\mathcal{F})$.*

Notice that the converse is trivially true: every map $f(X) = \lambda X + \eta(X)$ with η satisfying the condition in Theorem 1.1 is a (non-additive) Lie centralizer.

2. PROOFS

Let's start with some basic properties of Lie centralizers.

LEMMA 2.1. *Let f be a non-additive Lie centralizer of $N_n(\mathcal{F})$. Then:*

- (1) $f(0) = 0$;
- (2) for every $X, Y \in N_n(\mathcal{F})$, we have $f([X, Y]) = [X, f(Y)]$;

(3) f is a commuting map, i.e., $f(X)X = Xf(X)$ for all $X \in N_n(\mathcal{F})$.

Proof. To prove (1) it suffices to notice that

$$f(0) = f([0, 0]) = [f(0), 0] = 0.$$

(2) Observe that if $f([X, Y]) = [f(X), Y]$, then we have

$$f(XY - YX) = f(X)Y - Yf(X).$$

Interchanging X and Y in the above identity, we have

$$f(YX - XY) = f(Y)X - Xf(Y).$$

Replacing X with $-X$ in the above relation, we arrive at $f(XY - YX) = Xf(Y) - f(Y)X$ which can be written as $f([X, Y]) = [X, f(Y)]$.

From (1) one also gets (3):

$$[f(X), X] = f([X, X]) = f(0) = 0. \quad \blacksquare$$

Remark 2.1. Let f be a non-additive Lie centralizer of $N_n(\mathcal{F})$ and $X \in C_{N_n(\mathcal{F})}(Y)$. Then $f(X) \in C_{N_n(\mathcal{F})}(Y)$. Indeed, if $X \in C_{N_n(\mathcal{F})}(Y)$, then $[X, Y] = 0$ and

$$0 = f(0) = f([X, Y]) = [f(X), Y].$$

LEMMA 2.2. Let f be a non-additive Lie centralizer of $N_n(\mathcal{F})$. Then:

- (1) $f\left(\sum_{i=1}^{n-1} a_i E_{i,i+1}\right) = \sum_{i=1}^{n-1} b_i E_{i,i+1}$;
- (2) there exists $\lambda \in \mathcal{F}$ such that $f(J) = \lambda J$.

Proof. Let $D_0 = \sum_{i=1}^n (n-i) E_{i,i}$.

(1) Consider $A \in M_n(\mathcal{F})$. It is well known that $[D_0, A] = A$ if and only if $A = \sum_{i=1}^{n-1} a_i E_{i,i+1}$.

Hence, if $A = \sum_{i=1}^{n-1} a_i E_{i,i+1}$, we have $[D_0, A] = A$. Thus $f([D_0, A]) = [D_0, f(A)] = f(A)$. Therefore $f(A) = \sum_{i=1}^{n-1} b_i E_{i,i+1}$.

(2) As in (1), consider $A = \sum_{i=1}^{n-1} a_i E_{i,i+1}$ for some $a_i \in \mathcal{F}$. Then $[J, A] = 0$ if and only if $A = aJ$ for some $a \in \mathcal{F}$.

Indeed, $f(J) = \sum_{i=1}^{n-1} a_i E_{i,i+1}$ by (1). Thus, $0 = f(0) = f([J, J]) = [J, f(J)]$. Hence, there exists $\lambda \in \mathcal{F}$ such that $f(J) = \lambda J$. ■

We will need the following lemma.

LEMMA 2.3. (LEMMA 2.1, [14]) *Suppose that \mathcal{F} is an arbitrary field. If $G, H \in UT_n(\mathcal{F})$ are such that $g_{i,i+1} = h_{i,i+1} \neq 0$ for all $1 \leq i \leq n-1$, then G and H are conjugated in $UT_n(\mathcal{F})$.*

Here $UT_n(\mathcal{F})$ is the multiplicative group of $n \times n$ upper triangular matrices with only 1's in the main diagonal. From the lemma above we obtain the following corollary.

COROLLARY 2.1. *Let \mathcal{F} be a field. For every $A = \sum_{1 \leq i < j \leq n} a_{ij} E_{ij}$, where $a_{i,i+1} \neq 0$ for all $1 \leq i \leq n-1$, there exists $B \in T_n(\mathcal{F})$ such that $B^{-1}AB = J$ and $T_n(\mathcal{F})$ is the ring of upper triangular matrices.*

Proof. Let A be a matrix in $N_n(\mathcal{F})$ of the mentioned form. Then $I_n + A$ is a unitriangular matrix. Let's notice first that there exists $B_1 \in D_n(\mathcal{F})$ such that $(B_1^{-1}AB_1)_{i,i+1} = 1$ for all $i \in \mathbb{N}$. We can construct $B_1 \in D_n(\mathcal{F})$ recursively by:

$$(B_1)_{11} = 1, \quad (B_1)_{i+1,i+1} = (B_1)_{ii} \cdot (A_{i,i+1})^{-1} \quad \text{for } i \geq 1.$$

Consider the matrix $I_n + B_1^{-1}AB \in UT_n(\mathcal{F})$. The unitriangular matrices $I_n + J$ and $I_n + B_1^{-1}AB$ fulfill the condition in Lemma 2.3. Hence, there exists $B_2 \in UT_n(\mathcal{F})$ such that

$$I_n + J = B_2^{-1}(I_n + B_1^{-1}AB)B_2.$$

Then $J = B_2^{-1}(B_1^{-1}AB)B_2$. Taking $B = B_1B_2 \in T_n(\mathcal{F})$, we get $J = B^{-1}AB$ as wanted. ■

LEMMA 2.4. *Let $A = \sum_{i < j} a_{ij} E_{ij}$ be a matrix in $N_n(\mathcal{F})$ with $a_{i,i+1} \neq 0$ for every $i = 1, \dots, n-1$. Then there exists $\lambda_A \in \mathcal{F}$ such that $f(A) = \lambda_A A$.*

Proof. Since $A = \sum_{1 \leq i < j \leq n} a_{ij} E_{ij}$, where $a_{i,i+1} \neq 0$, there exists $T \in T_n(\mathcal{F})$ such that $TAT^{-1} = J$ by the previous corollary. Define $h : N_n(\mathcal{F}) \rightarrow N_n(\mathcal{F})$ by $h(X) = Tf(T^{-1}XT)T^{-1}$. Then h is a non-additive Lie centralizer. Indeed, for all $A, B \in N_n(\mathcal{F})$ we have:

$$\begin{aligned}
h([A, B]) &= Tf(T^{-1}[A, B]T)T^{-1} \\
&= Tf(T^{-1}(AB - BA)T)T^{-1} \\
&= Tf(T^{-1}ATT^{-1}BT - T^{-1}BTT^{-1}AT)T^{-1} \\
&= Tf([T^{-1}AT, T^{-1}BT])T^{-1} \\
&= T[f(T^{-1}AT), T^{-1}BT]T^{-1} \\
&= T(f(T^{-1}AT)T^{-1}BT - T^{-1}BTf(T^{-1}AT))T^{-1} \\
&= Tf(T^{-1}AT)T^{-1}B - BTf(T^{-1}AT)T^{-1} \\
&= [Tf(T^{-1}AT)T^{-1}, B] \\
&= [h(A), B].
\end{aligned}$$

Hence, $h(J) = \lambda_A J$ by Lemma 2.2. Then

$$Tf(A)T^{-1} = Tf(T^{-1}(TAT^{-1})T)T^{-1} = h(J) = \lambda_A J = \lambda_A TAT^{-1}.$$

Multiplying the left and right sides by T^{-1} and T respectively yields $f(A) = \lambda_A A$. ■

Now we wish to extend Lemma 2.4 to all elements of $N_n(\mathcal{F})$. In order to do this, let's introduce the following set:

$$\mathcal{S} = \{B = (b_{ij}) \in N_n(\mathcal{F}) : b_{i,i+1} \neq 0 \quad \forall i = 1, \dots, n-1\}.$$

This set has an important property that is established below.

LEMMA 2.5. *Let \mathcal{F} be a field. Every element of $N_n(\mathcal{F})$ can be written as a sum of at most two elements of \mathcal{S} .*

Proof. If $a_{i,i+1} \neq 0$ for all $i = 1, \dots, n-1$, then A belongs to \mathcal{S} , so there is nothing to prove. If A is not in \mathcal{S} , then we can define B_1 and B_2 as follows:

$$(B_1)_{ij} = \begin{cases} a_{i,i+1} - b_i & \text{if } j = i+1, \\ a_{ij} & \text{if } j > i+1, \end{cases} \quad (B_2)_{ij} = \begin{cases} b_i & \text{if } j = i+1, \\ 0 & \text{otherwise,} \end{cases}$$

where b_i is an element in \mathcal{F} different from $a_{i,i+1}$. It is easy to see that B_1, B_2 are in \mathcal{S} , and $A = B_1 + B_2$, so we wanted. ■

LEMMA 2.6. *Let \mathcal{F} be a field. For arbitrary elements A, B of $N_n(\mathcal{F})$, there exists $\lambda_{A,B} \in \mathcal{F}$ such that*

$$f(A + B) = f(A) + f(B) + \lambda_{A,B}E_{1n}.$$

Proof. For any A, B, X of $N_n(\mathcal{F})$, we have

$$\begin{aligned} [f(A + B), X] &= f([A + B, X]) \\ &= [A + B, f(X)] \\ &= [A, f(X)] + [B, f(X)] \\ &= [f(A), X] + [f(B), X] \\ &= [f(A) + f(B), X], \end{aligned}$$

which implies that $f(A + B) - f(A) - f(B) \in \mathcal{Z}(N_n(\mathcal{F}))$. Thus, there exists $\lambda_{A,B} \in \mathcal{F}$ such that $f(A + B) = f(A) + f(B) + \lambda_{A,B}E_{1n}$. ■

Now we can prove the main theorem.

Proof of Theorem 1.1. Let $A, B \in \mathcal{S}$ be two non-commuting elements. By Lemma 2.4, $f(A) = \lambda_A A$, $f(B) = \lambda_B B$, $\lambda_A, \lambda_B \in \mathcal{F}$.

Since f is a non-additive Lie centralizer, we get,

$$\begin{aligned} f([A, B]) &= [f(A), B] = \lambda_A[A, B] \\ &= [A, f(B)] = \lambda_B[A, B]. \end{aligned}$$

Then, $[A, B] \neq 0$ implies that $\lambda_A = \lambda_B$.

If $A, B \in \mathcal{S}$ commute, then we take $C \in \mathcal{S}$ that does not commute neither with A nor with B . As we have just seen, $\lambda_A = \lambda_C$ and $\lambda_B = \lambda_C$. So $\lambda_A = \lambda_B = \lambda$ for arbitrary elements $A, B \in \mathcal{S}$. Given $X \in N_n(\mathcal{F})$ we know, by Lemma 2.5, that there exists $A, B \in \mathcal{S}$ such that $X = A + B$ (we can assume that $X \notin \mathcal{S}$). Then $f(X) - f(A) - f(B) \in \mathcal{Z}(N_n(\mathcal{F}))$ by Lemma 2.6.

That is $f(X) - \lambda_A A - \lambda_B B = f(X) - \lambda X \in \mathcal{Z}(N_n(\mathcal{F}))$ for $\lambda \in \mathcal{F}$ such that $f(A) = \lambda A$ for each $A \in \mathcal{S}$.

We can define $\eta : N_n(\mathcal{F}) \rightarrow \mathcal{Z}(N_n(\mathcal{F}))$ such that $\eta(X) = f(X) - \lambda X$, that is, $f(X) = \lambda X + \eta(X)$.

Notice that $\eta(A) = 0$ for each $A \in \mathcal{S}$. Furthermore, if $X, Y \in N_n(\mathcal{F})$, then

$$\begin{aligned} f([X, Y]) &= \lambda[X, Y] + \eta([X, Y]) = [f(X), Y] \\ &= [\lambda X + \eta(X), Y] = \lambda[X, Y], \end{aligned}$$

since $\eta(X) \in \mathcal{Z}(N_n(\mathcal{F}))$.

Consequently, $\eta([X, Y]) = 0$ and Theorem 1.1 is proved. ■

ACKNOWLEDGEMENTS

The author would like to thank the referee for providing useful suggestions which served to improve this paper.

REFERENCES

- [1] J. BOUNDS, Commuting maps over the ring of strictly upper triangular matrices, *Linear Algebra Appl.* **507** (2016), 132–136.
- [2] M. BREŠAR, Centralizing mappings on von Neumann algebra, *Proc. Amer. Math. Soc.* **111** (1991), 501–510.
- [3] M. BREŠAR, Centralizing mappings and derivations in prime rings, *J. Algebra* **156** (1993), 385–394.
- [4] M. BREŠAR, Commuting traces of biadditive mappings, commutativity-preserving mappings and Lie mappings, *Trans. Amer. Math. Soc.* **335** (1993), 525–546.
- [5] W.-S. CHEUNG, Commuting maps of triangular algebras, *J. London Math. Soc. (2)* **63** (2001), 117–127.
- [6] D. EREMITA, Commuting traces of upper triangular matrix rings, *Aequationes Math.* **91** (2017), 563–578.
- [7] A. FOŠNER, W. JING, Lie centralizers on triangular rings and nest algebras, *Adv. Oper. Theory* **4** (2) (2019), 342–350.
- [8] W. FRANCA, Commuting maps on some subsets of matrices that are not closed under addition, *Linear Algebra Appl.* **437** (2012), 388–391.
- [9] F. GHOMANJANI, M.A. BAHMANI, A note on Lie centralizer maps, *Palest. J. Math.* **7** (2) (2018), 468–471.
- [10] T.K. LEE, Derivations and centralizing mappings in prime rings, *Taiwanese J. Math.* **1** (3) (1997), 333–342.
- [11] T.K. LEE, T.C. LEE, Commuting additive mappings in semiprime rings, *Bull. Inst. Math. Acad. Sinica* **24** (1996), 259–268.
- [12] J. VUKMAN, An identity related to centralizers in semiprime rings, *Comment. Math. Univ. Carolin.* **40** (3) (1999), 447–456.
- [13] B. ZALAR, On centralizers of semiprime rings, *Comment. Math. Univ. Carolin.* **32** (4) (1991), 609–614.
- [14] R. SLOWIK, Expressing infinite matrices as products of involutions, *Linear Algebra Appl.* **438** (2013), 399–404.
- [15] L. CHEN, J.H. ZHANG, Nonlinear Lie derivations on upper triangular matrices, *Linear Multilinear Algebra* **56** (2008), 725–730.
- [16] D. AIAT HADJ AHMED, R. SLOWIK, m -Commuting maps of the rings of infinite triangular and strictly triangular matrices (*in preparation*).